

Gut microbiota and attention deficit hyperactivity disorder: new perspectives for a challenging condition

María Carmen Cenit^{1,2} · Isabel Campillo Nuevo¹ · Pilar Codoñer-Franch^{2,3} · Timothy G. Dinan^{4,5} · Yolanda Sanz¹

Received: 4 October 2016 / Accepted: 22 February 2017
© Springer-Verlag Berlin Heidelberg 2017

Abstract A bidirectional communication between the gut and the brain (gut–brain axis) is well recognized with the gut microbiota viewed as a key regulator of this cross-talk. Currently, a body of preclinical and to a lesser extent epidemiological evidence supports the notion that host–microbe interactions play a key role in brain development and function and in the etiology of neurodevelopmental disorders. Early life events and shifts away from traditional lifestyles are known to impact gut microbiota composition and function and, thereby, may increase the risk of developing neurodevelopmental disorders. Attention deficit hyperactivity disorder (ADHD) is nowadays the most prevalent neurodevelopmental disorder. Despite many years of research its

etiology is unclear and its diagnosis and treatment are still challenging. Different factors reported to be associated with the risk of developing ADHD and/or linked to different ADHD manifestations have also been linked to shifts in gut microbiota composition, suggesting a link between the microbiota and the disorder. Evidence from preliminary human studies also suggests that dietary components that modulate gut microbiota may also influence ADHD development or symptoms, although further studies are warranted to confirm this hypothesis. Here, we firstly review the potential mechanisms by which the gut microbiota may regulate the brain–gut axis and influence behavior and neurodevelopmental disorders. Secondly, we discuss the current knowledge about the different factors and dietary components reported to be associated with the risk of developing ADHD or its manifestations and with shifts in gut microbiota composition. Finally, we briefly highlight the need to progress our understanding regarding the role of the gut microbiota in ADHD, since this could open new avenues for early intervention and improved management of the disease.

This article is part of the Focused Issue “The role of nutrition in child and adolescent onset mental disorders” of European Child and Adolescent Psychiatry.

✉ María Carmen Cenit
mccenit81@gmail.com

✉ Yolanda Sanz
yolsanz@iata.csic.es

¹ Microbial Ecology, Nutrition and Health Research Group, Institute of Agrochemistry and Food Technology, National Research Council (IATA-CSIC), Avd. Agustín Escardino, 7, 46980 Paterna, Valencia, Spain

² Department of Pediatrics, Dr. Peset University Hospital, Avd. De Gaspar Aguilar, 80, 46017 Valencia, Spain

³ Department of Pediatrics, Obstetrics and Gynecology, University of Valencia, Av Blasco Ibáñez, 13, 46010 Valencia, Spain

⁴ APC Microbiome Institute, University College Cork, Cork, Ireland

⁵ Department of Psychiatry and Neurobehavioural Science, University College Cork, Cork, Ireland

Keywords Microbiota · ADHD · Microbiota · Gut–brain axis · Dysbiosis

Introduction

Neurodevelopmental disorders constitute a heterogeneous group of conditions characterized by developmental deficits of the central nervous system that typically have an early onset. The cluster of neurodevelopmental conditions includes highly prevalent disorders such as autism, autism spectrum disorders, attention deficit hyperactivity disorder (ADHD) and schizophrenia among others.

These disorders, varying from mild to severe phenotypes, lead to impairments in personal, social, academic and/or occupational functioning. The specific underlying mechanisms that bring about the development of the majority of neurodevelopmental disorders have not been fully elucidated. Although genetics has been described to play the most important role in the etiology of most of the neurodevelopmental conditions, environmental factors are also strongly believed to contribute to their development and/or manifestation [1]. ADHD is currently the most frequent neurodevelopmental and neurobehavioral disorder in childhood affecting roughly 5% of children worldwide. ADHD is characterized by developmentally inappropriate levels of hyperactivity, impulsivity and/or attention problems. The current global burden of ADHD in the society is highly significant and its prevalence is alarmingly rising over time [2]. Furthermore, ADHD is persisting in many cases into adolescence and adulthood being thus considered a life-long issue for many patients [3]. According to the fifth edition of the Diagnostic and Statistical Manual (DSM-V) based on the types of symptoms, mainly three types of ADHD presentations can occur; a hyperactive and/or impulsive subtype predominant in males, a inattentive subtype predominant in females and a combined subtype [4]. In addition, ADHD is often associated with other psychiatric comorbidities, being therefore considered as a very heterogeneous disorder [5]. It could be suggested that the effect of different combinations of genetic, environmental factors and complex gene–environment interactions may lead to divergent neurobiological alterations and in turn to different neuropsychological impairments and symptom profiles in ADHD. Thus, it has been proposed that the prominent clinical heterogeneity observed between the different subgroups of ADHD patients might be partially due to differences in the disease etiology [6].

Currently, ADHD diagnosis is based on clinical symptoms, an approach widely criticized on the grounds of reliability. In addition, ADHD treatment is currently based on a multimodal approach with the combination of psychotherapy and pharmacotherapy, but no reliable markers of treatment response have been identified yet and 20–35% of subjects in clinical trials may have an inadequate response to the treatment [7]. Furthermore, pharmacotherapy is often associated with adverse side effects [8, 9] and has shown short-term effectiveness on ADHD treatment although its long-term efficacy is still questionable. Due to these concerns, many families choose not to use pharmacotherapies to treat ADHD, and even when pharmacotherapy is used, drug holidays are recommended. Therefore, future studies would be required to identify risk factors involved in ADHD which could offer new opportunities to improve ADHD diagnosis and management.

So far, family studies, including adoption studies, have suggested a high genetic component for ADHD, showing levels of heritability comparable to other neurodevelopmental and psychiatric disorders such as schizophrenia or bipolar disorder (2- to 8-fold increased risk of ADHD in parents and siblings of children with the disease and rates of ADHD significantly higher between biological relatives compared with adoptive relatives) [10]. Twins studies have confirmed the high level of ADHD heritability ($h^2 = 0.77\text{--}0.82$), but they have also shown a 20% of discordance between monozygotic twins which indicates that environmental factors are also involved in disease development. Indeed, the contribution of environmental factors to ADHD has been estimated to be 20–30% although the exact etiological factors involved remain unknown [11]. Neurotransmission system dysfunctions (dopamine, noradrenaline and/or serotonin deficits) play an important role in the ADHD pathophysiology and its manifestations and currently the most widely used treatments for ADHD are based on targeting monoaminergic systems [12]. However, the precise underlying mechanisms that lead to these neurobiological alterations still remain unidentified. Intensive investigations by different genetic approaches carried out in recent years have had limited success in identifying associations at the critical significance level, very likely due to the high heterogeneity; a large part of ADHD heritability has yet to be defined [13]. Remarkably, most of the studies performed until now in order to decipher the etiology of ADHD have focused on the study of the influence of the human genome in the risk of developing ADHD and/or their phenotypes, overlooking the fact that our second genome, the genome of our microbiota, is also involved in brain development and function and may play a crucial role in neurotransmission and neuronal plasticity at least according to experimental evidence from animal models [14].

It is well established that the gut microbiome, the entire genome of the gut microbiota, influences host development and physiology, playing a key role in the balance between health and disease. In recent years, emerging evidence has suggested a role of the gut microbiota in brain function and behavior and, in turn, intestinal dysbiosis has been linked to different behavioral features [14]. Gut microbiota undergoes a dynamic nonrandom process of maturation until the age of 2–3 years when an adult-like microbiota structure is established. Although later in life the microbiota may still experience changes, the symbiotic link between the host and the microbiota is thought to be mainly established early in life and occurs in parallel with neurodevelopment. In fact, different evidences have demonstrated that early life perturbations of the developing gut microbiota, very vulnerable because of its high instability and immaturity, can impact neurodevelopment and potentially lead to adverse

mental health outcomes later in life [15]. Understanding the early interaction between the intestinal microbiota, the environment and the host would open new avenues for nutritional/therapeutic interventions in at-risk populations for neurodevelopmental disorders. Several preclinical studies and a few human studies have investigated the role of different probiotics in behaviors associated with psychiatric disorders with promising results. Such bacteria are termed psychobiotics (probiotics with a mental health benefit). The potential regulation of the brain–gut axis via intervention in the gut ecosystem [16] has been neglected for many years in neuropsychiatric research, but now it is being intensively explored. Nevertheless, most of the current evidence comes from animal studies and more human studies are urgently needed to move forward in this field. In particular, genomic studies in ADHD including analysis of the gut microbiome are needed, especially at early stages in life and during further neurodevelopmental periods (childhood and adolescence), where many psychiatric disorders first become evident and when the role of the microbiota and its functions could be amenable [17].

In this review, we discuss the potential mechanisms by which the gut microbiota may influence the brain–gut axis and consequently behavior and neurodevelopmental disorders such as ADHD. We also highlight the current knowledge about the different environmental factors reported to be associated with both the risk of developing ADHD or its manifestations and shifts in gut microbiota composition. The ultimate aim of the review is to identify gaps of knowledge that will be necessary to investigate for getting a better understanding of the potential role played by the gut microbiota in ADHD, since this could open new possibilities for early intervention and improved management of the disorder.

Gut microbiota mechanisms regulating gut–brain axis and behavior

Studies in animal models, where the intestinal microbiota can be easily manipulated, have provided multiple insights into how the microbiota may be involved in the development of brain-related disorders [18, 19]. It has also been established that behavioral traits can be transferred by fecal microbiota transplantation, suggesting that microbiota changes could be rather a cause than a consequence of behavioral alterations [20, 21]. Particularly, in the context of mental health, the proposed mechanisms by which the gut microbiota may modulate brain development, function and behavior include immune (cytokines), metabolic (short-chain fatty acids) endocrine (cortisol) and neural (vagus and enteric nervous system) pathways [22]. Microbiota has a substantial impact on the host metabolome and

a large array of essential molecules with neuroactive functions is known to be produced by gut microbes [23]. Different bacterial strains have been shown to mediate behavioral effects via the vagus nerve as demonstrated by vagotomy in a number of preclinical studies, although not all effects depend on the vagus nerve [24]. Likewise, different mechanisms by which the brain influences gut microbiota composition have been identified [16]. Indeed, host stress hormones such as noradrenaline might influence bacterial gene expression or signaling between bacteria, and this might change the microbial composition and activity [25]. Certain bacteria including inhabitants of the intestinal tract are capable of producing different neuroactive compounds such as neurotransmitters. It has been determined that *Lactobacillus* spp. and *Bifidobacterium* spp. can produce GABA [26]; *Escherichia* spp., *Bacillus* spp. and *Saccharomyces* spp. can produce noradrenalin; *Candida* spp., *Streptococcus* spp., *Escherichia* spp. and *Enterococcus* spp. produce serotonin; *Bacillus* spp. produce dopamine; and *Lactobacillus* spp. produce acetylcholine [27]. In addition, plasma serotonin levels of conventional mice have been found to be significantly higher than those in germ-free (GF) mice, which have no intestinal microbiota, demonstrating the capacity of the microbiota to influence this neurotransmitter. It has been suggested that secreted neurotransmitters from bacteria in the intestinal lumen may induce epithelial cells to release molecules such as hormones and cytokines that, in turn, have the ability to modulate neural signaling within the enteric nervous system and subsequently control brain function and behavior [18]. This evidence suggests that the gut microbiota might be a therapeutic target for brain disorders mediated by deregulation in neurotransmitters such as ADHD. The dopamine and norepinephrine pathways, which project to the prefrontal cortex and striatum, are responsible for modulating cognitive control of behavior, motivation and reward perception and these pathways are known to play a central role in the pathophysiology of ADHD. However, no studies have been conducted so far to explore how these pathways may be modulated by the gut microbiota in the context of ADHD and further investigations are needed to develop and validate evidence-based therapeutic strategies targeting the microbiota for ADHD.

Tryptophan is an essential amino acid precursor to many biologically active molecules, including the neurotransmitter serotonin and metabolites of the kynurenine pathway. Only around 5% of systemic tryptophan is metabolized into serotonin and the rest is metabolized along the kynurenine pathway. This depends on the expression of two enzymes, indoleamine-2,3-dioxygenase (IDO), which is found in all tissues in mammals and in yeast, and tryptophan-2,3-dioxygenase (TDO), which is localized within the liver in mammals and is also produced by eukaryote and

prokaryote organisms [28]. The activity of both enzymes is strongly controlled by inflammatory mediators such as cytokines and corticosteroids [29]. The kynurenine pathway is stimulated both in the periphery and in the brain under inflammatory conditions. The increased activation of these two enzymes could induce serotonin depletion and depressive mood. On the other hand, the downstream metabolites of the kynurenine pathway are neuroactive metabolites which can also modulate neurotransmission and immunity [30]. Kynurenine, kynurenic acid, xanthurenic acid, and quinolinic acid are the major downstream breakdown products of tryptophan metabolism which increase after immune stimulation. Those different metabolites have shown to play multiple and diverse effects. In this context, few years ago a study specifically showed that kynurenine, kynurenic acid, and xanthurenic acid have anti-inflammatory effects through a reduction of IFN γ , whereas quinolinic acid, produced mainly by activated microglia and macrophages, exerts pro-inflammatory effects further aggravating the initial inflammation (21). It is important to highlight that quinolinic acid also acts as an important neurotoxin, gliotoxin and pro-oxidant molecule [31]. Gut microbial colonization and development of the gut microbiome occur in parallel with cognitive development and overlap with the ontogeny of the serotonergic system [32]. It has been recently reported that the gut microbiota regulates the host immune system and modulates HPA axis and, therefore, glucocorticoids production, with a subsequent impact on tryptophan metabolism leading to changes in the serotonergic neurotransmission with neuropsychiatric consequences [33]. Of note, all bifidobacteria have the machinery to produce tryptophan and particularly oral ingestion of a strain of *Bifidobacterium infantis* showed to increase, as expected, the levels of the serotonin precursor, tryptophan, in the plasma of rats and potential antidepressant properties measured by the forced swim test [34]. Although a relationship between serotonin depletion and tryptophan availability has not been found in ADHD yet, further studies are awaited to explore the role of the microbiota in the balance between these neural mediators.

In addition, short-chain fatty acids (SCFAs) such as butyrate, acetate and propionate that are produced by microbial fermentation mainly from dietary fiber in the gut are known to have neuroactive properties. For instance, the administration of a high dose of propionate in rats induced a neuroinflammatory process and behavioral alterations related with neurodevelopmental disorders [35]. Propionate is also a common preservative in food products that has been demonstrated to exacerbate autism spectrum disorder symptomatology [36]. On the other hand, butyrate, via an epigenetic mechanism related to its capacity to inhibit histone deacetylase, exerts beneficial behavioral effects [37]. SCFAs also regulate immunity and this may

have consequences on the central nervous system. In fact, it has been recently demonstrated that the treatment with a mix of SCFAs could rescue microglial function previously impaired in GF animals [38].

The microbiota has also been shown to modulate a range of neurotrophins and proteins involved in brain development and plasticity. Particularly, brain-derived neurotrophic factor (BDNF) is a neurotrophin known for its important effects on promoting neurogenesis and neuronal survival. Some findings clearly indicate that the gut microbiota can influence BDNF levels mainly through the effect of SCFAs and this links the microbiota to neuroplasticity-related CNS systems [24]. A reduction in the level of BDNF expression in the hippocampus of germ-free mice, as well as in mice after the administration of oral antimicrobials [24], has been shown to contribute to working-memory impairment. It has also been demonstrated that the intervention with a strain of *Bifidobacterium longum* normalizes the changes in behavior and BDNF mRNA induced by an intestinal parasite. Recent studies have shown that plasma BDNF levels in ADHD patients differ from normal controls suggesting that BDNF may play a role in its pathogenesis [39]. Although it is still not clear the extent to which the human genome and microbiome contribute to BDNF expression, there are indications that both could influence its variation. Therefore, further studies would be desired to explore whether gut microbiota could be manipulated to favorably modulate BDNF levels in the context of ADHD.

Environmental factors influencing gut microbiome composition and risk of developing ADHD

The etiology and reasons for the increasing ADHD prevalence are not fully understood, although recent evidence suggests that environmental factors might be playing a key role [11]. Recently, the risk of developing ADHD has been suggested to be associated with many perinatal risk factors, including delivery mode (vaginal delivery or cesarean section delivery), gestational age, type of feeding (breastfeeding and formula), maternal health and early life stressors among others, supporting the idea that ADHD is a multifactorial disorder triggered by certain environmental factors such as gut microbiota among others in genetically susceptible individuals [40, 41]. Of note, many of those environmental factors associated with the risk of developing ADHD are also known to influence the early gut microbiota composition (Fig. 1) [42, 43]. The perinatal period is a key period in the establishment of the initial microbiota of the infant and it has been considered as a window of opportunity for influencing brain development via gut microbiota modulation [15].




Fig. 1 Schematic representation of environmental factors that influence the development of gut microbiota in childhood. Many of these factors have also been associated with the risk of developing ADHD, suggesting a link between the microbiota and this disorder. There-

fore, it is hypothesized that microbiota configuration during infancy and childhood, under the influence of different environmental factors, may have long-term consequences, protecting against or contributing to ADHD development or its manifestation

Delivery mode and gestational age

The mode of delivery (vaginal delivery or delivery by cesarean section) has a strong influence on shaping the initial gut microbiota composition. Cesarean section (C-section) correlates with decreased gut microbiota diversity and delays in colonization of *Bacteroidetes*, *Bifidobacterium* and *Lactobacillus* [44, 45]. In fact, the analysis of the meconium of newborn infants revealed a strong correlation between the first infant gut microbial communities and those found in either the mother's vagina in the case of vaginal delivery or the mother's skin in the case of delivery by CS [46].

Interestingly, using animal models delivery by CS has been shown to be associated with changes in dopamine response [47]. Furthermore, CS delivery compared to vaginal delivery has been linked to the risk of developing many diseases, including neurodevelopmental disorders such as autism, although confounding factors may play a role driven this association [48]. In addition, although a recent meta-analysis have not confirmed the significant

effect of CS on the risk of developing ADHD [49], some individual studies have reported a link between developing the disorder and this type of delivery mode [50, 51]. Therefore, there is still a need for further studies considering confounding factors to draw more definitive conclusions about the relationship between risk of ADHD and delivery mode. In addition, significant divergences in gut microbial composition of the neonate have been found depending on the duration of pregnancy, being gestational age at birth a major driver of gut microbiota assembly in preterm neonates. Preterm neonates, who have been described to have a higher risk of developing ADHD and to have significantly more severe ADHD symptoms [52, 53], seem to lack two of the main bacterial genera present in healthy term infants, *Bifidobacterium* and *Lactobacillus* [43]. Preterm birth increases also the risk of infection which in turn is associated with the increased risk of neurodevelopmental impairment. In addition, it is well established that antibiotics use, which is related to infection, promotes a less diverse microbiota [54] and antibiotic intake and altered brain function are also associated [55].

Maternal health and early life stressors

Maternal stress during pregnancy has also been demonstrated to have profound effects on offspring, being associated with an increased incidence of neurodevelopmental disorders including ADHD [56–58]. Furthermore, children with ADHD whose mothers were exposed to moderate and severe stress during pregnancy were reported to develop more severe symptoms than children with ADHD whose mothers were not exposed to prenatal stress [58]. The exact mechanisms by which maternal stress causes developmental problems in offspring are still unclear although it is well known that chronic stress during pregnancy alters vaginal host immunity and resident microbiota composition [59] and this, in turn, could impact the microbial assembly of the neonatal gut having significant consequences for offspring development and disease risk. Maternal separation, an early life stressor that results in long-term hypothalamic–pituitary–adrenal (HPA) alterations, has been shown to have long-term effects on the microbiome composition and behavior in animal models (45). It has been hypothesized that non-genetic factors, such as early life experiences, may interact with genetic predisposition, leading to ADHD development. The development of the HPA stress axis overlaps with the early postnatal microbial colonization of the neonatal gut and the establishment of the microbiota and has been demonstrated to be sensitive to gut microbial disruption in animal models. In particular, Sudo and colleagues published that gut microbes were able to affect the postnatal development of the HPA stress response in mice, being the gut commensal microbiota required at a specific developmental stage for the HPA system to become completely susceptible to inhibitory neural regulation [60]. Glucocorticoids, the end product of HPA axis activation, are required for normal brain development. The appropriate development of the HPA axis is essential for the balanced functioning of the gut–brain axis. HPA dysregulation has been recognized as a hallmark of inflammatory and psychiatric disorders. A meta-analysis of case–control studies found significantly lower cortisol levels in children suffering from ADHD indicating a dysregulation of the HPA axis in the disorder (46). Contradictory findings have also been published. In the spontaneously hypertensive rat, an animal model of ADHD, maternal separation induced excessive activation of the HPA-mediated stress response and also affects dopamine transporter function and norepinephrine release in the hippocampus [61, 62]. It has been reported that the administration of specific strains of *Lactobacillus* and *Bifidobacterium* species seems to be able to normalize corticosterone release as well as to ameliorate colonic dysfunction and

behavioural alterations induced by maternal separation, supporting the beneficial role of interventions in the gut ecosystem in restoring HPA-axis balance, at least in this model of chronic stress [60, 63].

Maternal infection, strongly linked to microbiota changes and immune activation, has also been suggested to be associated with ADHD onset in offspring [64, 65]. A preclinical study assessing the effect of maternal infection on mental health in the offspring determined that maternal immune activation could lead to a disrupted gut microbiota as well as to gut barrier dysfunction and behavioral alterations in the offspring [66]. The integrity of the adaptive immune system, and in particular T lymphocyte responses, has been shown to be crucial for normal learning and memory in mice [67]. Ceylan and co-workers found significantly higher levels of neopterin, a good indicator of cellular immunity, in ADHD compared with controls, suggesting that changes in cellular immunity may be involved in ADHD brain dysfunction [68]. Remarkably, the significant impact of the prenatal immune activation on the gut–brain axis has been shown to be reversible by the administration of *Bacteroides fragilis*, a common inhabitant of gut microbiota, to the offspring [66].

It has been demonstrated that maternal obesity and metabolic disease, including diabetes and hypertension, and unhealthy maternal diets have a long-term impact on offspring behavior and physiology [69], which may also be related to changes in gut microbiota composition induced by these conditions and the diet reported elsewhere. Different studies have suggested that maternal body mass index (BMI) and obesity prior and during pregnancy are linked to the risk of developing ADHD [70]. Rodriguez and colleagues analyzed data from three separate cohorts which examined whether pregnancy weight (BMI and/or weight gain) shows a relationship to symptoms of ADHD in offspring [71]. They found a significant association between pre-pregnancy overweight or obesity and a high ADHD symptom score in offspring while gestational weight gain was unrelated to ADHD. Evidence from animal models supports human epidemiological studies and demonstrates that maternal high-fat-diet (HFD)-induced obesity impacts behavioral programming of offspring resulting in impairments in social behavior and hyperactivity, increased anxiety and depressive behaviors and reduced cognitive development [72]. Remarkably, a recent study in mice has shown that dietary intervention can rescue maternal obesity induced behavioral deficits and neuroinflammation in offspring [72]. However, further studies in humans should be conducted to investigate the role of dietary intervention in rescuing behavioral alterations and understand whether gut microbiota modulation mediates the effects. Breast milk microbiota composition has been demonstrated to be

influenced by several factors including maternal weight [73]. Thus, the effect of maternal weight on milk microbiota, which in turn influences the offspring microbiota, could also be an additional mechanism by which the offspring behavior is modulated.

Although detailed analysis of the microbiota in patients with ADHD is lacking, there is evidence about the link between obesity and ADHD [74], and between obesity and alteration of the gut microbiota [75]. Obesity induces a low-grade inflammatory state which has been associated with behavioral and cognitive alterations, being gut microbiota most likely an important mediator between inflammation and altered behaviors [76]. Overall, data from GF mice studies, antibiotic treatment studies, and probiotic interventions suggest that alterations in gut microbiota that reduce the inflammatory state also reduce stress-related behaviors, supporting the role of the gut microbiota as a mediator between inflammation and behavioral alterations. This evidence points for a potential role of intervention in the gut ecosystem in obese mothers, mothers who experience stress during pregnancy and/or are at risk of delivering prematurely to reduce the risk of induced behavioral alterations in offspring in the future.

Infant nutrition and childhood diet

Infant nutrition in early life (mainly type of feeding, the time of the introduction of solid food and the pattern of complementary food introduced) is a key element that shapes gut microbiota composition and, remarkably, a protective role of breastfeeding in ADHD development has been suggested [77, 78]. Furthermore, diet is considered one of the most crucial factors impacting human gut microbiota composition and functionality [79] and has emerged as one of the modifiable factors that can help protect against mental health disorders. To date, many studies have reported associations between diet and the risk of developing psychiatric disorders with gut microbiota dysbiosis considered one of the important players mediating this association. Unhealthy diet is a risk factor for depression and recent evidence suggests that dietary manipulation of the gut microbiota may have significant value in preventing and treating emotional conditions [80]. In this context, a meta-analysis including 22 studies investigating the protective effects of the Mediterranean-style diet on brain diseases demonstrated that higher adherence to this specific diet was associated with a reduced risk for depression and cognitive decline [81]. A recent prospective study in a large cohort showed that unhealthy maternal and early postnatal dietary patterns, independently of other potential confounding factors, elevate the risk of behavioral and emotional problems in the offspring [82]. In addition,

an overall shift away from traditional lifestyles has been linked to changes in the microbiota composition as well as to the increased rates of depression and other mental health disorders including ADHD in independent studies. Diet has been proposed as one of the most important environmental factors that seem to influence the risk of developing ADHD in children and its prognosis or manifestations [83]. Specific diets have been suggested to reduce ADHD symptoms including sugar-restricted, additive/preservative-free, oligoantigenic elimination diet, and fatty acid supplements. In addition, it has been proposed that chronic deficiencies of certain minerals such as zinc, iron, magnesium and iodine and insufficient dietary intake of long-chain polyunsaturated fatty acids (PUFAs) may have a significant impact on the development and symptoms of ADHD in children [83].

Few years ago a study showed considerable improvement effects of a restricted elimination diet in a group of children with ADHD [84]. In addition, a recent systematic review and meta-analysis indicated that artificial food color exclusion, and to a lesser extent free fatty acid supplementation, significantly reduce symptom severity in ADHD [85]. Nowadays, it is still unclear how both artificial food coloring and free fatty acids could impact ADHD outcomes. However, the well-known immunomodulatory effects of dietary fatty acids [86] could be one of the underlying mechanisms that, in turn, could be linked to gut microbiota changes [87, 88]. Indeed, recently common food additives called emulsifiers have been found to promote colitis and metabolic syndrome in mice by altering gut microbiota [89]. Although further studies are needed to generalize and shed light into the underlying mechanisms that could explain the link between the intake of artificial food coloring and ADHD, we may speculate that theoretically changes on microbiota composition could be involved.

According to the results of a recent meta-analysis, omega-3 levels are reduced in children with ADHD and omega-3 dietary supplementation appears to provide modest although significant evidence of clinical efficacy for improving ADHD symptoms [90]. Omega-3 influences the levels of neurotrophins, molecules that increase neuronal growth and survival. Particularly, changes on the levels of BDNF have been reported to be associated with dietary omega-3 fatty acids intake [91]. A prospective study identified an association between a “Western” dietary pattern (high in refined sugar and animal fats) and ADHD onset in adolescents [92]. Western diets, rich in saturated fat and simple sugars, are known to be major contributors to gut dysbiosis. In contrast, rural or traditional diets rich in complex carbohydrates (fiber) are associated with increased proportions of beneficial bacterial species in the human gut microbiota. A major characteristic of the modern Western diet is the increase in the consumption of omega-6 PUFAs

relative to the consumption of the anti-inflammatory omega-3. A decreased dietary intake of omega-3 PUFAs has been associated with many inflammatory-related conditions including psychiatric disorders [93] whereas omega-3 supplementation has been shown to be protective against depression, dementia and age-related cognitive decline [94]. It is noteworthy that the proposed mechanisms by which omega-3 PUFAs may beneficially influence mental health are the regulation of BDNF and neurotransmitters levels as well as regulation of HPA axis activity, which have been linked to changes in behaviors; nevertheless, whether PUFA-induced gut microbiota changes could be involved in those effects remains to be determined.

Diets rich in polyphenols have been suggested to help maintain normal brain function in both large-scale epidemiological studies [95] and intervention studies [96]. In the context of ADHD, a polyphenolic extract from pine bark has been shown to reduce hyperactivity in ADHD children together with a decrease in catecholamine excretion and oxidative stress [97]. Although different mechanisms have been proposed to account for the positive effects of polyphenols in mental health, including their anti-oxidant and anti-inflammatory properties, an indirect effect via modulation of gut microbiota composition cannot be rule out. In fact, approximately 90% of dietary polyphenols are accumulated in the large intestine where they are broken down into less complex metabolites by gut microbiota [98]. It has been described that polyphenol metabolites modulate gut microbiota composition, favoring the growth of potentially beneficial bacteria such as *Bifidobacterium* spp. and reducing that of potentially pathogenic microorganisms in both animal and human studies [99, 100].

There is an increased prevalence of both constipation and fecal incontinence in children who suffer ADHD and this occurs independently of any pharmacotherapy [101]. These findings suggest that gut dysbiosis could be associated with the gastrointestinal dysfunctions observed in ADHD patients and could play a role in brain functioning and behavior in these cases. Increasing evidence also suggests that food allergies in children and adults are associated with behavioral problems and neuropsychiatric disorders, including ADHD, and dietary-based treatments in children with allergic disorders have shown to reduce ADHD-like behavior [102].

Of note, most of the aforementioned factors associated with both ADHD development and alterations in gut microbiota composition are also factors related to the development of many other diseases. Therefore, the combination of the effects of host genetics and microbiome together with those of environmental factors could be indispensable for explaining ADHD development and its different manifestations.

In a recent human study, the administration of a specific probiotic during the first 6 months of life seems to reduce

the risk of ADHD development at 13 years of age, although differences in gut microbiota composition were not detected [103]. This finding, although encouraging, requires confirmation in well-powered studies. In addition, further studies are also needed to progress in the understanding of the possible role of intestinal dysbiosis in the risk of developing ADHD and clinical manifestation of the disease by applying high-throughput omics technologies. In addition, the role of the activity of gut microbiota in drug metabolism should also be investigated since this might partially explain the inter-individual variability observed in the efficacy and side effects of ADHD treatment.

Gut microbiota, oxidative stress and immune imbalance in ADHD

Based on the described current evidences, a new potential theory of immunopathogenesis for ADHD might be proposed. Nowadays, it is well recognized that under normal physiological circumstances there is a balance within the gut microbiota composition, known as eubiosis state, strongly associated with immune homeostasis. However, many environmental factors such as diet or even endogenous signals, caused by genetic or epigenetic factors, may disturb gut microbiota homeostasis leading to a microbial imbalance recognized as dysbiosis. This microbial imbalance is characterized by an increase of potentially inflammatory microbes and associated with an immune homeostasis breakdown linked to many immune mediated disorders. In turn, those microbiota alterations, by disrupting the intestinal permeability, may trigger migration of intestinal bacteria into the systemic circulation (microbial translocation) which might contribute to systemic inflammation [104–106]. Systemic inflammation and also other factors may contribute to the breakdown of the blood–brain barrier and, consequently, to neuroinflammation associated with different mental health problems such as ADHD. So far several studies have reported increased serum levels of pro-inflammatory cytokines such as IFN- γ and IL-16 in ADHD patients [107]. Furthermore, currently it is well recognized that the human immune system and gut microbiota interact with each other in such a way that the microbiota strongly modulate the immune system and vice versa. Therefore, it is tempting to speculate that the compromised immune balance linked to ADHD might be triggered by alterations in the gut microbiota composition and also that the dysregulated immune profile, in turn, may lead to a more compromised gut microbiota balance. Gut microbiota composition also modulate oxidative stress [108, 109] which has also been reported to be significantly increased in ADHD patients [110–112]. Oxidative stress, known as the imbalance

between production of reactive oxygen species and their elimination by protective mechanisms such as antioxidants, may be contributing to the neuronal damage and the abnormal neurotransmission linked to ADHD. Therefore, it could hypothesize that both features, inflammation and oxidative stress, partly triggered by alterations in gut microbiota composition and associated with ADHD may be playing an important role in the etiopathogenesis of the disorder contributing to the development of ADHD symptoms by inducing neuroinflammation.

Conclusions and future perspectives

Nowadays we are beginning to understand the extent to which the gut microbiota influences brain function and behavior via the gut–brain axis, being this axis an appealing therapeutic target for treating developmental disorders such as ADHD, possibly without side effects. There is preliminary evidence indicating that specific diets or dietary components (including probiotics) may alter brain activity in regions of relevance to cognitive performance, behavior and specific ADHD symptoms and diet is well known to modify the gut microbiota. In addition, it has been reported mainly in preclinical studies but also in a few human studies that brain function and behavior are highly dependent on the gut microbiota composition and many evidences suggest that gut microbiota might be linked to the risk of developing ADHD and/or its manifestations. However, so far there is no a single specific study investigating the role of the gut microbiota in ADHD and thus, we strongly argue that the study of the role of the gut microbiota in ADHD is currently more than warranted. In addition, it would be interesting to examine whether or not some microbiome alterations associated with ADHD in humans are causally related to the disorder or its different manifestations in experimental models using a translational approach. Furthermore, it would be desirable to find out the exact altered routes of communication between the brain and the gut microbiota involved in ADHD to better understand the mechanistic basis of this disorder. These altogether could pave the way towards microbiota-targeted interventions that might improve the disease management.

Acknowledgements This work was supported by Grant AGL2014-52101-P from the Spanish Ministry of Economy and Competitiveness (MINECO). The Sara Borrell postdoctoral contract of MCC from ISCIII and the PTA contract of IC from MINECO are fully acknowledged.

Compliance with ethical standards

Conflict of interest The authors declare no conflict of interest.

References

1. Martens G, van Loo K (2007) Genetic and environmental factors in complex neurodevelopmental disorders. *Curr Genom* 8:429–444
2. Polanczyk G, de Lima MS, Horta BL, Biederman J, Rohde LA (2007) The worldwide prevalence of ADHD: a systematic review and metaregression analysis. *Am J Psychiatry* 164:942–948
3. Kooij SJJ et al (2010) European consensus statement on diagnosis and treatment of adult ADHD: the European network adult ADHD. *BMC Psychiatry* 10:67
4. Trull TJ, Tragesser SL, Solhan M, Schwartz-Mette R (2007) Dimensional models of personality disorder: diagnostic and statistical manual of mental disorders fifth edition and beyond. *Curr Opin Psychiatry* 20:52–56
5. Larson K, Russ SA, Kahn RS, Halfon N (2011) Patterns of comorbidity, functioning, and service use for US children with ADHD, 2007. *Pediatrics* 127:462–470
6. Dias TGC et al (2013) Developments and challenges in the diagnosis and treatment of ADHD. *Rev Bras Psiquiatr* 35:S40–S50
7. Childress AC, Sallee FR (2014) Attention-deficit/hyperactivity disorder with inadequate response to stimulants: approaches to management. *CNS Drugs* 28:121–129
8. Shyu Y-C et al (2015) Attention-deficit/hyperactivity disorder, methylphenidate use and the risk of developing schizophrenia spectrum disorders: a nationwide population-based study in Taiwan. *Schizophr Res* 168:161–167
9. Martinez-Raga J, Knecht C, Szerman N, Martinez MI (2013) Risk of serious cardiovascular problems with medications for attention-deficit hyperactivity disorder. *CNS Drugs* 27:15–30
10. Sprich S, Biederman J, Crawford MH, Mundy E, Faraone SV (2000) Adoptive and biological families of children and adolescents with ADHD. *J Am Acad Child Adolesc Psychiatry* 39:1432–1437
11. Thapar A, Cooper M, Eyre O, Langley K (2013) Practitioner review: what have we learnt about the causes of ADHD? *J Child Psychol Psychiatry* 54:3–16
12. Sharma A, Couture J (2014) A review of the pathophysiology, etiology, and treatment of attention-deficit hyperactivity disorder (ADHD). *Ann Pharmacother* 48:209–225
13. Banaschewski T, Becker K, Scherag S, Franke B, Coghill D (2010) Molecular genetics of attention-deficit/hyperactivity disorder: an overview. *Eur Child Adolesc Psychiatry* 19:237–257
14. Dinan TG, Cryan JF (2015) The impact of gut microbiota on brain and behaviour: implications for psychiatry. *Curr Opin Clin Nutr Metab Care* 18:552–558
15. Borre YE et al (2014) Microbiota and neurodevelopmental windows: implications for brain disorders. *Trends Mol Med*. doi:10.1016/j.molmed.2014.05.002
16. Cryan JF, Dinan TG (2012) Mind-altering microorganisms: the impact of the gut microbiota on brain and behaviour. *Nat Rev Neurosci* 13:701–712
17. McVey Neufeld K-A, Luczynski P, Seira Oriach C, Dinan TG, Cryan JF (2016) What's bugging your teen? The microbiota and adolescent mental health. *Neurosci Biobehav Rev*. doi:10.1016/j.neubiorev.2016.06.005
18. Luczynski P et al (2016) Growing up in a bubble: using germ-free animals to assess the influence of the gut microbiota on brain and behavior. *Int J Neuropsychopharmacol*. 19(8). doi:10.1093/ijnp/pyw020
19. Cryan JF, Dinan TG (2015) More than a gut feeling: the microbiota regulates neurodevelopment and behavior. *Neuropsychopharmacology* 40:241–242

20. De Palma G et al (2015) Microbiota and host determinants of behavioural phenotype in maternally separated mice. *Nat Commun* 6:7735
21. Collins SM, Kassam Z, Bercik P (2013) The adoptive transfer of behavioral phenotype via the intestinal microbiota: experimental evidence and clinical implications. *Curr Opin Microbiol* 16:240–245
22. Rogers GB et al (2016) From gut dysbiosis to altered brain function and mental illness: mechanisms and pathways. *Mol Psychiatry* 21:738–748
23. Lyte M (2011) Probiotics function mechanistically as delivery vehicles for neuroactive compounds: microbial endocrinology in the design and use of probiotics. *Bioessays* 33:574–581
24. Bercik P et al (2011) The intestinal microbiota affect central levels of brain-derived neurotrophic factor and behavior in mice. *Gastroenterology* 141:599–609 (**609.e1–3**)
25. Collins SM, Bercik P (2009) The relationship between intestinal microbiota and the central nervous system in normal gastrointestinal function and disease. *Gastroenterology* 136:2003–2014
26. Barrett E, Ross RP, O'Toole PW, Fitzgerald GF, Stanton C (2012) γ -Aminobutyric acid production by culturable bacteria from the human intestine. *J Appl Microbiol* 113:411–417
27. Dinan TG, Stanton C, Cryan JF (2013) Psychobiotics: a novel class of psychotropic. *Biol Psychiatry* 74:720–726
28. Clarke G et al (2012) A distinct profile of tryptophan metabolism along the kynurenine pathway downstream of toll-like receptor activation in irritable bowel syndrome. *Front Pharmacol* 3:90
29. O'Mahony SM, Clarke G, Borre YE, Dinan TG, Cryan JF (2014) Serotonin, tryptophan metabolism and the brain–gut–microbiome axis. *Behav Brain Res* 277:32–48
30. Maes M, Mihaylova I, Ruyter M De, Kubera M, Bosmans E (2007) The immune effects of TRYCATs (tryptophan catabolites along the IDO pathway): relevance for depression—and other conditions characterized by tryptophan depletion induced by inflammation. *Neuro Endocrinol Lett* 28:826–831
31. Guillemin GJ (2012) Quinolinic acid, the inescapable neurotoxin. *FEBS J* 279:1356–1365
32. O'Mahony SM, Clarke G, Dinan TG, Cryan JF (2015) Early life adversity and brain development: is the microbiome a missing piece of the puzzle? *Neuroscience*. doi:[10.1016/j.neuroscience.2015.09.068](https://doi.org/10.1016/j.neuroscience.2015.09.068)
33. Clarke G et al (2013) The microbiome–gut–brain axis during early life regulates the hippocampal serotonergic system in a sex-dependent manner. *Mol Psychiatry* 18:666–673
34. Desbonnet L, Garrett L, Clarke G, Bienenstock J, Dinan TG (2008) The probiotic *Bifidobacteria infantis*: an assessment of potential antidepressant properties in the rat. *J Psychiatr Res* 43:164–174
35. Foley KA, MacFabe DF, Vaz A, Ossenkopp K-P, Kavaliers M (2014) Sexually dimorphic effects of prenatal exposure to propionic acid and lipopolysaccharide on social behavior in neonatal, adolescent, and adult rats: implications for autism spectrum disorders. *Int J Dev Neurosci* 39:68–78
36. MacFabe DF et al (2007) Neurobiological effects of intraventricular propionic acid in rats: possible role of short chain fatty acids on the pathogenesis and characteristics of autism spectrum disorders. *Behav Brain Res* 176:149–169
37. Stilling RM, Dinan TG, Cryan JF (2014) Microbial genes, brain and behaviour—epigenetic regulation of the gut–brain axis. *Genes Brain Behav* 13:69–86
38. Erny D et al (2015) Host microbiota constantly control maturation and function of microglia in the CNS. *Nat Neurosci* 18:965–977
39. Corominas-Roso M et al (2013) Decreased serum levels of brain-derived neurotrophic factor in adults with attention-deficit hyperactivity disorder. *Int J Neuropsychopharmacol* 1–9. doi:[10.1017/S1461145712001629](https://doi.org/10.1017/S1461145712001629)
40. Sucksdorff M et al (2015) Preterm birth and poor fetal growth as risk factors of attention-deficit/hyperactivity disorder. *Pediatrics*. doi:[10.1542/peds.2015-1043](https://doi.org/10.1542/peds.2015-1043)
41. Rodriguez A, Bohlin G (2005) Are maternal smoking and stress during pregnancy related to ADHD symptoms in children? *J Child Psychol Psychiatry* 46:246–254
42. Zijlmans MAC, Korpela K, Riksen-Walraven JM, de Vos WM, de Weerth C (2015) Maternal prenatal stress is associated with the infant intestinal microbiota. *Psychoneuroendocrinology* 53:233–245
43. Barrett E et al (2013) The individual-specific and diverse nature of the preterm infant microbiota. *Arch Dis Child Fetal Neonatal Ed* 98:F334–F340
44. Jakobsson HE et al (2014) Decreased gut microbiota diversity, delayed bacteroidetes colonisation and reduced Th1 responses in infants delivered by caesarean section. *Gut* 63:559–566
45. Grönlund MM, Lehtonen OP, Eerola E, Kero P (1999) Fecal microflora in healthy infants born by different methods of delivery: permanent changes in intestinal flora after cesarean delivery. *J Pediatr Gastroenterol Nutr* 28:19–25
46. Dominguez-Bello MG et al (2010) Delivery mode shapes the acquisition and structure of the initial microbiota across multiple body habitats in newborns. *Proc Natl Acad Sci USA* 107:11971–11975
47. Boksa P, El-Khodori BF (2003) Birth insult interacts with stress at adulthood to alter dopaminergic function in animal models: possible implications for schizophrenia and other disorders. *Neurosci Biobehav Rev* 27:91–101
48. Curran EA et al (2015) Association between obstetric mode of delivery and autism spectrum disorder: a population-based sibling design study. *JAMA psychiatry* 72:935–942
49. Curran EA et al (2015) Research review: birth by caesarean section and development of autism spectrum disorder and attention-deficit/hyperactivity disorder: a systematic review and meta-analysis. *J Child Psychol Psychiatry* 56:500–508
50. Talge NM, Allswede DM, Holzman C (2016) Gestational age at term, delivery circumstance, and their association with childhood attention deficit hyperactivity disorder symptoms. *Paediatr Perinat Epidemiol* 30:171–180
51. Amiri S, Malek A, Sadegfard M, Abdi S (2012) Pregnancy-related maternal risk factors of attention-deficit hyperactivity disorder: a case-control study. *ISRN Pediatr* 2012:458064
52. Chu S-M et al (2012) The relationship between attention deficit hyperactivity disorder and premature infants in Taiwanese: a case control study. *BMC Psychiatry* 12:85
53. Johnson S et al (2016) Antecedents of attention-deficit/hyperactivity disorder symptoms in children born extremely preterm. *J Dev Behav Pediatr* 37:285–297
54. Becattini S, Taur Y, Pamer EG (2016) Antibiotic-induced changes in the intestinal microbiota and disease. *Trends Mol Med* 22:458–478
55. Sternbach H, State R (1997) Antibiotics: neuropsychiatric effects and psychotropic interactions. *Harv Rev Psychiatry* 5:214–226
56. Van den Bergh BRH, Marcoen A (2004) High antenatal maternal anxiety is related to ADHD symptoms, externalizing problems, and anxiety in 8- and 9-year-olds. *Child Dev* 75:1085–1097
57. Li J, Olsen J, Vestergaard M, Obel C (2010) Attention-deficit/hyperactivity disorder in the offspring following prenatal maternal bereavement: a nationwide follow-up study in Denmark. *Eur Child Adolesc Psychiatry* 19:747–753
58. Grizenko N, Shayan YR, Polotskaia A, Ter-Stepanian M, Joober R (2008) Relation of maternal stress during pregnancy to

- symptom severity and response to treatment in children with ADHD. *J Psychiatry Neurosci* 33:10–16
59. Culhane JF et al (2001) Maternal stress is associated with bacterial vaginosis in human pregnancy. *Matern Child Health J* 5:127–134
 60. Sudo N et al (2004) Postnatal microbial colonization programs the hypothalamic-pituitary-adrenal system for stress response in mice. *J Physiol* 558:263–275
 61. Sterley T-L, Howells FM, Russell VA (2013) Maternal separation increases GABA(A) receptor-mediated modulation of norepinephrine release in the hippocampus of a rat model of ADHD, the spontaneously hypertensive rat. *Brain Res* 1497:23–31
 62. Womersley JS, Hsieh JH, Kellaway LA, Gerhardt GA, Russell VA (2011) Maternal separation affects dopamine transporter function in the spontaneously hypertensive rat: an in vivo electrochemical study. *Behav Brain Funct* 7:49
 63. Sanz Y (2016) Bifidobacteria in Foods: health Effects. *Encycl Food Heal* 1:388–394
 64. Mann JR, McDermott S (2011) Are maternal genitourinary infection and pre-eclampsia associated with ADHD in school-aged children? *J Atten Disord* 15:667–673
 65. Silva D, Colvin L, Hagemann E, Bower C (2014) Environmental risk factors by gender associated with attention-deficit/hyperactivity disorder. *Pediatrics* 133:e14–e22
 66. Hsiao EY et al (2013) Microbiota modulate behavioral and physiological abnormalities associated with neurodevelopmental disorders. *Cell* 155:1451–1463
 67. Derecki NC et al (2010) Regulation of learning and memory by meningeal immunity: a key role for IL-4. *J Exp Med* 207:1067–1080
 68. Ceylan MF et al (2014) Increased levels of serum neopterin in attention deficit/hyperactivity disorder (ADHD). *J Neuroimmunol*. doi:[10.1016/j.jneuroim.2014.06.002](https://doi.org/10.1016/j.jneuroim.2014.06.002)
 69. Rivera HM, Christiansen KJ, Sullivan EL (2015) The role of maternal obesity in the risk of neuropsychiatric disorders. *Front Neurosci* 9:194
 70. Jo H et al (2015) Maternal prepregnancy body mass index and child psychosocial development at 6 years of age. *Pediatrics* 135:e1198–e1209
 71. Rodriguez A et al (2008) Maternal adiposity prior to pregnancy is associated with ADHD symptoms in offspring: evidence from three prospective pregnancy cohorts. *Int J Obes (Lond)* 32:550–557
 72. Kang SS, Kurti A, Fair DA, Fryer JD (2014) Dietary intervention rescues maternal obesity induced behavior deficits and neuroinflammation in offspring. *J Neuroinflammation* 11:156
 73. Cabrera-Rubio R et al (2012) The human milk microbiome changes over lactation and is shaped by maternal weight and mode of delivery. *Am J Clin Nutr* 96:544–551
 74. Cortese S et al (2015) Association between ADHD and obesity: a systematic review and meta-analysis. *Am J Psychiatry appi*. doi:[10.1176/appi.ajp.2015.15020266](https://doi.org/10.1176/appi.ajp.2015.15020266)
 75. Cani PD (2013) Gut microbiota and obesity: lessons from the microbiome. *Brief Funct Genom* 12:381–387
 76. Foster JA, McVey Neufeld K-A (2013) Gut–brain axis: how the microbiome influences anxiety and depression. *Trends Neurosci* 36:305–312
 77. Mimouni-Bloch A et al (2013) Breastfeeding may protect from developing attention-deficit/hyperactivity disorder. *Breastfeed Med* 8:363–367
 78. Park S et al (2014) Protective effect of breastfeeding with regard to children's behavioral and cognitive problems. *Nutr J* 13:111
 79. David LA et al (2013) Diet rapidly and reproducibly alters the human gut microbiome. *Nature*. doi:[10.1038/nature12820](https://doi.org/10.1038/nature12820)
 80. Dash S, Clarke G, Berk M, Jacka FN (2015) The gut microbiome and diet in psychiatry: focus on depression. *Curr Opin Psychiatry* 28:1–6
 81. Psaltopoulou T et al (2013) Mediterranean diet, stroke, cognitive impairment, and depression: a meta-analysis. *Ann Neurol* 74:580–591
 82. Jacka FN et al (2013) Maternal and early postnatal nutrition and mental health of offspring by age 5 years: a prospective cohort study. *J Am Acad Child Adolesc Psychiatry* 52:1038–1047
 83. Konikowska K, Regulska-Ilow B, Różańska D (2012) The influence of components of diet on the symptoms of ADHD in children. *Rocz Państwowego Zakładu Hig* 63:127–134
 84. Pelsser LM et al (2011) Effects of a restricted elimination diet on the behaviour of children with attention-deficit hyperactivity disorder (INCA study): a randomised controlled trial. *Lancet* 377:494–503
 85. Sonuga-Barke EJS et al (2013) Nonpharmacological interventions for ADHD: systematic review and meta-analyses of randomized controlled trials of dietary and psychological treatments. *Am J Psychiatry* 170:275–289
 86. Stulnig TM, Zeyda M (2004) Immunomodulation by polyunsaturated fatty acids: impact on T-cell signaling. *Lipids* 39:1171–1175
 87. Kaliannan K, Wang B, Li X-Y, Kim K-J, Kang JX (2015) A host–microbiome interaction mediates the opposing effects of omega-6 and omega-3 fatty acids on metabolic endotoxemia. *Sci Rep* 5:11276
 88. Yu H-N et al (2014) Effects of fish oil with a high content of n-3 polyunsaturated fatty acids on mouse gut microbiota. *Arch Med Res* 45:195–202
 89. Chassaing B et al (2015) Dietary emulsifiers impact the mouse gut microbiota promoting colitis and metabolic syndrome. *Nature* 519:92–96
 90. Hawkey E, Nigg JT (2014) Omega-3 fatty acid and ADHD: blood level analysis and meta-analytic extension of supplementation trials. *Clin Psychol Rev* 34:496–505
 91. Ferreira CF et al (2014) Correlation between n-3 polyunsaturated fatty acids consumption and BDNF peripheral levels in adolescents. *Lipids Health Dis* 13:44
 92. Howard AL et al (2011) ADHD is associated with a 'Western' dietary pattern in adolescents. *J Atten Disord* 15:403–411
 93. Freeman MP et al (2006) Omega-3 fatty acids: evidence basis for treatment and future research in psychiatry. *J Clin Psychiatry* 67:1954–1967
 94. Zhang X-W, Hou W-S, Li M, Tang Z-Y (2015) Omega-3 fatty acids and risk of cognitive decline in the elderly: a meta-analysis of randomized controlled trials. *Aging Clin Exp Res*. doi:[10.1007/s40520-015-0381-9](https://doi.org/10.1007/s40520-015-0381-9)
 95. Letenneur L, Proust-Lima C, Le Gouge A, Dartigues JF, Barberger-Gateau P (2007) Flavonoid intake and cognitive decline over a 10-year period. *Am J Epidemiol* 165:1364–1371
 96. Desideri G et al (2012) Benefits in cognitive function, blood pressure, and insulin resistance through cocoa flavanol consumption in elderly subjects with mild cognitive impairment: the cocoa, cognition, and aging (CoCoA) study. *Hypertension* 60:794–801
 97. Dvořáková M et al (2007) Urinary catecholamines in children with attention deficit hyperactivity disorder (ADHD): modulation by a polyphenolic extract from pine bark (pycnogenol). *Nutr Neurosci* 10:151–157
 98. Selma MV, Espín JC, Tomás-Barberán FA (2009) Interaction between phenolics and gut microbiota: role in human health. *J Agric Food Chem* 57:6485–6501
 99. Duda-Chodak A, Tarko T, Satora P, Sroka P (2015) Interaction of dietary compounds, especially polyphenols, with the intestinal microbiota: a review. *Eur J Nutr* 54:325–341

100. Laparra JM, Sanz Y (2010) Interactions of gut microbiota with functional food components and nutraceuticals. *Pharmacol Res* 61:219–225
101. McKeown C, Hisle-Gorman E, Eide M, Gorman GH, Nyland CM (2013) Association of constipation and fecal incontinence with attention-deficit/hyperactivity disorder. *Pediatrics* 132:e1210–e1215
102. Stevens LJ, Kuczek T, Burgess JR, Hurt E, Arnold LE (2011) Dietary sensitivities and ADHD symptoms: thirty-five years of research. *Clin Pediatr (Phila)* 50:279–293
103. Pärty A, Kalliomäki M, Wacklin P, Salminen S, Isolauri E (2015) A possible link between early probiotic intervention and the risk of neuropsychiatric disorders later in childhood: a randomized trial. *Pediatr Res* 77:823–828
104. Sekirov I, Russell SL, Antunes LCM, Finlay BB (2010) Gut microbiota in health and disease. *Physiol Rev* 90:859–904
105. Chung H, Kasper DL (2010) Microbiota-stimulated immune mechanisms to maintain gut homeostasis. *Curr Opin Immunol* 22:455–460
106. Atarashi K, Honda K (2011) Microbiota in autoimmunity and tolerance. *Curr Opin Immunol* 23:761–768
107. Verlaet AAJ, Noriega DB, Hermans N, Savelkoul HFJ (2014) Nutrition, immunological mechanisms and dietary immunomodulation in ADHD. *Eur Child Adolesc Psychiatry* 23:519–529
108. Toral M et al (2014) The probiotic *Lactobacillus coryniformis* CECT5711 reduces the vascular pro-oxidant and pro-inflammatory status in obese mice. *Clin Sci* 127:33–45
109. Chen L et al (2013) *Lactobacillus acidophilus* ATCC 4356 attenuates the atherosclerotic progression through modulation of oxidative stress and inflammatory process. *Int Immunopharmacol* 17:108–115
110. Joseph N, Zhang-James Y, Perl A, Faraone SV (2015) Oxidative stress and ADHD: a meta-analysis. *J Atten Disord* 19:915–924
111. Kul M et al (2015) Evaluation of oxidative metabolism in child and adolescent patients with attention deficit hyperactivity disorder. *Psychiatry Investig* 12:361–366
112. Sezen H et al (2016) Increased oxidative stress in children with attention deficit hyperactivity disorder. *Redox Rep* 21:248–253