

Meta-Analysis

Maternal Thyroid Dysfunction During Pregnancy and the Risk of Adverse Outcomes in the Offspring: A Systematic Review and Meta-Analysis

Grace Mengqin Ge,¹ Miriam T. Y. Leung,¹ Kenneth K. C. Man,^{1,2} Wing Cheong Leung,³ Patrick Ip,⁴ Gloria H.Y. Li,^{1,5} Ian C. K. Wong,^{1,2} Annie W.C. Kung,⁶ and Ching-Lung Cheung¹

¹Centre for Safe Medication Practice and Research, Department of Pharmacology and Pharmacy, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Hong Kong Special Administrative Region, China; ²Research Department of Practice and Policy, UCL, School of Pharmacy, London WC1E 6BT, UK; ³Department of Obstetrics and Gynecology, Kwong Wah Hospital, Hong Kong; ⁴Department of Paediatrics and Adolescent Medicine, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Hong Kong Special Administrative Region, China; ⁵Department of Health Technology and Informatics, The Hong Kong Polytechnic University, Hung Hom, Hong Kong; and ⁶Department of Medicine, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Hong Kong Special Administrative Region, China

ORCID numbers: 0000-0001-7341-2039 (G. M. Ge); 0000-0002-6233-9144 (C.-L. Cheung).

Abbreviations: ADHD, attention deficit hyperactivity disorder; ASD, autism spectrum disorder; CBCL, Child Behavior Checklist; CI, confidence interval; FT4, free T4; HR, hazard ratio; PRISMA, Preferred Reporting Items for Systematic Reviews and Meta-Analyses; RCT, randomized controlled trial; TPO-Ab, thyroid peroxidase antibody.

Received: 15 May 2020; Accepted: 13 August 2020; First Published Online: 18 August 2020; Corrected and Typeset: 29 September 2020.

Abstract

Context: Previous studies suggested a potential link of maternal thyroid dysfunction with adverse neurocognitive outcomes and impaired development of internal organs in offspring.

Objective: To review the association between maternal thyroid dysfunction and the risk of adverse outcomes in offspring.

Data Sources: PubMed, EMBASE, and Cochrane Library.

Study Selections: Eligible studies reported the association between maternal thyroid hormone function and the risk of adverse outcomes in their children.

Data Extraction: Reviewers extracted data on study characteristics and results independently.

Data Synthesis: Estimates were pooled and reported as odds ratio (OR) with 95% confidence interval (CI). I^2 tests were applied to assess the heterogeneity across studies.

Results: We identified 29 eligible articles and found an association between maternal hyperthyroidism and attention deficit hyperactivity disorder (ADHD) (OR: 1.18, 95% CI: 1.04–1.34, $I^2 = 0\%$) and epilepsy (OR: 1.19, 95% CI: 1.08–1.31, $I^2 = 0\%$) in offspring; as well as an association of maternal hypothyroidism with increased risk of ADHD (OR: 1.14, 95% CI: 1.03–1.26, $I^2 = 25\%$), autism spectrum disorder (OR: 1.41, 95% CI: 1.05–1.90, $I^2 = 63\%$), and epilepsy (OR: 1.21, 95% CI: 1.06–1.39, $I^2 = 0\%$) in offspring.

Conclusion: Routine measurement and timely treatment on thyroid function should be considered for pregnant women.

Key Words: gestation, child development, offspring, pregnancy, thyroid, neurodevelopmental disorders

Thyroid dysfunction is among the most prevalent endocrine disorders during pregnancy (1, 2). Hyperthyroidism occurs in about 0.2% of pregnant women, with Graves' disease being the major underlying cause (3). Maternal hyperthyroidism is associated with severe adverse effects in both mother and fetus, including preeclampsia, preterm delivery, heart failure, and in utero growth retardation (4, 5). Conversely, the incidence of hypothyroidism in pregnancy is estimated to be from 0.3% to 3%, and it is particularly common in regions with iodine deficiency (1). Maternal hypothyroidism is associated with an increased risk of in utero growth retardation, miscarriage, preeclampsia, placental abruption, and fetal death (1, 6, 7). In addition to hyperthyroidism and hypothyroidism, as high as 18% (8) of pregnant women are tested positive for thyroid peroxidase antibody (TPO-Ab) or thyroglobulin antibody. TPO-Ab positivity regulates the effect of maternal thyroid status unfavorably on pregnancy and the developing fetus (8, 9). Thus, maintaining a normal thyroid status in pregnancy is important for both the mother and the offspring.

The growth and development of the fetus are totally dependent on maternal thyroid hormones until the onset of fetal thyroid function near mid-gestation (10). Thus, the transfer of maternal thyroid hormones to the developing fetus is critical during the early stage of pregnancy (10–12). The hypothesis of fetal programming (10, 13) suggested that maternal thyroid dysfunction can disturb early fetal brain development and lead to subsequent onset of neurodevelopmental disorders during the childhood of the offspring. This makes it biologically plausible to investigate the potential association between maternal thyroid dysfunction during pregnancy and the long-term health outcome of the offspring (14, 15). Moreover, because the thyroid hormone receptor is widely expressed in multiple tissues, abnormal maternal thyroid status may also lead to non-neurodevelopmental disorders in the offspring, such as metabolic or endocrine disorders (16, 17), impaired cardiovascular function (18, 19), and respiratory system (20, 21). Although maternal thyroid status may have a profound impact on the offspring, no systematic reviews were conducted to evaluate the relationship.

Therefore, we conducted a systematic review and meta-analysis of published studies to review and meta-analyze the association between maternal thyroid dysfunction during pregnancy and the risk of various adverse outcomes in the offspring.

Methods

This systematic review was reported following The Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines (22) in Appendix 1 (<https://osf.io/k9jpf> (23)).

Study search

We performed a systematic literature search in PubMed, EMBASE, and Cochrane Library from inception to 1 November 2019, with an updated search conducted on 1 February 2020. Articles from previous reviews or included studies were also reviewed. No restrictions on study design or language were applied in the search, but only original research studies published in peer-reviewed journals were included. The search strategy was developed for PubMed and applied to other databases. The keywords were developed based on 3 concepts: namely (1) mothers and pregnancy, (2) thyroid hormone status, and (3) offspring. The full predefined search terms and search strategy are provided in the Supplementary Table 1 (<https://osf.io/k9jpf> (23)).

Study selection and data extraction

Only full-text articles reporting the association between maternal thyroid hormone function during pregnancy and the risk of adverse outcomes in their children were included in the present systematic review and meta-analysis. Such adverse outcomes included neurodevelopmental disorders (attention deficit hyperactivity disorder [ADHD], autism spectrum disorder [ASD], epilepsy, language and speech impairment, depression, schizophrenia, and cerebral palsy), and other clinically diagnosed outcomes. However, markers of these clinical outcomes, such as fasting glucose and blood pressure, were excluded from this systematic review and meta-analysis. Whereas studies related to the thyroid function of the offspring which measured levels of TSH, free T4 (FT4), and TPO-Ab were included. Nonhuman studies, commentary reviews, or descriptive studies without adequate statistical analysis were excluded.

The titles and abstracts of the included articles were screened by 2 independent researchers (G.G. and M.L.). The 2 authors (G.G. and M.L.) independently decided on the inclusion and exclusion of the studies and conducted data extraction. Discrepancies were resolved by consensus. Information in the data extraction form included year and

country of the publication, study design, sample size, exposure timing, maternal thyroid status, child outcomes with measurement tools, age of the children at assessment, summary statistics, and variables used for potential confounder adjustment. Relative risk, incidence rate ratio, or comparable statistics from the fully adjusted model were extracted from each included study. Because the prevalence of clinical outcomes in offspring are all < 10%, odds ratio (OR) is similar to and considered as comparable to relative risk. For duplicated studies using the same cohort, the study with a larger sample size was included in the meta-analysis. Because Andersen et al, 2018 (15), is an update of Andersen et al, 2013 (24), and Andersen et al, 2014 (25), with different study design and smaller sample size, Andersen et al, 2018 (15), was excluded from the systematic review and meta-analysis.

Quality assessment

The quality of the included studies was assessed using The Newcastle-Ottawa Scale by 2 authors (G.G. and M.L.) independently. The scale consists of 3 domains (selection, comparability, and outcome), and provides a summary measure of quality for each study. Disagreements were resolved through discussion with I.W. and C.L.C.

Data analysis

All included studies were described narratively and presented in Table 1. Pooled OR was calculated using the inverse variance weighting model. Meta-analysis was conducted when data of 2 or more studies investigating the same exposure and outcome were available. The original studies' definition of maternal hyperthyroidism, hypothyroidism, and hypothyroxinemia was adopted in this meta-analysis. Maternal thyroid parameters (FT4 and TSH) were fitted as continuous variables and meta-analyzed if they had the same unit (eg, per SD). Where multiple cutoffs of FT4 or TPO-Ab were measured, we included results derived from all cutoffs. We extracted OR/hazard ratio (HR) from the adjusted model when applicable. We estimated the OR/HR based on the lower confidence limits in the original studies if only data with 1 decimal point was available. The heterogeneity of the studies was evaluated using I^2 tests. I^2 no more than 25% represented low heterogeneity; I^2 between 25% and 50% represented moderate heterogeneity; I^2 greater than 50% represented high heterogeneity. The overall effects were analyzed with a fixed-effects model if the heterogeneity was low; otherwise, a random-effects model was applied. Forest plots were generated for further visual inspection of heterogeneity. Statistical analyses were

conducted using Review Manager (RevMan 5.3; Cochrane Collaboration, Oxford, UK).

Results

Study selection

A total of 9975 records were identified by the electronic search engine. After removal of duplicates, 7914 items were evaluated based on title and abstract screening, of which 151 full-text articles were assessed for eligibility. Of these, 29 studies (14, 16-21, 24-45) met our inclusion criteria and were included in this study. The PRISMA flow diagram summarizing the search results is presented in Fig. 1.

Characteristics of included studies

Study characteristics are summarized and shown in Table 1. All included studies were observational studies based on population-based prospective birth cohorts or national register-based retrospective cohorts. Among the 29 studies included, 22 articles investigated the association between maternal thyroid dysfunction and neurodevelopmental disorders in offspring. One of these 22 articles, together with the remaining 7 articles, also investigated non-neurodevelopmental outcomes. In all these 29 studies, the nonexposure group comprised pregnant mothers without thyroid dysfunction. Estimates of maternal thyroid status and offspring outcomes extracted from these 29 studies are provided in Tables 2-5. Pooled estimates and the forest plots are provided in Figs 2-4. The corresponding funnel plots are provided in Supplementary Figures 1-3 (<https://osf.io/k9jpf> (23)).

Neurodevelopmental disorders in the offspring (N = 22)

Most studies (N = 17) were conducted in European countries, except 5 conducted in India (N = 1), Israel (N = 1), and the United States (N = 3). There are 18 cohort studies and 4 case-control studies. Maternal thyroid dysfunctions discussed were hyperthyroidism, hypothyroidism, hypothyroxinemia, and levels of maternal thyroid hormone (including TSH, FT4, and thyroperoxidase antibodies positive [TPO-Ab⁺]).

Hyperthyroidism. Seven studies investigated hyperthyroidism as exposure (Table 2), with ADHD ($n_{\text{study}} = 2$; $n_{\text{sample size}} = 15,422$) (25, 29), ASD ($n_{\text{study}} = 2$; $n_{\text{sample size}} = 14,204$) (25, 26), epilepsy ($n_{\text{study}} = 2$; $n_{\text{sample size}} = 25,495$) (14, 24), and schizophrenia ($n_{\text{study}} = 2$; $n_{\text{sample size}} = 4638$) (14, 28) in offspring as the outcomes. The pooled OR of offspring

Table 1. Characteristics of Studies with Maternal Thyroid Dysfunction Exposure and Adverse Outcomes in the Offspring

Item	Author (Y), Country	Study Design (Sample Size)	Exposure Measurement ^a (Timing), Measurement kit for laboratory assays	Mean Maternal Age (y $\pm$ SD)	Outcome Measurement ^b	Age of Children (y)	Adjusted Variables in the Model
1	Brown et al (2015), Finland	Case-control (1920)	TFT: TSH, FT4, TPO-Ab (1st and early 2nd trimesters), Abbott Diagnostics, Abbott Park, IL	TPO-Ab ⁺ : 29.62 $\pm$ 4.65 TPO-Ab ⁻ : 29.22 $\pm$ 5.30	ASD: Dx from FHDR, assessed with the ADI-R	NA	No adjusted covariate
2	Grattan et al (2014), Canada	Case-control (998)	Self-report maternal history of hypothyroidism with a standardized questionnaire (during pregnancy)	NA	CHD: Dx using echocardiography	Range 0-18, median of 2.7	Age and risk of child chromosomal disorder, maternal diabetes mellitus, maternal age, and family history of CHD
3	George et al (2014), India	Case-control (343)	Self-report Dx by interview (during pregnancy)	NA	ASD: Dx from autism clinic of Child Development Centre, using a Childhood Autism Rating Scale, score over 30	Range 2-6	Antenatal and natal risk factors for autism
4	Gyllenberg et al (2016), Finland	Case-control (2020)	TFT: TSH, FT4 (1st trimester) <i>Abbott Diagnostics, Abbott Park, IL</i>	Cases: 28.5 $\pm$ 5.5 control: 28.2 $\pm$ 5.1	Schizophrenia: FHDR, using ICD-10: F20 and ICD-10: F25	NA	Maternal psychiatric history, maternal smoking, degree of urbanization of birth municipality, birth province, twinning
5	Instanes et al (2017), Norway	Case-control (2 332 657)	Dx based on medical registry diagnosis records from MBRN and NorPD (before or during pregnancy)	NA	ADHD: Rx of ADHD medications methylphenidate (ATC code N06BA04), atomoxetine (ATC code N06BA09) and racemic amphetamine (ATC code N06BA01) extracted from the NorPD	> 3	Year of birth, parity, mother's age at birth, mother's educational level, mother's marital status, maternal and paternal use of ADHD medication
6	Andersen et al (2014), Denmark	Cohort (857 014)	Dx and Rx from DNHR and DNPR (before and after delivery)	NA	ADHD, ASD: Dx and Rx from DNHR, DNPR and DNPR ADHD: ICD-10: F90; if no Dx then at least 2 Rx of ADHD medication (ATC N06NA) ASD: ICD-10: F84	Range 3-18	Gender of the child, year of birth of the child, maternal age, maternal origin, maternal residence, maternal marriage status, income, and parity

Table 1. Continued

Item	Author (Y), Country	Study Design (Sample Size)	Exposure Measurement ^a (Timing), Measurement kit for laboratory assays	Mean Maternal Age (y $\pm$ SD)	Outcome Measurement ^b	Age of Children (y)	Adjusted Variables in the Model
7	Andersen et al (2013), Denmark	Cohort (1 699 693)	Dx from DNHR (before and after delivery)	NA	Epilepsy: Dx after the neonatal period, data from DNHR using ICD-8: 345.09-345.99 and ICD-10: G40.0-G41.9	Median of 5.3	Gender of the child, birth year, maternal age, parity including index pregnancy, maternal marriage status, income, maternal origin, maternal residence, and maternal diagnosis of febrile seizure and/or epilepsy registered in the DNHR
8	Chevrier et al (2011), U.S.	Cohort (287)	TFT: TSH, FT4 (26.9 $\pm$ 3.4 wk gestation); Siemens Healthcare Diagnostics, Deerfield, IL	N.A.	ADHD: Maternal report on the Attention Problems scale of the CBCL or child performance on the KCPT	5	Maternal age, income, employment status at 6 mo, country of birth, diet quality index, delivery complications, Peabody Picture Vocabulary Test score, child 5-min APGAR, hospitalization at 1 y; the number of children in the home at 1 and 2 y, home density at 1 y, the family structure at 1 y; the season of assessment
9	Cowett et al (1975), U.S.	Cohort (1394)	Butanol-extractable iodine determination and clinical history (during pregnancy)	NA	Respiratory distress syndrome: clinical evidence of respiratory distress syndrome (NA)	NA	No adjusted covariate
10	Eshkoli et al (2019), Israel	Cohort (217 910)	Dx based on the maternal report as well as on a routine review of all computerized medical records from the hospital and ambulatory setting from SUMC (during pregnancy)	Hypothyroidism: 30.8 $\pm$ 5 euthyroid: 28.2 $\pm$ 5	Pediatric endocrine morbidity: Dx using ICD-9 codes	NA	Gestational and pregestational diabetes mellitus, chronic, gestational or pre-eclampsia, delivery before 37 wk of gestation
11	Fetene et al (2019), U.K.	Cohort (2920)	TFT: TSH, FT4 (1st trimester); Abbott Diagnostics, Abbott Park, IL	28.9 $\pm$ 4.6	Depression: DAWBA coded according to parent-reported or self-reported DMS-IV criteria	7.5 for parent-reported depression; 1.5 for self-reported depression	Maternal age, BMI, gender, birth weight, maternal smoking, maternal depression, gestational hypertension, and zinc and iodine intake during pregnancy

Table 1. Continued

Item	Author (Y), Country	Study Design (Sample Size)	Exposure Measurements (Timing), Measurement kit for laboratory assays	Mean Maternal Age (y $\pm$ SD)	Outcome Measurements	Age of Children (y)	Adjusted Variables in the Model
12	Fetene et al (2018), U.K.	Cohort (2912)	TFT: TSH, FT4, TPO-Ab (1st trimester); Abbott Diagnostics, Abbott Park, IL	28.9 $\pm$ 4.6	ADHD: DAWBA coded according to parent-reported or self-reported DSM-IV criteria	7.5 for parent-reported ADHD; 15 for self-reported ADHD	Maternal age, BMI, sex, gestational hypertension, antenatal depression, maternal smoking, partner smoking, maternal alcohol use, birth weight, maternal iodine, and zinc intake during pregnancy
13	Frank et al (2019), Norway	Cohort (1204)	Rx records from MBRN and NorPD based on ATC classification System (during pregnancy)	NA	Language impairment: ICD-10: F80 and parent-reported symptoms of language and communication skill deficits	8	Maternal age, educational level, income, parity, BMI at conception, use of folic acid and other supplements, lifetime history of major depression, comedication for somatic and mental comorbidities, and smoking and alcohol use during pregnancy
14	Gerahun et al (2018), U.S.	Cohort (397 201)	Dx using ICD-9: 244.X and medication prescribed to treat the condition (during pregnancy)	NA	ASD: at least 1 documented DSM-IV code for ASD on any 2 separate visits	Range 2-17	maternal age, education, smoking during pregnancy, prenatal care, parity, year of diagnosis, median household income, child's sex and race/ethnicity
15	Ghassabian et al (2012), Netherland	Cohort (3139)	TFT: TSH, FT4, TPO-Ab (13.5 $\pm$ 1.8 wk of gestation); Vitros ECI Immunodiagnostic System Ortho Clinical Diagnostics, Rochester, NY	TPO-Ab ⁺ : 31.2 $\pm$ 4.3 TPO-Ab ⁻ : 31.1 $\pm$ 4.3	ADHD: CBCL completed by parents, using scales consistent with diagnostic categories of DSM-IV	3	Child's gender and ethnicity, maternal age, cigarette smoking, and time of thyroid sampling during pregnancy
16	Ghassabian et al (2011), Netherland	Cohort (3736)	TFT: TSH, FT4 (13.3 $\pm$ 1.7 wk of gestation)	30.9 $\pm$ 4.5	ADHD: CBCL completed by parents, using externalizing (attention problems and aggressive behavior) scale consistent with diagnostic categories of DSM	1.5 and 3	Maternal age, education level, and psychopathology, child's gender, ethnicity, mode of delivery, and gestational age at the time of maternal thyroid sampling

Table 1. Continued

Item	Author (Y), Country	Study Design (Sample Size)	Exposure Measurement ^a (Timing), Measurement kit for laboratory assays	Mean Maternal Age (y $\pm$ SD)	Outcome Measurement ^b	Age of Children (y)	Adjusted Variables in the Model
17	Gutvirth et al (2019), Israel	Cohort (217 910)	Dx based on the maternal report as well as medical records from the hospital and/or ambulatory settings, data from SUMC (during pregnancy)	Hypothyroidism: 30.8 $\pm$ 5.2 no hypothyroidism: 28.2 $\pm$ 5.7	ADHD, ASD, cerebral palsy: computerized pediatric hospitalization database of SUMC and the computerized perinatal database of the Obstetrics and Gynecology department	< 18	Maternal age, birth weight, pregestational and gestational diabetes and hypertensive disorders of pregnancy (pregestational, gestational hypertension and pre-eclampsia)
18	Haddow et al (1999), U.S.	Cohort (186)	TFT: TSH, FT4, TPO-Ab (2nd trimester); TSH: Diagnostic Products, LA; FT4: Wallac Oy, Turku, Finland; TPO-Ab: Kronus, San Clemente, CA	NA	ADHD: the Conners' Continuous Performance Test to measure sustained vigilance and WISC-III scores	8	No adjusted covariate
19	Heikkinen et al (2017), Finland	Cohort (3 229)	TFT: thyrotropin, FT4, TPO-Ab, Tg-Abs (before 20th week of gestation); Abbott Diagnostics, Abbott Park, IL	Euthyroid: 28.1 $\pm$ 5.3 hypothyroid: 28.4 $\pm$ 5.5 hyperthyroid: 30.0 $\pm$ 5.4 hypothyroxinemia: 30.0 $\pm$ 6.3 TPO-Ab ⁻ : 28.2 $\pm$ 5.4 TPO-Ab ⁺ : 28.6 $\pm$ 5.0	Cardiometabolic risk factor: blood sampling	16	Maternal age, smoking, parity, and overweight/obesity
20	Jølling et al (2018), Denmark	Cohort (2 618)	Dx of autoimmune hyperthyroidism/ Graves' disease (ICD-8 codes: 242.00, 242.01, 242.08, 242.09; ICD-10 code: E05.0) or autoimmune hypothyroidism/Hashimoto thyroiditis (ICD-8 code: 244.01; ICD-10 codes: E06.3, E06.3A and E06.3B) (within 1 y before delivery)	NA	1.5 disease groups covering major nonmalignant somatic and psychiatric disease categories	> 5	Sex of the child, year of birth, maternal age at birth, delivery mode, multiple births, birth order, preterm birth, small for gestational age, and maternal comorbidity of the outcome disease

Table 1. Continued

Item	Author (Y), Country	Study Design (Sample Size)	Exposure Measurement ^a (Timing), Measurement kit for laboratory assays	Mean Maternal Age (y $\pm$ SD)	Outcome Measurement ^b	Age of Children (y)	Adjusted Variables in the Model
21	Liu et al (2018), Denmark	Cohort (14 302)	Dx using ICD-8 codes: 243.99-244.09; or E03 and ICD-10 code E89.0; Rx using ATC: H03A; data from DNPR (before 5 y after delivery);	NA	Asthma: Dx using ICD-10: J45 and J46; Rx using ATC codes of inhaled β 2-agonists (R03AC02-04, -12, and -13), inhaled glucocorticoids (R03BA01, -02 and -05), fixed-dose combination of inhaled β 2-agonists and glucocorticoids (R03AK06 and -07), leukotriene receptor antagonists (R03DC03), and anti-IgE therapies (R03DX05)	> 5	Maternal age, primiparity, smoking during pregnancy, education status, income status, calendar year of birth, maternal diabetes, maternal asthma, paternal asthma at delivery, and the index child's age at observation during the study period
22	Modesto et al (2015), Finland	Cohort (3873)	TFT: thyrotropin, FT4 (range 6.6-17.9 wk, mean of 13.6 wk gestation); FT4 & TSH: Vitros Eciimmunodiagnostic system; Ortho Clinical Diagnostics; TPO-Ab: Phadia 250; Thermo Scientific	30.0 $\pm$ 5.0	ADHD: Conner's Parent Rating Scale-Revised Short Form with ADHD index	Range 7.5 to 10.5, mean (SD) of 8.1 (0.2)	Child age, sex, and ethnic background and maternal educational level, age, history of smoking, psychopathologic symptoms during pregnancy, parity, marital status, household income, BMI, and time of blood sampling in pregnancy
23	Oostenbroek et al (2017), Netherland	Cohort (2000)	TFT: TSH, FT4, TPO-Ab (median of 12.9 wk gestation); FT4 & TSH: Beckman Coulter Inc (Fullerton, California) TPO-Ab+: E-CK-96; ZenTech, Angleur, Belgium	Hypothyroxinemic: 32.2 $\pm$ 4.4 nonhypothyroxinemic: 31.8 $\pm$ 4.3	ADHD: parent and teacher reports of SDQ using borderline clinical cutoff	Range 5-6, mean (SD) of 5.1 (0.2)	Ethnicity, years of education, pregnancy BMI, hypertension, smoking during pregnancy of at least 1 cigarette per day and anxiety level
24	Pakkila et al (2013), Finland	Cohort (3673)	TFT: TSH, FT4, TPO-Ab (10.7 $\pm$ 2.8 wk gestation) Architect i2000 automatic analyzer (Abbott Diagnostics)	NA	Same serum thyroid biomarkers level as mothers: TFT	8	Maternal continuous TPO-Ab levels

Table 1. Continued

Item	Author (Y), Country	Study Design (Sample Size)	Exposure Measurement ^a (Timing), Measurement kit for laboratory assays	Mean Maternal Age (y $\pm$ SD)	Outcome Measurement ^b	Age of Children (y)	Adjusted Variables in the Model
25	Pakkila et al (2014), Finland	Cohort (5131)	TFT: TSH, FT4, TPO-Ab (10.7 $\pm$ 2.8 weeks gestation); Abbott Architect i2000 method (Abbott Diagnostics)	NA	ADHD: Parents reported Rutter B2 Scale-Finnish version	8	Having more than 2 children in the family, maternal smoking, maternal education, and maternal age
26	Petersen et al (2018), Denmark	Cohort (1 270 079)	Hypothyroidism: Dx based on ICD-8 243.99 and 244.00-244.09, ICD-10: E00, E03.0-E03.9 AND E89.0, excluding 244.02, E03.0A, E03.1B, and E03.4; Rx based on thyroid hormone, ACT: h03A hyperthyroidism: Dx based on ICD-8 as 242.00-242.29 and by ICD-10 as E05.0-E05.9, excluding E05.4, E05.8A, and E05.9A; Rx based on antithyroid medication, ACT: H03A (before and during pregnancy, and within 5 years after pregnancy)	NA	Cerebral palsy: Dx through the Danish National Cerebral Palsy Registry	Range 1-6	Birth year, maternal age, maternal diabetes, and maternal socioeconomic status and smoking and alcohol consumption in pregnancy
27	Roman et al (2013), Netherland	Cohort (4039)	TFT: TSH, FT4, TPO-Ab (range 5.9-7.913.4 $\pm$ 1.9 wk gestation); FT4&TSH: Vitros ECI Immunodiagnostic System; Ortho Clinical Diagnostics, Rochester, NY; TPO-Ab: Phadia 250 immunoassay; Phadia, Uppsala, Sweden	31.2 $\pm$ 4.7	ASD: parents reported CBCL using pervasive developmental problems scale	6	Child's sex, ethnicity, gestational age at birth, birth weight, maternal age, educational level, smoking history, prenatal psychopathology, thyroid medication during pregnancy, parity, marital status, maternal folate and C-reactive protein levels in early pregnancy, time of thyroid sampling during pregnancy, and paternal age
28	Vermiglio et al (2004), Italy	Cohort (27)	TFT: TSH, FT4 (5-10, 11-14, 18-20 wk gestation)	NA	ADHD: DMS-IV-TR items by clinical observation	Range 8-10	No adjusted covariate

Table 1. Continued

Item	Author (Y), Country	Study Design (Sample Size)	Exposure Measurement ^a (Timing), Measurement kit for laboratory assays	Mean Maternal Age (y $\pm$ SD)	Outcome Measurement ^b	Age of Children (y)	Adjusted Variables in the Model
29	Wasserman et al (2008), U.S.	Cohort (1731)	TFT: TPO-Ab (3rd trimester); QUANTA Lite TPO; INOVA Diagnostics, San Diego, CA	24.1 (range 11-46)	Sensorineural hearing loss: Dx according to pediatric case definition	8	Maternal age, race and treated clinical hypothyroidism

^aExposure refers to maternal thyroid status. Timing refers to the time of the diagnosis or prescription or the function test conducted. TFT indicates thyroid function test; Dx indicates the diagnosis of thyroid diseases; Rx indicates prescription for thyroid diseases.

^bOutcomes refer to diagnosed or assessed disorders in the offspring. Dx indicates diagnosis and Rx indicates prescriptions.

Abbreviations: ADHD, attention deficit hyperactivity disorder; ADI-R, Autism Diagnostic Interview-Revised; ASD, autism spectrum disorders; ATC, Anatomical Therapeutic Chemical Classification System; BMI, Body Mass Index; CBCL, Child Behavior Checklist; CHD, congenital heart defect; DAWBA, Developmental and Well-Being Assessment; DMS-IV-TR, Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition-Text Revision; DNH, Danish National Hospital Registry; DNPR, Danish National Prescription Registry, DPCR, Danish Psychiatric Central Register; FHDR, Finnish Hospital, and Outpatient Discharge Registry; FT4, free thyroxine; ICD, International Classification of Diseases; KCPT, Kiddie Continuous Performance Test; MBRN, Medical Birth Registry of Norway; NorPD, Norwegian Prescription Database; NA, not available; SDQ, Strengths and Difficulties Questionnaire; SUMC, Soroka University Medical Center; TgAb, thyroglobulin antibodies; TPO-Ab(+/-), thyroid peroxidase antibodies (positive/negative); WISC-III, Wechsler Intelligence Scale for Children- Third edition.

ADHD, ASD, epilepsy, and schizophrenia was 1.18 (95% confidence interval [CI]: 1.04-1.34), 1.17 (95% CI: 0.96-1.42), 1.19 (95% CI: 1.08-1.31), and 0.96 (95% CI: 0.53-1.72), respectively. The heterogeneity was estimated to be low ($I^2 = 0\%$) in all outcomes (Fig. 2, Supplementary Figure 1 [<https://osf.io/k9jpf>] (23))).

Only 1 study investigated the association between maternal hyperthyroidism and cerebral palsy, revealing a null association (42) (Table 2). Meta-analysis was not performed for this outcome.

Hypothyroidism. Among all maternal thyroid status, maternal hypothyroidism was the most frequently studied ($N = 12$). There are 4 ($n_{\text{sample size}} = 32827$), 5 ($n_{\text{sample size}} = 419878$), 2 ($n_{\text{sample size}} = 12465$), 2 ($n_{\text{sample size}} = 2780$), and 2 studies ($n_{\text{sample size}} = 29070$) investigating ADHD (25, 29, 37, 38), ASD (25-27, 34, 37), epilepsy (14, 24), schizophrenia (14, 28), and cerebral palsy (37, 42) as the clinical outcomes, respectively (Table 3). The pooled OR for offspring ADHD, ASD, epilepsy, schizophrenia, and cerebral palsy was 1.14 (95% CI: 1.03-1.26), 1.41 (95% CI: 1.05-1.90), 1.21 (95% CI: 1.06-1.39), 3.02 (95% CI: 0.26-35.51), and 0.95 (95% CI: 0.64-1.40), respectively, with low (ADHD: $I^2 = 25\%$; epilepsy/cerebral palsy: $I^2 = 0\%$), to high (ASD: $I^2 = 63\%$, schizophrenia: $I^2 = 86\%$) heterogeneity detected for each outcome (Fig. 3, Supplementary Figure 2 [<https://osf.io/k9jpf>] (23))).

Only 1 study by Frank et al (33) investigated language and speech impairment as the clinical outcome but null association was observed in the study (HR: 0.75, 95% CI: 0.38-1.43; Table 3).

The definition of hypothyroidism in the included studies were mainly overt hypothyroidism. Only 3 studies performed an exploratory analysis of maternal subclinical hypothyroidism and investigated offspring ASD (26), schizophrenia (28), and ADHD (39) as the clinical outcomes, but null association was observed.

Hypothyroxinemia. Five studies investigated the association between maternal hypothyroxinemia and offspring neurodevelopmental disorders (Table 4). Three studies investigated ADHD as a clinical outcome. Modesto et al (39) found children exposed to maternal hypothyroxinemia during pregnancy had higher ADHD scores compared with nonexposed children (β : 0.07, 95% CI: 0.003-0.14). Similarly, Oostenbroek et al (40) found maternal hypothyroxinemia with FT4 < 5th percentile was associated with a 1.70-fold (95% CI: 1.01-2.86) increase in odds of teacher-reported ADHD, although a null association was observed when parent-reported ADHD was used as the outcome. Vermiglio et al (44) reported a significantly higher prevalence of ADHD among the offspring of




Figure 1. Preferred Reporting Items for Systematic Reviews and Meta-Analyses flow diagram of included studies.

Table 2. Estimated Associations Between Maternal Hyperthyroidism and Offspring Neurodevelopmental Disorders

Item	Author (Y)	Outcome Measurement	Estimate (95% CI)
<i>ADHD as outcome</i>			
1	Andersen et al (2014)	ICD-10 diagnosis and prescription of drugs	HR = 1.18 (1.03-1.36)
2	Instanes et al (2017)	Prescription of drugs	OR = 1.2 (0.9-1.5)
<i>ASD as outcome</i>			
1	Andersen et al. (2014)	ICD-10 diagnosis	HR = 1.18 (0.96- 1.45)
2	Brown et al. (2015)	ICD-10 diagnosis assessed with the Autism Diagnostic Interview-Revised	OR = 1.06 (0.54-2.10)
<i>Epilepsy as outcome</i>			
1	Andersen et al. (2013)	ICD-10 diagnosis	HR = 1.20 (1.09-1.32)
2	Jolving et al. (2018)	ICD-10 diagnosis	HR = 1.04 (0.70-1.55)
<i>Cerebral palsy as outcome</i>			
1	Petersen et al. (2018)	ICD-8 and ICD-10 diagnosis	OR = 1.11 (0.94-1.66)
<i>Schizophrenia as outcome</i>			
1	Gyllenberg et al. (2016)	ICD-10 diagnosis	OR = 0.87 (0.41-1.82)
2	Jolving et al. (2018)	ICD-10 diagnosis	HR = 1.13 (0.43-3.01)

Abbreviations: ADHD, attention deficit hyperactivity disorder; ASD, autism spectrum disorders; CI, confidence interval; HR, hazard ratio; ICD: International Classification of Diseases; OR, odds ratio.

mothers with hypothyroxinemia in the moderately iodine-deficient region.

Only 1 study was conducted for the association of maternal hypothyroxinemia with offspring ASD and

schizophrenia (Table 4). Roman et al (43) reported severe maternal hypothyroxinemia ($FT4 < 5$ th percentile) was associated with increased odds of pervasive developmental problems in offspring (OR: 2.60, 95% CI: 1.30-5.18).

Table 3. Estimated Associations Between Maternal Hypothyroidism and Offspring Neurodevelopmental Disorders

Item	Author (Y)	Outcome Measurement	Estimate (95% CI)
<i>ADHD as outcome</i>			
1	Andersen et al (2014)	Prescription of drugs	HR = 1.10 (0.98-1.25)
2	Gutvirtz et al (2018)	ICD-9 diagnosis	HR = 2.25 (0.31-16.16)
3	Haddow et al (1999) ^a	Conner's Continuous Performance Test > 8 ^a WISC-III freedom from distractibility score ^b	OR = 3 (1-5) Mean difference ± SE = -3 ± 2
4	Instanes et al (2017)	Prescription of drugs	OR = 1.2 (1.0-1.4)
<i>ASD as outcome</i>			
1	Andersen et al (2014)	ICD-10 diagnosis	HR = 1.30 (1.11-1.53)
2	Brown et al (2015)	ICD-10 diagnosis assessed with the Autism Diagnostic Interview-Revised	OR = 0.67 (0.27-1.63)
3	Getahun et al (2018)	Clinical diagnosis	HR = 1.31 (1.13-1.53)
4	Gutvirtz et al (2018)	ICD-9 diagnosis	HR = 5.22 (0.70-38.56)
5	George et al (2014)	Clinical diagnosis	OR = 4.25 (1.38-13.07)
<i>Epilepsy as outcome</i>			
1	Andersen et al (2013)	ICD-10 diagnosis	HR = 1.22 (1.06-1.40)
2	Jolving et al (2018)	ICD-10 diagnosis	HR = 0.94 (0.39-2.27)
<i>Language and speech impairment as outcome</i>			
1	Frank et al (2019)	ICD-10 diagnosis	HR = 0.75 (0.38-1.43)
<i>Schizophrenia as outcome</i>			
1	Gyllenberg et al (2016)	ICD-10 diagnosis	OR = 0.86 (0.23-3.24)
2	Jolving et al. (2018)	ICD-10 diagnosis	HR = 10.64 (2.80-40.41)
<i>Cerebral palsy as outcome</i>			
1	Gutvirtz et al (2018)	ICD-9 diagnosis	HR = 0.88 (0.13-6.31)
2	Petersen et al (2018)	ICD diagnosis	OR = 0.95 (0.64-1.39)

Haddow et al. used Continuous Performance Test score and WISC-III freedom-from-distractibility score to measure offspring ADHD.

^aA higher score indicates more problems in this scale. The author took score 8 as cutoff point and this row provides OR for the children of mothers with hypothyroidism as compared with the control children.

^bThis row provides the difference as the value in the case minus the average of the values in the control; the value expressed as means ± standard error of the individual differences in each matched set.

Abbreviations: ADHD, attention deficit hyperactivity disorder; ASD, autism spectrum disorders; CI, confidence interval; HR, hazard ratio; ICD: International Classification of Diseases; OR, odds ratio; WISC-III: Wechsler Intelligence Scale for Children, Third edition.

Gyllenberg et al (28). reported a positive association between maternal hypothyroxinemia and offspring schizophrenia (OR: 1.75, 95% CI: 1.22-2.50).

Thyroid hormone levels. Eleven studies investigated the association between maternal thyroid hormone parameters and offspring neurodevelopmental disorders (Table 5). Fetene et al (32) found null association between maternal TPO-Ab⁺ with offspring ADHD, but Ghassabian et al (36) observed an association between elevated titers of TPO-Abs and the risk of ADHD (OR = 1.77, 95% CI: 1.15-2.72; Table 5). Two studies (26, 43) investigated the association of maternal TPO-Ab⁺ with offspring ASD (n_{sample size} = 5959), and the pooled OR is 1.30 (95% CI: 0.58-2.94, I² = 60%) (Fig. 4, Supplementary Figure 3 [https://osf.io/k9jpf (23)]). Notably, the 2 studies used different cutoff points to define TPO-Ab⁺.

Four studies investigated the association of maternal FT4 level with offspring ADHD. Chevrier et al (30) and Modesto et al (39) found a null association between maternal FT4

level and offspring scores on ADHD (Table 5). The other 2 studies (32, 40) included offspring ADHD as a binary outcome and also found the level of maternal FT4 was not associated with ADHD in children.

Six studies investigated the association of maternal TSH level with offspring ADHD. Chevrier et al (30) found doubling in TSH levels was associated with a 0.65-point decrease (95% CI = -1.26 to -0.04) on the ADHD subscale of the Child Behavior Checklist (CBCL), but there was no significant association between maternal TSH level and child's performance on the Conner's Kiddie Continuous Performance Test. Ghassabian et al (35) reported that per SD increase in plasma level of TSH was associated with higher CBCL externalizing (ADHD) scores. On the other hand, Modesto et al (39) found null association between maternal TSH level and offspring scores on ADHD using Conner's Parent Rating Scale-Revised Short Form. Three studies (32, 40, 41) fitted maternal TSH level as continuous variables and included offspring ADHD as a binary outcome. Fetene et al (32) observed null association between

Table 4. Estimated Associations Between Maternal Hypothyroxinemia and Offspring Neurodevelopmental Disorders

Item	Author (Y)	Definition of Maternal Hypothyroxinemia	Outcome Measurement	Estimate (95% CI)
<i>ADHD as outcome</i>				
1	Modesto et al (2015)	TSH levels of 0.1 to 2.5 mIU/L and FT4 levels below the 5th percentile of the sample	Conner's Parent Rating Scale- Revised Short Form	$\beta = 0.07$ (0.003-0.14)
2	Oostenbroek et al (2017)	FT4 level below the 10th or 5th percentile of distribution in the maternal FT4 level	Strengths and Difficulties Questionnaire	Teacher-reported: OR _{FT4 < 10th percentile} = 1.47 (0.99, 2.20) OR _{FT4 < 5th percentile} = 1.70 (1.01, 2.86) Parent-reported: OR _{FT4 < 10th percentile} = 0.85 (0.50, 1.46) OR _{FT4 < 5th percentile} = 0.78 (0.36-1.66)
3	Vermiglio et al (2004)	Normal TSH concentrations (0.4-4.0 μ U/mL) with low serum FT4 value as compared with the range values calculated at the same stage of pregnancy	DMS-IV-TR items by clinical observation	$\chi^2 = 2.34$, $P = 0.01$
<i>ASD as outcome</i>				
1	Roman et al (2013)	Mild: normal TSH levels of 0.03 to 2.5 mIU/L and FT4 levels below the 10 th percentile of the sample (< 11.82 pmol/L); severe: FT4 levels below the 5th percentile of the sample (< 10.99 pmol/L)	Pervasive Developmental Problems subscale from CBCL	OR _{FT4 < 10th percentile} = 1.41 (0.78, 2.57) OR _{FT4 < 5th percentile} = 2.60 (1.30-5.18)
<i>Schizophrenia as outcome</i>				
1	Gyllenberg et al (2016)	TSH levels over 5th to 95th percentile and FT4 levels below 10th percentile of the sample	ICD-10 diagnosis	OR = 1.75 (1.22-2.50)

Abbreviations: ADHD, attention deficit hyperactivity disorder; ASD, autism spectrum disorders; CBCL, Child Behavior Checklists; DMS-IV-TR, Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition-Text Revision; FT4, free T4.

maternal TSH level and offspring ADHD. Similarly, Oostenbroek et al (40) observed null association between maternal TSH level and offspring ADHD, either reported by teachers or by parents. However, Pakkila et al (41) found an association for each log increase in maternal TSH with ADHD symptoms in both boys (OR: 1.17, 95% CI: 1.00-1.36) and girls (OR: 1.39, 95% CI: 1.07-1.80; Table 5).

Only 1 study was conducted for the following comparisons between maternal thyroid hormone levels and offspring neurodevelopmental disorders. Roman et al (43) investigated the association of maternal FT4 and TSH levels with offspring ASD while null association was observed. Fetene et al (31) was the only study investigating offspring depression as a clinical outcome. They reported a null association of maternal TPO-Ab⁺ and TSH levels with offspring depression. Meanwhile, a significant association of maternal FT4 level with higher odds of depression in offspring was observed (OR: 1.21, 95% CI: 1.00-1.47). Gyllenberg et al (28) investigated the association of maternal FT4 and TSH levels with offspring schizophrenia. They found the odds of

schizophrenia decreased by 54% per log unit increase of maternal FT4 (95% CI: 0.31-0.94), but the association between log-transformed level of TSH and schizophrenia was not significant (Table 5).

Non-neurodevelopmental outcomes (N = 8)

Because most of the non-neurodevelopmental outcomes were observed in 1 study (Supplementary Table 2, <https://osf.io/k9jpf> (23)), meta-analysis was not conducted, and the individual studies were described in Supplementary Text (<https://osf.io/k9jpf> (23)).

Quality assessment

Study quality assessments were presented in Appendix 2 (<https://osf.io/k9jpf> (23)). Most studies (accounted for 60% of total included studies) were of good quality except for 9 studies. The overall quality of these nine studies was low because of the small sample size, poor identification of exposure and outcomes, or they did not adjust for any potential covariates in the model. Six of the 9 studies were not included

Table 5. Estimated Associations Between Maternal Thyroid Hormone Parameters and Offspring Neurodevelopmental Disorders

Item	Author (Y)	Outcome Measurement	Estimate (95% CI)
Maternal TPO-Ab⁺			
<i>ADHD as outcome</i>			
1	Fetene et al (2018)	DAWBA	OR = 1.00 (0.84-1.20) per SD
2	Ghassabian et al. (2012)	CBCL (Maternal and paternal rating)	OR = 1.77 (1.15-2.72), TPO-Ab ⁺ defined as >100 IU/mL
<i>ASD as outcome</i>			
1	Brown et al (2015) ^a	ICD-10 diagnosis	OR = 1.78 (1.16-2.75), TP-Ab ⁺ defined as >156 IU/mL
2	Roman et al (2013) ^b	Pervasive Developmental Problems subscale from CBCL	OR = 0.78 (0.28-2.16), TPO-Ab ⁺ defined as >100 IU/mL
<i>Depression as outcome</i>			
1	Fetene et al (2019)	DAWBA	OR = 1.02 (0.85-1.23) per SD
Maternal FT4 level			
<i>ADHD as outcome</i>			
1	Chevrier et al (2011)	CBCL KCPT	β = -0.10 (-2.03 to 1.82) per 1 ng/dL β = 7.52 (-4.86 to 19.91) per 1 ng/dL
2	Modesto et al (2015)	Conner's Parent Rating Scale-Revised Short Form	β = -0.01 (-0.02 to 0.01) per SD
3	Fetene et al (2018)	DAWBA	OR = 1.10 (0.89 to 1.36) per SD
4	Oostenbroek et al (2017)	SDQ	Teacher-reported: OR = 0.95 (0.85-1.05) per 1 pmol/L Parent-reported: OR = 1.01 (0.90-1.13) per 1 pmol/L
<i>ASD as outcome</i>			
1	Roman et al (2013)	Pervasive Developmental Problems subscale from CBCL	OR = 0.95 (0.77-1.17) per SD
<i>Depression as outcome</i>			
1	Fetene et al (2019)	DAWBA	OR = 1.21 (1.00-1.47) per SD
<i>Schizophrenia as outcome</i>			
1	Gyllenberg et al (2016)	ICD-10 diagnosis	OR = 0.54 (0.31-0.94) per LN unit
Maternal TSH level			
<i>ADHD as outcome</i>			
1	Chevrier et al (2011)	CBCL KCPT	β = -0.65 (-1.26 to -0.04) per LN unit β = -0.75 (-4.61 to 3.12) per LN unit
2	Ghassabian et al (2011)	CBCL	β = 0.08 (0.01-0.15) per SD
3	Modesto et al (2015)	Conner's Parent Rating Scale-Revised Short Form	β = -0.01 (-0.02 to 0.01) per SD
4	Fetene et al (2018)	DAWBA	OR = 0.83 (0.37-1.85) per SD
5	Oostenbroek et al (2017)	SDQ	Teacher-reported: OR = 1.02 (0.96-1.09) per mU/L Parent-reported: OR = 1.01 (0.92-1.11) per mU/L
6	Pakkila et al (2014)	Rutter B2 Scale-Finnish version	OR _{boys} = 1.17 (1.00-1.36) per LN unit OR _{girls} = 1.39 (1.07-1.80) per LN unit
<i>ASD as outcome</i>			
1	Roman et al (2013)	Pervasive Developmental Problems subscale from CBCL	OR = 0.92 (0.72-1.17) per SD
<i>Depression as outcome</i>			
1	Fetene et al (2019)	DAWBA	OR = 1.14 (0.98-1.32) per SD
<i>Schizophrenia as outcome</i>			
1	Gyllenberg et al (2016)	ICD-10 diagnosis	OR = 0.93 (0.84-1.02) per LN unit

^aThe cutoff point for TPO-Ab⁺ corresponded to serum concentrations >156 IU/mL.^bThe cutoff point for TPO-Ab⁺ corresponded to plasma concentrations $\geq$ 100 IU/mL.

Abbreviations: ADHD, attention deficit hyperactivity disorder; ASD, autism spectrum disorders; CBCL, Child Behavior Checklist; DAWBA: Developmental and Well-Being Assessment; FT4, free T4; SDQ: Strengths and Difficulties Questionnaire; TPO-Ab: thyroid peroxidase antibodies.





a. ADHD**b. ASD****c. epilepsy****d. schizophrenia**

Figure 2. Meta-analysis of studies on the association between maternal hyperthyroidism and (a) ADHD, (b) ASD, (c) epilepsy, and (d) schizophrenia in the offspring. ADHD, attention deficit hyperactivity disorder; ASD, autism spectrum disorder; CI, confidence interval; SE, standard error.

in meta-analysis. For the remaining 3, we performed additional sensitivity analyses by removing lower quality studies in the meta-analysis. An exception is the association analysis between maternal TPO-Ab⁺ and offspring ASD where only 2 studies were included in the meta-analysis, indicating cautious interpretation may be required. Similar results were observed in all the sensitivity analyses when compared with the original analysis without excluding lower quality studies (Fig. 3).

Discussion

This is a comprehensive systematic review with meta-analysis providing evidence for the association between maternal thyroid status and various offspring clinical outcomes. In general, maternal hyperthyroidism during pregnancy was significantly associated with increased risk of

offspring ADHD, epilepsy, and hyperthyroidism. Whereas maternal hypothyroidism during pregnancy was significantly associated with increased risk of ADHD, ASD, epilepsy, and hypothyroidism. These findings suggest that either maternal hyper- or hypothyroidism could affect neurodevelopment and thyroid health in the offspring.

Maternal thyroid dysfunction and neurodevelopmental disorders in offspring

Most studies included in this systematic review and meta-analysis studied maternal hypothyroidism as exposure. Although not a highly prevalent condition during pregnancy, maternal hypothyroidism is known to be associated with increased risk of spontaneous miscarriage, stillbirth, and perinatal death. Regarding the ascertainment of thyroid





a. ADHD**i.)****ii.) removing Haddow 1999 et al.****b. ASD****i.)****ii.) removing George 2014 et al.**

Figure 3. Meta-analysis of studies on the association between maternal hypothyroidism and (a) ADHD, (b) ASD, (c) epilepsy, (d) schizophrenia, and (e) cerebral palsy in the offspring. Sensitivity analysis was conducted by (i) including or (ii) removing the comparatively lower quality study. ADHD, attention deficit hyperactivity disorder; ASD, autism spectrum disorder; CI, confidence interval; SE, standard error.

status, 13 of 22 studies measured maternal exposure using thyroid function test, whereas the other 9 studies used clinical diagnosis or prescription records from hospitals. The majority of the studies used laboratory tests to measure maternal thyroid levels during the first and early second

trimesters. Notably, most studies available did not adjust for treatment status, whereas this potential factor can play a role in the association. A recently published article by Rotem et al (46), which is out of our investigation period, examined the association between maternal thyroid disorders





c. epilepsy**d. schizophrenia****e. cerebral palsy**

Figure 3. Continued.

Figure 4. Meta-analysis of studies on the association between maternal TPO-Ab⁺ and offspring ASD. ASD, autism spectrum disorder; TPO-Ab, thyroid peroxidase positive antibody.

and the risk of ASD in offspring, taking into account gestational thyroid hormone level and thyroid medications use. They found that maternal thyroid disorders, especially hypothyroidism, were associated with increased ASD risk in the offspring. This observation is indeed in line with our meta-analysis, showing hypothyroidism was significantly associated with offspring ASD. Nevertheless, they proposed that the association between maternal thyroid status and offspring ASD was independent of maternal gestational thyroid hormones and unaffected by medication treatment. By including a more comprehensive range of confounding factors and biologically relevant covariates (eg, thyroxine treatment), future studies are warranted to revisit the association between maternal thyroid dysfunction and various health outcomes in offspring. Together with larger sample

size and longer follow-up time, future work may particularly focus on the underlying mechanisms, which is largely unknown.

Notably, in the setting of register-based study cohorts using diagnosis codes or prescription records for exposure measurement, most included studies were unable to identify individuals with subclinical hypothyroidism. Only 3 studies investigated the association between maternal subclinical hypothyroidism and clinical outcomes in offspring while null associations were reported. The current data did not show a consistent association between subclinical hypothyroidism during pregnancy and adverse pregnancy outcomes (8, 47-49), but this was observed in a limited number of studies with small sample size. In fact, subclinical hypothyroidism is more common than overt hypothyroidism

(8, 50) Future studies are required to examine individuals with subclinical hypothyroidism. On the other hand, we observed in this review that there was some evidence of association between maternal hypothyroxinemia and offspring ADHD, ASD, and schizophrenia. To date, treatment for thyroid dysfunction during pregnancy is still a controversial issue (48, 49). The potential risks of maternal hypothyroxinemia on the offspring unraveled in this review may raise the awareness of the need to devise an optimum clinical management strategy for pregnant women, such as levothyroxine treatment.

The most commonly studied outcome is neurodevelopmental disorders in offspring, and there is a plausible explanation of the impact of maternal thyroid status during the first and early stages of second trimesters on the neurodevelopment in offspring (51-53). The fetal thyroid heavily depends on the maternal supply of thyroid hormone before the onset of thyroid function in the fetus which occurs around 16 to 20 weeks of gestation, and there is a significant transfer of thyroid hormones from the mother to the fetus (10, 12). The supply of thyroid hormone to the fetus leads to increased FT4 and decreased TSH concentrations from around the eighth week until the first half of pregnancy in the pregnant mothers, resulting in different reference intervals for TSH and FT4 when compared with the nonpregnant status (54). Besides, Oostenbroek et al (40) investigated maternal thyroid hormone levels throughout the whole duration of pregnancy. They partially confirmed early disruption in maternal thyroid function may be associated with ADHD but found a null association with other behavioral problems. In this systematic review and meta-analysis, we provided evidence that maternal hyper- and hypothyroidism are associated with increased risk of a range of neurodevelopmental disorders, suggesting that the impact of maternal thyroid status may not be limited to ADHD only in offspring.

The impact of maternal thyroid status after the early stage of the second trimester is less frequently studied. Chevrier et al (30) was the only one who measured maternal thyroid hormone during the third trimester. They reported a higher maternal TSH level was associated with a lower risk of having ADHD symptoms in offspring. Some studies identified a positive association between higher maternal thyroid hormone levels in the first 2 trimesters and increased risk of ADHD in offspring (35, 41). This indicated a balanced level of maternal thyroid hormone at each stage of pregnancy would be optimal for fetus neurodevelopment. However, the presence of abnormal thyroid status in the third trimester could be a marker of abnormal thyroid status in earlier trimesters. Nevertheless, the effect of timing of maternal thyroid function measurement requires further studies.

Strength and limitations

Previously, 3 published systematic reviews investigated the association between maternal thyroid hormone abnormalities and the risk of neurodevelopmental disorders in the offspring (55-57). Two systematic reviews examined the relationship between maternal thyroid dysfunction and psychiatric disorders (mainly focused on ADHD) (58, 59). Compared with these previous reviews, our study had a wider scope in respect of both the exposure and the outcome. Moreover, we only included outcomes that are clinically diagnosed or ascertained by well-established measurement tools. The setting of our inclusion criteria is more likely to produce consistent and reliable conclusions.

Nevertheless, there are limitations. First, we observed marginally significant association in the meta-analysis, which could be due to the limited number of studies, and the small sample size in each study, echoing the presence of the research gap and arousing the need to have more studies in this important but underexplored area. Cautious interpretation is also required in the current meta-analysis. Second, the definitions of thyroid dysfunction were heterogeneous across studies because there was no consensus in the definition of thyroid dysfunction during pregnancy at the time of this current study. Third, the definition of the clinical outcomes was not standardized in the included studies. Although we only included outcomes that were clinically diagnosed or assessed by validated tools, 10 of 22 studies that examined neurodevelopmental outcomes in offspring used behavior checklist rather than diagnosis or prescription records to identify the outcomes. The questionnaires adopted were reported by parents (30-32, 35, 36, 39-41, 43), teachers (40), self-report (31, 32), or clinical observation (44), leading to the discrepancy of the estimates. Fourth, the timing for the measurement of offspring outcomes plays a key role in revealing the association. Outcomes such as schizophrenia and depression usually develop at a late stage in childhood or even adolescence. Thus, a long follow-up time is essential to detect the potential association. Fifth, clinical outcomes ascertained using the diagnosis code or prescription records could lead to overestimation by misclassification. Because these limitations may increase heterogeneity between studies, cautious interpretation is required regarding the findings of meta-analyses with high heterogeneity.

Clinical interpretations

Although measurement of maternal thyroid hormone levels during pregnancy can be used for determining maternal thyroid status, current guidelines from the American Thyroid Association and the European Thyroid Association differ in

their recommendations regarding the screening of thyroid hormone during pregnancy (2, 50). The current systematic review and meta-analysis highlighted the potential association between maternal thyroid dysfunction during pregnancy and various disorders in the offspring, suggesting the importance of prevention during pregnancy in maintaining the health of the offspring. To cater for fetal iodine requirement, increase in maternal thyroid hormone production and renal iodine clearance, there is an elevated iodine requirement during pregnancy. Having sufficient iodine intake during pregnancy is one of the effective preventive measures of maternal thyroid dysfunction. In cases of pregnant mothers who are diagnosed with abnormal thyroid function, they should be promptly treated and carefully monitored throughout pregnancy to avoid hyper- and hypothyroidism. Regarding the thyroid hormone supplementation during pregnancy, a recent systematic review and meta-analysis (60) of randomized controlled trials (RCTs) and retrospective studies including 7970 patients demonstrated that LT4 supplementation had beneficial effects in pregnant women with subclinical hypothyroidism and/or thyroid autoimmunity by reducing their risks of pregnancy loss and preterm birth. Consistent results were obtained by subgroup analysis of RCTs. Depending on the types of pregnancy (naturally conceived pregnancies vs pregnancy achieved by assisted reproduction technology [such as in vitro fertilization and intracytoplasmic sperm injection]), LT4 supplementation may reduce the risk of pregnancy loss and/or preterm birth better in women with thyroid autoimmunity or subclinical hypothyroidism. On the other hand, a meta-analysis (61) of RCTs demonstrated that euthyroid women with individualized initial dosages of LT4 had lower risk of miscarriage, whereas those with fixed dosages of LT4 showed no significant difference. In addition, euthyroid women with thyroid autoimmunity given initial LT4 treatment in early pregnancy had a lower risk of preterm birth when compared with those receiving placebo or no treatment. According to the findings discussed here, detailed treatment strategies (such as timing and dosage of initial treatment) should be developed for pregnant women based on their types of pregnancy and thyroid status. After such strategies are developed, individualized therapeutic strategy of LT4 supplementation might be encouraged for pregnant women with euthyroidism, subclinical hypothyroidism or thyroid autoimmunity. Nevertheless, the number of RCTs to date is limited, and the sample size of available studies are small. Future large RCTs are warranted in this area.

Conclusion

Our review suggested that maternal thyroid dysfunction during pregnancy is a potential risk factor for

neurodevelopmental disorders (including ADHD, ASD, epilepsy), other cardiometabolic and respiratory conditions, and thyroid dysfunction in offspring during childhood. Currently, the optimum management strategy of maternal thyroid dysfunction during pregnancy is still largely unexplored. In view of these findings, routine measurement of thyroid function test during early pregnancy should be considered for all pregnant women. Further studies are warranted to delineate the association of thyroid status, especially levels of anti-TPO antibodies, in pregnancy and fetal outcome.

Acknowledgments

Author Contributions: C.L.C. contributed to the conception of the work and designed the study. G.M.G. and M.T.Y.L. decided on the inclusion and exclusion of studies and conducted data extraction. G.M.G. performed the meta-analysis, wrote the first draft of the manuscript, had full access to all the data in the study, and takes responsibility for the integrity of the data and the accuracy of the data analysis. K.K.C.M., W.C.L., P.I., G.H.Y.L., I.C.K.W., and A.W.C.K. interpreted, critically evaluated, and improved the study design and manuscript, and shared the responsibility for the final manuscript and the decision to submit.

Additional Information

Correspondence and Reprint Requests: Ching-Lung Cheung, Department of Pharmacology and Pharmacy, 2/F, 21 Sassoon Road, Li Ka Shing Faculty of Medicine, Laboratory Block, Hong Kong SAR, China. E-mail: lung1212@hku.hk.

Disclosure Summary: K.K.C.M. is the recipient of the CW Maplethorpe Fellowship and personal fees from IQVIA Ltd., unrelated to the submitted work. P.I. has received funding from the Hong Kong Research Grants Council, Health & Medical Research Fund, and Hong Kong Jockey Club Charities Trust, unrelated to the submitted work. I.C.K.W. reports research funding outside the submitted work from Amgen, Bristol-Myers Squibb, Pfizer, Janssen, Bayer, GSK, Novartis, the Hong Kong RGC, and the Hong Kong Health and Medical Research Fund, National Institute for Health Research in England, European Commission, National Health and Medical Research Council in Australia, and also received speaker fees from Janssen and Medicine in the previous 3 years, unrelated to the submitted work. The other authors declare no conflict of interest.

Data Availability: Data sharing is not applicable to this article as no datasets were generated or analyzed during the current study.

References

1. Carney LA, Quinlan JD, West JM. Thyroid disease in pregnancy. *Am Fam Physician*. 2014;89(4):273-278.
2. De Groot L, Abalovich M, Alexander EK, et al. Management of thyroid dysfunction during pregnancy and postpartum: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab*. 2012;97(8):2543-2565.
3. Mestman JH. Hyperthyroidism in pregnancy. *Curr Opin Endocrinol Diabetes Obes*. 2012;19(5):394-401.

4. Stagnaro-Green A, Pearce E. Thyroid disorders in pregnancy. *Nat Rev Endocrinol*. 2012;8(11):650-658.
5. Marx H, Amin P, Lazarus JH. Hyperthyroidism and pregnancy. *BMJ*. 2008;336(7645):663-667.
6. Lazarus JH. Thyroid function in pregnancy. *Br Med Bull*. 2011;97:137-148.
7. Gärtner R. Thyroid diseases in pregnancy. *Curr Opin Obstet Gynecol*. 2009;21(6):501-507.
8. Alexander EK, Pearce EN, Brent GA, et al. 2017 Guidelines of the American Thyroid Association for the diagnosis and management of thyroid disease during pregnancy and the postpartum. *Thyroid*. 2017;27(3):315-389.
9. Lazarus JH. Thyroid dysfunction in pregnancy: offspring effects. *Lancet Diabetes Endocrinol*. 2013;1(3):174-175.
10. Moog NK, Entringer S, Heim C, Wadhwa PD, Kathmann N, Buss C. Influence of maternal thyroid hormones during gestation on fetal brain development. *Neuroscience*. 2017;342:68-100.
11. Pop VJ, de Vries E, van Baar AL, et al. Maternal thyroid peroxidase antibodies during pregnancy: a marker of impaired child development? *J Clin Endocrinol Metab*. 1995;80(12):3561-3566.
12. Sahay RK, Nagesh VS. Hypothyroidism in pregnancy. *Indian J Endocrinol Metab*. 2012;16(3):364-370.
13. Andersen SL, Carlé A, Karmisholt J, Pedersen IB, Andersen S. Mechanisms in endocrinology: neurodevelopmental disorders in children born to mothers with thyroid dysfunction: evidence of fetal programming? *Eur J Endocrinol*. 2017;177(1):R27-R36.
14. Jølvig LR, Nielsen J, Kesmodel US, Nielsen RG, Nørgård BM, Beck-Nielsen SS. Chronic diseases in the children of women with maternal thyroid dysfunction: a nationwide cohort study. *Clin Epidemiol*. 2018;10:1381-1390.
15. Andersen SL, Andersen S, Vestergaard P, Olsen J. Maternal thyroid function in early pregnancy and child neurodevelopmental disorders: a Danish nationwide case-cohort study. *Thyroid*. 2018;28(4):537-546.
16. Eshkoli T, Wainstock T, Sheiner E, Beharier O, Fraenkel M, Walfisch A. Maternal hypothyroidism during pregnancy and the risk of pediatric endocrine morbidity in the offspring. *Am J Perinatol*. 2019;36(9):975-980.
17. Pääkkilä F, Männistö T, Surcel HM, et al. Maternal thyroid dysfunction during pregnancy and thyroid function of her child in adolescence. *J Clin Endocrinol Metab*. 2013;98(3):965-972.
18. Grattan MJ, Thomas DS, Hornberger LK, Hamilton RM, Midodzi WK, Vohra S. Maternal hypothyroidism may be associated with CHD in offspring. *Cardiol Young*. 2015;25(7):1247-1253.
19. Heikkinen AL, Pääkkilä F, Hartikainen AL, Väärasmäki M, Männistö T, Suvanto E. Maternal thyroid antibodies associates with cardiometabolic risk factors in children at the age of 16. *J Clin Endocrinol Metab*. 2017;102(11):4184-4190.
20. Liu X, Andersen SL, Olsen J, et al. Maternal hypothyroidism in the perinatal period and childhood asthma in the offspring. *Allergy*. 2018;73(4):932-939.
21. Cowett RM, Stern L. Maternal thyroid status and the incidence of respiratory distress syndrome: evaluation of a proposed relationship. *Pediatrics*. 1975;55(4):497-499.
22. Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ*. 2009;339:b2535.
23. Ge GM. Maternal Thyroid Dysfunction and Adverse Outcomes in the Offspring. 2020. <https://osf.io/k9jpf>. Accessed July 14, 2020.
24. Andersen SL, Laurberg P, Wu CS, Olsen J. Maternal thyroid dysfunction and risk of seizure in the child: a Danish nationwide cohort study. *J Pregnancy*. 2013;2013:636705.
25. Andersen SL, Laurberg P, Wu CS, Olsen J. Attention deficit hyperactivity disorder and autism spectrum disorder in children born to mothers with thyroid dysfunction: a Danish nationwide cohort study. *BJOG*. 2014;121(11):1365-1374.
26. Brown AS, Surcel HM, Hinkka-Yli-Salomäki S, Cheslack-Postava K, Bao Y, Sourander A. Maternal thyroid auto-antibody and elevated risk of autism in a national birth cohort. *Prog Neuropsychopharmacol Biol Psychiatry*. 2015;57:86-92.
27. George B, Padmam MS, Nair MK, Leena ML, Russell PS. CDC Kerala 13: antenatal, natal and postnatal factors among children (2-6 y) with autism—a case control study. *Indian J Pediatr*. 2014;81(Suppl 2):S133-S137.
28. Gyllenberg D, Sourander A, Surcel HM, Hinkka-Yli-Salomäki S, McKeague IW, Brown AS. Hypothyroxinemia during gestation and offspring schizophrenia in a national birth cohort. *Biol Psychiatry*. 2016;79(12):962-970.
29. Instanes JT, Halmøy A, Engeland A, Haavik J, Furu K, Klungsoyr K. Attention-deficit/hyperactivity disorder in offspring of mothers with inflammatory and immune system diseases. *Biol Psychiatry*. 2017;81(5):452-459.
30. Chevrier J, Harley KG, Kogut K, Holland N, Johnson C, Eskenazi B. Maternal thyroid function during the second half of pregnancy and child neurodevelopment at 6, 12, 24, and 60 months of age. *J Thyroid Res*. 2011;2011:426427.
31. Fetene DM, Betts KS, Alati R. The role of maternal prenatal thyroid function on offspring depression: findings from the ALSPAC cohort. *Dev Psychopathol*. 2020;32(1):189-196.
32. Fetene DM, Betts KS, Alati R. Maternal prenatal thyroid function and offspring ADHD: findings from the ALSPAC cohort. *J Nerv Ment Dis*. 2018;206(11):859-864.
33. Frank AS, Lupattelli A, Brandlistuen RE, Nordeng H. Maternal thyroid hormone replacement therapy exposure and language and communication skills of offspring at 8 years of age. *JAMA Netw Open*. 2019;2(10):e1912424.
34. Getahun D, Jacobsen SJ, Fassett MJ, et al. Association between maternal hypothyroidism and autism spectrum disorders in children. *Pediatr Res*. 2018;83(3):580-588.
35. Ghassabian A, Bongers-Schokking JJ, Henrichs J, et al. Maternal thyroid function during pregnancy and behavioral problems in the offspring: the generation R study. *Pediatr Res*. 2011;69(5 Pt 1):454-459.
36. Ghassabian A, Bongers-Schokking JJ, de Rijke YB, et al. Maternal thyroid autoimmunity during pregnancy and the risk of attention deficit/hyperactivity problems in children: the Generation R Study. *Thyroid*. 2012;22(2):178-186.
37. Gutvirtz G, Walfisch A, Wainstock T, Landau D, Sheiner E. Maternal hypothyroidism and future pediatric neurological morbidity of the offspring. *Arch Gynecol Obstet*. 2019;299(4):975-981.
38. Haddow JE, Palomaki GE, Allan WC, et al. Maternal thyroid deficiency during pregnancy and subsequent neuropsychological development of the child. *N Engl J Med*. 1999;341(8):549-555.

39. Modesto T, Tiemeier H, Peeters RP, et al. Maternal mild thyroid hormone insufficiency in early pregnancy and attention-deficit/hyperactivity disorder symptoms in children. *JAMA Pediatr.* 2015;169(9):838-845.
40. Oostenbroek MHW, Kersten RHJ, Tros B, Kunst AE, Vrijkotte TGM, Finken MJJ. Maternal hypothyroxinaemia in early pregnancy and problem behavior in 5-year-old offspring. *Psychoneuroendocrinology.* 2017;81:29-35.
41. Pääkilä F, Männistö T, Pouta A, et al. The impact of gestational thyroid hormone concentrations on ADHD symptoms of the child. *J Clin Endocrinol Metab.* 2014;99(1):E1-E8.
42. Petersen TG, Andersen AN, Uldall P, et al. Maternal thyroid disorder in pregnancy and risk of cerebral palsy in the child: a population-based cohort study. *BMC Pediatr.* 2018;18(1):181.
43. Román GC, Ghassabian A, Bongers-Schokking JJ, et al. Association of gestational maternal hypothyroxinemia and increased autism risk. *Ann Neurol.* 2013;74(5):733-742.
44. Vermiglio F, Lo Presti VP, Moleti M, et al. Attention deficit and hyperactivity disorders in the offspring of mothers exposed to mild-moderate iodine deficiency: a possible novel iodine deficiency disorder in developed countries. *J Clin Endocrinol Metab.* 2004;89(12):6054-6060.
45. Wasserman EE, Nelson K, Rose NR, et al. Maternal thyroid autoantibodies during the third trimester and hearing deficits in children: an epidemiologic assessment. *Am J Epidemiol.* 2008;167(6):701-710.
46. Rotem RS, Chodick G, Shalev V, et al. Maternal thyroid disorders and risk of autism spectrum disorder in progeny. *Epidemiology.* 2020;31(3):409-417.
47. Wiles KS, Jarvis S, Nelson-Piercy C. Are we overtreating subclinical hypothyroidism in pregnancy? *BMJ.* 2015;351:h4726.
48. Maraka S, Mwangi R, McCoy RG, et al. Thyroid hormone treatment among pregnant women with subclinical hypothyroidism: US national assessment. *Bmj.* 2017;356:i6865.
49. Casey BM, Thom EA, Peaceman AM, et al.; Eunice Kennedy Shriver National Institute of Child Health and Human Development Maternal-Fetal Medicine Units Network. Treatment of subclinical hypothyroidism or hypothyroxinemia in pregnancy. *N Engl J Med.* 2017;376(9):815-825.
50. Lazarus J, Brown RS, Daumerie C, Hubalewska-Dydejczyk A, Negro R, Vaidya B. 2014 European thyroid association guidelines for the management of subclinical hypothyroidism in pregnancy and in children. *Eur Thyroid J.* 2014;3(2):76-94.
51. Préau L, Fini JB, Morvan-Dubois G, Demeneix B. Thyroid hormone signaling during early neurogenesis and its significance as a vulnerable window for endocrine disruption. *Biochim Biophys Acta.* 2015;1849(2):112-121.
52. Glinioer D. The regulation of thyroid function in pregnancy: pathways of endocrine adaptation from physiology to pathology. *Endocr Rev.* 1997;18(3):404-433.
53. Krassas GE, Poppe K, Glinioer D. Thyroid function and human reproductive health. *Endocr Rev.* 2010;31(5):702-755.
54. Marco A, Vicente A, Castro E, et al. Patterns of iodine intake and urinary iodine concentrations during pregnancy and blood thyroid-stimulating hormone concentrations in the newborn progeny. *Thyroid.* 2010;20(11):1295-1299.
55. Fan X, Wu L. The impact of thyroid abnormalities during pregnancy on subsequent neuropsychological development of the offspring: a meta-analysis. *J Matern Fetal Neonatal Med.* 2016;29(24):3971-3976.
56. Wang P, Gao J, Zhao S, Guo Y, Wang Z, Qi F. Maternal thyroxine levels during pregnancy and outcomes of cognitive development in children. *Mol Neurobiol.* 2016;53(4):2241-2248.
57. Thompson W, Russell G, Baragwanath G, Matthews J, Vaidya B, Thompson-Coon J. Maternal thyroid hormone insufficiency during pregnancy and risk of neurodevelopmental disorders in offspring: a systematic review and meta-analysis. *Clin Endocrinol (Oxf).* 2018;88(4):575-584.
58. Fetene DM, Betts KS, Alati R. Mechanisms in Endocrinology: maternal thyroid dysfunction during pregnancy and behavioural and psychiatric disorders of children: a systematic review. *Eur J Endocrinol.* 2017;177(5):R261-R273.
59. Drover SSM, Villanger GD, Aase H, et al. Maternal thyroid function during pregnancy or neonatal thyroid function and attention deficit hyperactivity disorder: a systematic review. *Epidemiology.* 2019;30(1):130-144.
60. Rao M, Zeng Z, Zhou F, et al. Effect of levothyroxine supplementation on pregnancy loss and preterm birth in women with subclinical hypothyroidism and thyroid autoimmunity: a systematic review and meta-analysis. *Hum Reprod Update.* 2019;25(3):344-361.
61. Sun X, Hou N, Wang H, Ma L, Sun J, Liu Y. A meta-analysis of pregnancy outcomes with levothyroxine treatment in euthyroid women with thyroid autoimmunity. *J Clin Endocrinol Metab.* 2020;105(4):1009-1019.