

AOGS REVIEW ARTICLE

Pre-, peri- and neonatal risk factors for autism

VINCENT GUINCHAT¹, POUL THORSEN², CLAUDINE LAURENT^{1,3}, CHRISTINE CANS⁴, NICOLAS BODEAU¹ & DAVID COHEN¹

¹Department of Child and Adolescent Psychiatry, Pitié-Salpêtrière University Hospital, Paris, France, ²Department of Obstetrics and Gynecology, Lillebaelt Hospital, Kolding, Denmark, ³Biotechnology and Biotherapy Unit, Research Centre, Brain and Spinal Cord Institute, Pierre and Marie Curie University, INSERM UMR_S 975, CNRS UMR 7225, Pitié-Salpêtrière Hospital, Paris, France, and ⁴Register for Disabled Children and the Isère County Perinatal Survey, Grenoble, France

Key words

Autism, prenatal, perinatal, neonatal, pervasive developmental disorders, risk factor, pregnancy

Correspondence

Dr Vincent Guinchat, Service de psychiatrie de l'enfant et de l'adolescent, Groupe Hospitalier Pitié-Salpêtrière, 47–83 Boulevard de l'hôpital, 75013 Paris, France.
E-mail: vincent.guinchat@psl.aphp.fr

Conflict of interest

The authors have stated explicitly that there are no conflicts of interest in connection with this article.

Please cite this article as: Guinchat V, Thorsen P, Laurent C, Cans C, Bodeau N, Cohen D. Pre-, peri- and neonatal risk factors for autism. Acta Obstet Gynecol Scand 2012; 91:287–300.

Received: 9 September 2011

Accepted: 4 November 2011

DOI: 10.1111/j.1600-0412.2011.01325.x

Abstract

Objective. To identify pre-, peri- and neonatal risk factors for pervasive developmental disorders (PDD). **Methods.** We searched the Medline database through March 2011 for relevant case-control and population-based studies on pre-, peri- and neonatal hazards related to PDD, including autism. We identified 85 studies for this review. Data were extracted systematically and organized according to risk factors related to family history, pregnancy, gestational age, delivery, birth milestones and the neonate's condition at birth. **Results.** During the prenatal period, risk factors for PDD were advanced maternal or paternal ages, being firstborn vs. third or later, maternal prenatal medication use and mother's status as foreign born. During the perinatal and neonatal periods, the risk factors for PDD were preterm birth, breech presentation, planned cesarean section, low Apgar scores, hyperbilirubinemia, birth defect and a birthweight small for gestational age. The influence of maternal pre-eclampsia, diabetes, vomiting, infections and stress during pregnancy requires further study in order to determine risk for PDD. **Discussion.** Despite evidence for the association of some pre-, peri- and neonatal risk factors associated with PDD, it remains unclear whether these risks are causal or play a secondary role in shaping clinical expression in individuals with genetic vulnerability. A plausible hypothesis is that improvements in obstetric and neonatal management have led to an increased rate of survivors with pre-existing brain damage. Given the variety of risk factors, we propose that future studies should investigate combinations of multiple factors, rather than focusing on a single factor.

Abbreviations: AOR, adjusted odd ratio; DSM, Diagnostic and Statistical Manual of Mental Disorders; ICD, International Classification of Diseases; IQ, intellectual quotient; LBW, low birthweight; PDD, pervasive developmental disorders.

Introduction

Autism and other pervasive developmental disorders (PDD) are common behavioral syndromes characterized by impairments in social interaction, abnormalities in verbal and non-verbal communication, and restricted and stereotyped interests and behaviors. Their onset occurs in early childhood and often results in severe lifelong impairments. In the USA, the Autism and Developmental Disabilities Monitoring Network reported an overall average prevalence of autism spectrum disorders in nine of 1000 children aged 8 years, with a 57%

increase in prevalence between 2002 and 2006 (1), which demonstrates that PDD is an urgent public health concern (1).

The first research on a link between complications during pregnancy and autism by Pasamanick et al. dates back to 1956, only a few years after the syndrome was first described (2). Since then, various conclusions have been suggested in studies on autism risk factors but have failed to clarify the relation between autism and adverse exposures during the pre-, peri- and neonatal periods. Increasing evidence also suggests a role of genetic factors in the origins of autism. Therefore, it remains unclear whether certain complications at birth are

causal, play a secondary role in shaping clinical expression in individuals with genetic vulnerability, or represent some of the shared causal factors in the development of PDD.

Numerous considerations surrounding PDD make research on pre-, peri- and neonatal factors worthwhile. Firstly, concordance in monozygotic twin pairs is incomplete, suggesting that nonheritable factors contribute to the risk of autism (3). For example, pregnancy-induced central nervous system insults may result in relevant epigenetic changes. Secondly, increasing evidence indicates that the prevalence of PDD has increased over the past 20 years at a rate not explained by improved detection of PDD in the population (4). This phenomenon raises the probability that environmental factors play a role (5). Thirdly, a growing body of literature suggests that histological and anatomical disturbances in the brain play an important role in the etiology of PDD (6,7). Such research suggests that, irrespective of the cause of these structural anomalies, the etiologically relevant period could be the early in utero stages, similar to auto-immune processes during pregnancy that lead to biological differences (8). Fourthly, the proportion of children with a major gene defect is limited to a small proportion of PDD cases. Thus, a multifactorial approach towards PDD risk may serve as a more appropriate perspective in the study of the genesis of autism. Additionally, these genetic anomalies appear not to be specific for autism but rather to share a role in the etiology of intellectual disability (9) and perhaps schizophrenia (10). An explanation for how two children with the same genetic vulnerability develop autism rather than intellectual disability or schizophrenia remains elusive. Finally, identification of environmental factors for autism during pregnancy carries clinical implications in terms of primary prevention.

To draw conclusions about the role of obstetric factors and the magnitude of their effect, analysis of the current data is warranted. Brasic and Holland (11) detailed a reliable procedure to identify case-control reports to be used for meta-analytic purposes. However, a review of 156 articles (11) yielded only two studies that fulfilled that defined set of criteria, and those two studies had discordant results (12,13). Klevzon *et al.* defined another set of criteria (14). Seven studies from various countries were selected. Four of the seven studies were prospective, population-based cohort studies; the others were retrospective. Three studies had a partially overlapping sample. The authors identified the following four categories of risk factors: advanced parental age; maternal place of birth outside Europe or North America; low birthweight (LBW) and preterm delivery; and intrapartum asphyxia. Recently, Gardener *et al.* published a systematic review and meta-analysis of 64 studies published prior to March 2007 (15). Although those studies covered a full scope of pregnancy and birth complications, the authors presented results on pregnancy-related risk factors only. Criteria for inclusion in their meta-analysis were less strict than those of previ-

ous studies (11,14). The selected studies did not always use operational criteria for PDD defined by International Classification of Diseases, Tenth Revision (ICD-10; World Health Organization, 1993) and the Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (DSM-IV; American Psychiatric Association, 1994; 16,17). Forty studies were eligible for that meta-analysis (15), and the authors provided information about relative risk level and confidence intervals. For each factor, a summary effect estimate was calculated and heterogeneity was examined. A meta-regression analysis accounted for methodological factors that contributed to study variability. The results of the Gardener study are incorporated into this review.

The present review of case-control studies aimed to estimate the differential impact of pre-, peri- and neonatal factors. Our goal was to identify which factors are relevant to a better understanding of the pathogenesis of PDD and their early detection.

Material and methods

Study selection

Relevant studies on pre-, peri- and neonatal hazards in autism were identified from the bibliographies of recent reviews (11,14,15). Additionally, a PubMed search was conducted through March 2011 using the following keywords: 'autistic disorder', 'Asperger's syndrome', 'prenatal', 'perinatal', 'neonatal', 'obstetric', 'risk' and 'familial'. A separate search was performed for specific variables, such as 'birth defect' and 'clinical severity'. We selected the case-control studies that explored pre-, peri- or neonatal risk factors. We excluded case reports, letters to the editor, animal models and experiments, genetic studies and studies on biomarkers. We did not include publications on the season of birth, and we determined that research on exposure to toxins during pregnancy would not be exhaustive. Using a similar set of inclusion criteria, our results matched results of the Gardener meta-analysis (15), with the addition of articles published from March 2007 through March 2011. From the body of research selected, we identified the well-designed, population-based studies by using Klevzon's criteria in addition to the reporting on adjusted odd ratios (AOR; 14). Flowcharts illustrating the inclusion and exclusion criteria are presented in Figure 1.

Data extraction

The following characteristics of each study were recorded: (a) study design, including a prospective vs. a retrospective approach and a description of the databases; (b) sample size and characteristics, including diagnosis classification, clinical assessment, inclusion and exclusion criteria; (c) a description of the control group (healthy controls, IQ matches, siblings); (d) a description of the risk factors, including the format of




Figure 1. Flowchart of the studies included in the present systematic review of pre-, peri- and neonatal risk factors of pervasive developmental disorders.

these risks and the presence of aggregated scores; (e) source of obstetric information (parental interview, medical record, birth certificate); and (f) statistically significant results with or without multivariate adjustment. For each variable, we dis-

tinguished all statistically significant results from nonsignificant results. After a brief analysis of the factors that could explain any inconsistency, we extracted the well-designed, population-based studies for which an independent risk

factor emerged in the adjusted analysis. Unfortunately, the AOR varied according to the design and purpose of each study. Adjustment could be performed with either all significant variables or only some of the well-defined, confounding variables, such as parental age, parity and family factors. A total of 85 studies were included (12,13,18–100).

Results

Prenatal risk factors

In Table 1 we summarize prenatal (familial and pregnancy) risk factors based on Gardener's meta-analysis (15) and on studies published since March 2007. The following four familial factors were consistently associated with autism: advanced maternal age and advanced paternal age, both as independent risk factors; primiparous women; and having a mother born outside Europe, North America or Australia.

Regarding pregnancy, the following three risk factors emerged from Gardener's meta-analysis: bleeding; medication during pregnancy; and diabetes. Recent data confirm the association of medication and bleeding but not diabetes with autism (83). Pre-eclampsia, vomiting, infection and stress during pregnancy still need to be studied more thoroughly. Except for paternal age (over 40 years) with an AOR of around 3, all other documented AORs were between 1.5 and 2.

Perinatal risk factors

Most authors used an empirical definition of the perinatal period that included time of delivery. The purpose of distinguishing this period from the prenatal period lies in the potential contribution of external, mechanical procedures and the consequences of birth complications to the fetus. Although the relation between perinatal and prenatal risk factors is not easy to delineate, complications during delivery could at the very least serve as the observable event of a final, causal pathway, thus revealing previously unnoticed prenatal insults or abnormal fetal development. We present the various perinatal risk factors that emerged in more than two publications and distinguish a pathological duration of pregnancy (preterm and post-term pregnancies) and delivery-related risk factors (Table 2). During the perinatal period, the predominant risk factors were preterm birth, breech presentation and planned cesarean section. All the documented AOR were between 1.3 and 2.8. Nevertheless, in a stratified analysis the AOR for preterm birth increased by as much as 5 in a group of girls born before 33 weeks (72).

Neonatal risk factors

Numerous neonatal factors were suspected and investigated as possible risk factors for autism. We classified them as follows: (a) LBW and size; (b) poor condition at birth, including low Apgar scores and hypoxia; and (c) other conditions, such

as hyperbilirubinemia, encephalopathy and birth defects. Table 3 lists the potential risks, including positive and negative findings, from studies of these potential risk factors. During the neonatal period, the risk factors for PDD were low Apgar scores, neonatal encephalopathy, hyperbilirubinemia, birth defect and baby small for gestational age. The documented AORs were below 3 for most variables, but between 3 and 5 for neonatal encephalopathy and up to 9 for hyperbilirubinemia in children born at term.

Low birthweight, usually defined as less than 2500 g, led to inconsistent results. Nevertheless, in a stratified analysis, Schendel *et al.* credited LBW with an independent twofold increased risk for autism in children born at term and a 7.1 risk for girls born at term (72). Moreover, four population-based studies reported an association between autism and being small for gestational age (usually defined as a weight at birth less than 2 SD below the expected weight on customized curves). No studies have yet explored intrauterine growth restriction.

Discussion

No individual factor in the neonatal and perinatal periods has been consistently validated as a risk factor for autism. However, some have been associated with autism in several studies and should be considered as potential risk factors that provide small contributions to the etiology or causal pathway of autism (Table 4). Although heterogeneous results implicate a variety of events, the differences observed in the optimality scores indicate that rather than focusing on a single factor, future studies should investigate combinations of factors.

Some of the risk factors listed in Table 4 warrant individual discussion. Firstly, the prevalence of cesarean sections has increased in recent decades, for both social and medical reasons. The current data do not enable us to untangle the true effect of cesarean section from the underlying indications to this operative procedure (such as failure to progress in labor, fetal distress, multiple pregnancies, breech presentation and the increasing trend of women requesting a cesarean section). It appears that if mechanical interventions in delivery serve as environmental factors, the magnitude of those effects is low. Progress in neonatal management has led to increased survival of preterm infants, but as a consequence of this a growing number of severe disabilities may be anticipated during childhood. The contribution of increasing survivors of extreme prematurity to the dramatically increasing prevalence of autism has to be questioned. Regarding prematurity, evidence that moderate preterm birth (34–37 weeks gestation) is an independent risk factor has only emerged in three studies and has carried a moderate effect magnitude. However, the risk seems to increase and to be more reliable in cases of more severe preterm birth. Similar observations can be made for LBW babies. However, it is difficult to

Table 1. Prenatal risk factors for autism as reported by Gardener (13) or in at least two recent studies (since March 2007) after adjusted analyses.

Risk factors	Positive studies (univariate analysis, since March 2007)	Negative/null studies (univariate analysis, since March 2007)	P/T ^a	Adjusted effect estimates ^b
Familial risk factors				
Advanced maternal age	Bilder (2009; 79), Buchmayer (2009; 83), Burstyn (2010; 91), Croen (2007; 64), Dawson (2009; 80), Durkin (2008; 76), Grether (2009; 84), Haglund (2011; 99), King (2009; 81), Mann (2009), Shelton (2010; 87), Williams (2008; 69)	Croen (2008; 73), Hultman (2011; 93), Karmel (2010; 92), Tsuchiya (2008; 74), Sasanfar (2010; 86), Zhang (2010; 95)	20/47	Gardener (2009; 15) (>35 vs. <35): 1.6 (1.32–1.95)
Advanced paternal age	Croen (2007; 64), Dawson (2009; 80), Durkin (2008; 76), Grether (2009; 84), Hultman (2011; 93), King (2009; 81), Sasanfar (2010; 86), Shelton (2010; 87), Tsuchiya (2008; 74), Williams (2008; 69), Zhang (2010; 95)		15/20	Gardener (2009; 15) (>40 vs. <30): 2.1 (1.48–2.86) Gardener (2009; 15) (35+ vs. 25–29): 1.34 (1.16–1.54) Gardener (2009; 15) (40+ vs. <30): 3.10 (0.95–9.49)
Mother born abroad	Buchmayer (2009; 83), Haglund (2011; 99), Williams (2008; 69), Hultman (2011; 93),		7/9	Gardener (2009; 15): 1.58 (1.14–2.19) Haglund (2011; 99): 2.2 (1.6–3.1)
Parity	Bilder (2009; 79), Croen (2008; 73), Dawson (2009; 80), Durkin (2008; 76), Hultman (2011; 93), Sasanfar (2010; 86), Zhang (2010; 95)	Buchmayer (2009; 83), Burstyn (2010; 91), Haglund (2011; 99)	15/30 ^c	Gardener (2009; 15): first vs. ≥ third 1.61 (1.42–1.82)
Pregnancy risk factors				
Diabetes		Buchmayer (2009; 83), Burstyn (2010; 91), Dodds (2011; 100)	1/9	Gardener (2009; 15): 2.07 (1.24–3.47)
Bleeding	Burstyn (2010; 91), Dodds (2011; 100)	Bilder (2009; 79), Burstyn (2010; 91),	8/21	Gardener (2009; 15): 1.81 (1.14–2.86)
Psychotropic drugs	Dodds (2011; 100)		6/16	Gardener (2009; 15): 1.46 (1.08–1.96)
Pre-eclampsia	Buchmayer (2009; 83), Burstyn (2010; 91), Dodds (2011; 100), Mann (2010; 90)		5/17	Buchmayer (2009; 83): 1.64 (1.08–2.49) Burstyn (2010; 91): 1.49 (1.00–2.23) Mann (2010; 90): 1.69 (1.26–2.28)

Other risk factors with at least one positive result reported since March 2007 and no more than one significant AOR (P/T) are as follows: hypertension/edema ($n=2/15$); infection (5/20); nausea (3/8^c); stress during pregnancy (3/4); Rhesus sensitivity (2/11); smoking during pregnancy ($n=2/10$); threatened abortion (3/4); and weight gain during pregnancy (2/6).

Other risk factors with no recent positive results and for which no statistically significant AOR was reported by Gardener are as follows (number of studies): fever ($n=5$); physical injury ($n=4$); anemia ($n=4$); proteinuria ($n=4$); any illness during pregnancy ($n=6$); placental complication ($n=9$); number of prenatal visits ($n=2$); 1+ prenatal complications ($n=2$); genitourinary tract infection ($n=3$); previous miscarriage ($n=40$); and previous fetal loss (abortion, miscarriage, stillbirth; $n=15$).

Risk factors not evoked by Gardener and examined in only single studies are as follows: work with computers (95); X-ray exposure (95); tocolytic therapy (95); malnutrition (95); oligoamnios (79); and ultrasound exposure (85).

^aStatistically significant positive results/total.

^bEffect estimates were Relative Risk and Odds Ratio. When significant, only Gardener's AOR is indicated for one risk factor.

^cIndicates that p -value includes positive/negative and/or mixed results.

Table 2. Perinatal risk factors for autism, as reported in at least two positive studies.

Risk factors	Positive studies (univariate analysis)	Negative or null studies (univariate analysis)	P/T ^a	Adjusted effect estimates ^b
Prematurity and postmaturity				
Preterm birth <37 weeks	Brimacombe (2007; 63), Buchmayer (2009; 83), Dodds (2011; 100), Durkin (2008; 76), Guillem (2006; 62), Finegan (1979; 21), Haglund (2011; 99), Hultman (2002; 44), Maimburg (2006; 61), Mann (2010, 90), Wilkerson (2002; 45), Williams (2008; 69)	Bilder (2009; 79), Burstyn (2010; 91), Cryan (1996; 13), Eaton (2001; 41), Juul-Dam (2001; 40), Lord (1991; 32), Laxer (1988; 28), Larsson (2005; 50), Schendel (2008; 72), Stein (2006; 57), Wier (2006; 59)	12/23	Buchmayer (2009; 83); ^c 1.44 (1.07–1.94) Durkin (2008; 76): 1.4 (1.2–1.17) Williams (2008; 69): 2.2 (1.5–3.5)
<35 weeks	Larsson (2005; 50), Zhang (2010, 95)	Eaton (2001; 41)	2/3	Larsson (2005; 50): 2.57 (1.64–4.03)
<33 weeks	Schendel (2008; 72), Eaton (2001; 41)	Wier (2006; 59), Karmel (2010; 92)	2/4	Schendel (2008; 72); ^d 5.4 (1.1–27.7)
<28 weeks <26 weeks	Durkin (2008; 76), Johnson (2010; 88)		1/1 1/1	Durkin (2008; 76): 2.8 (1.6–3.9)
Post-term	Cryan (1996; 13), (male) Laxer (1988; 28), Lobasher (1970; 16), Lord (1991; 32), Sugie (2005; 56), Zambrino (1997; 37)	Bryson (1988; 27), Finegan (1979; 21), Juul-Dam (2001; 40), Hultman (2002; 44), Larsson (2005; 50), Mason-Brothers (1990; 30), Maimburg (2008; 75), Stein (2006; 57), Zhang (2010, 95)	6/15	
Delivery-related risk factors				
Breech presentation	Bilder (2009; 79), Burstyn (2010; 91), Finegan (1979; 21), Larsson (2005; 50), Levy (1988; 29), Maimburg (2006; 61), Wilkerson (2002; 45)	Deykin (1980; 23), Eaton (2001; 41), Gillberg (1983; 12), Juul-Dam (2001; 40), Lord (1991; 32), Mason-Brothers (1990; 30), Matsuishi (1999; 39), Piven (1993; 34), Stein (2006; 57), Bryson (1988; 27)	7/17	Burstyn (2010; 91): 1.31 (1.02–1.69) Bilder (2009; 79): 2.10 (1.11–3.97) Larsson (2005; 50): 1.63 (1.18–2.26)
Induced labor	Dodds (2011; 100), Glasson (2004; 48), Juul-Dam (2001; 40)	Burstyn (2010; 91), Brimacombe (2007; 63), Laxer (1988; 28), Mason-Brothers (1990; 30), Maimburg (2008; 75), Stein (2006; 57)	3/9	Dodds (2011; 100): 1.22 (1.03–1.44)
Precipitous labor	Finegan (1979; 21), Juul-Dam (2001; 40)	Deykin (1980; 23), Glasson (2004; 48), Stein (2006; 57)	2/5	
Prolonged labor	Juul-Dam (2001; 40), Finegan (1979; 21), Eaton (2001; 41), Karmel (2010; 92), Brimacombe (2007; 63), Wilkerson (2002; 45)	Deykin (1980; 23), Dodds (2011; 100), Laxer (1988; 28), Lord (1991; 32), Mason-Brothers (1990; 30), Stein (2006; 57)	5/11	
Cesarean section, all indications	Bilder (2009; 79), Brimacombe (2007; 63), Dodds (2011; 100), Eaton (2001; 41), Hultman (2002; 44), Zhang (2010, 95)	Burstyn (2010; 91), Finegan (1979; 21), Gillberg (1983; 12), Laxer (1988; 28), Lord (1991; 32), Mason-Brothers (1990; 30), Matsuishi (1999; 39)	6/13	Hultman (2002; 44): 1.6 (1.1–2.3)
Scheduled cesarean	Burstyn (2010; 91), Glasson (2004; 48), Haglund (2011; 99), Maimburg (2008; 75),		4/4	Glasson (2004; 48): 1.83 (1.32–2.5) Burstyn (2010; 91): 1.23 (1.01–1.49)

Table 2. Continued.

Risk factors	Positive studies (univariate analysis)	Negative or null studies (univariate analysis)	P/T ^a	Adjusted effect estimates ^b
Meconium	Bryson (1988; 27), Matsuishi (1999; 39)	Bilder (2009; 79), Finegan (1976), Gillberg (1983; 12), Levy (1988; 29), Lord (1991; 32), Maimburg (2006; 61), Mason-Brothers (1990; 30), Piven (1993; 34)	2/10	
Fetal distress	Glasson (2004; 48), Hultman (2010; 93),	Bilder (2009; 79), Brimacombe (2007; 63), Sugie (2005; 56)	2/5	Glasson (2004; 48): 1.52 (1.12–2.06) Hultman (2010; 93): 1.44 (1.02–2.05)

Delivery-related risk factors for autism that resulted in one positive result are as follows: cephalo-pelvic disproportion (48); cord complication (12,18,20,21,22,28,29,30,32,34,39,40,48,57,95); epidural anesthesia (12,20,27,48); forceps (12,18,21,23,28,30,32,34,40,48,57,94,99); placental insufficiency (41); placental abruption (30,57); postpartum hemorrhage (22,29,39,48,61); premature rupture of membranes (30,57,48,95); preterm birth <31 weeks (100); and unscheduled cesarean (22,48,57,61,99).

Delivery-related risk factors for autism that resulted in no positive results are as follows: vacuum extraction (12,21,23,32,34,40,44,48,57,61); vaginal birth after previous cesarean section (79); acidosis with a pH <7.20 in cord blood (61); pathological fetal heart rate in labor (61); umbilical cord infection (57); placenta calcified (21); abnormal placenta (57); placenta previa (13,27); placental infarcts (28,83); infected amniotic fluid (57); puncture of fetal membranes (61); general anesthesia (21,32,48); anesthesia, nonspecific (27,34) and trauma during delivery (12,21,28,34,40,49,83); and preterm birth <36 weeks (38).

^aStatistically significant positive results/total.

^bEffect estimate were Relative Risk and Odds Ratio.

^cAdjusted for mother/pregnancy characteristics only.

^dGirls only.

Table 3. Neonatal risk factors for autism, as reported in at least two positive studies.

Risk factors	Positive studies (univariate analysis)	Negative or null studies (univariate analysis)	P/T ^a	Adjusted effect estimates ^b
Birthweight and growth				
Low birthweight	Burd (1999; 38), Burstyn (2010; 91), Croen (2002; 43), Deykin (1980; 23), Eaton (2001; 41), Finegan (1979; 21), Haglund (2011; 99), Hultman (2002; 44), Hultman (2010; 93), Karmel (2010; 92), Knobloch (1975; 19), Larsson (2005; 50), Maimburg (2006; 61), Schendel (2008; 72)	Bilder (2009; 79), Brimacombe (2007; 63), Croen (2005; 52), Dodds (2011; 100), Juul-Dam (2001; 40), Levy (1988; 29), Mason-Brothers (1990; 30), Mann (2010; 90), Wier (2006; 59), Williams (2008; 69), Stein (2006; 57), Zhang (2010; 95)	14/27	Burstyn (2010; 91): 1.3 (1.1–1.75) Maimburg (2006; 61): 3.0 (1.7–5.1) Schendel (2008; 72), ^{d,f} : 7.1 (1.6–32.6)
Lower birthweight	Burd (1999; 38), Karmel (2010; 92), Mann (2010; 90), Mason-Brothers (1990; 30), ^e Wilkerson (2002; 45)	Bryson (1988; 26), Cryan (1996; 13), Croen (2005; 52, 2007; 64, 2008; 73), Gillberg (1983; 12), Glasson (2004; 48), Links (1980; 25), Torrey (1975; 20), Lobasher (1970; 18), Lord (1991; 32), Sugie (2005; 56)	5/17	
Small for gestational age	Buchmayer (2009; 83), Eaton (2001; 41), Hultman (2002; 44), Hultman (2010; 93), Larsson (2005; 50)	Durkin (2008; 76), Dodds (2011; 100), Gillberg (1983; 12), Glasson (2004; 48), Haglund (2011; 99), Karmel (2010; 92), Mann (2009), Piven (1993; 34), Williams (2008; 69)	5/14	Buchmayer (2009; 83): 1.86 (1.32–2.63) Hultman (2002; 44): 2.1 (1.1–3.9) Larsson (2005; 50): 1.32 (1.0–1.68)

Table 3. Continued.

Risk factors	Positive studies (univariate analysis)	Negative or null studies (univariate analysis)	P/T ^a	Adjusted effect estimates ^b
Head circumference	Courchesne (2003; 46), Karmel (2010; 92)	Glasson (2004; 48), Hazlet (2005; 55), Hultman (2002; 44), Laxer (1988; 28), Mason-Brothers (1990; 30), Torrey (2004; 49)	2/8	
Poor condition at birth				
Suboptimal Apgar 5	Bryson (1988; 26), Buchmayer (2009; 83), Burstyn (2010; 91), Finegan (1979; 21), Hultman (2002; 44), Larsson (2005; 50), Maimburg (2006; 61)	Bilder (2009; 79), Bryson (1988; 26), Burd (1999; 38), Dodds (2011; 100), Eaton (2001; 41), Gillberg (1983; 12), Haglund (2011; 99), Torrey (1975; 20), Juul-Dam (2001; 40), Karmel (2010; 92), Levy (1988; 29), Maimburg (2008; 75), Mason-Brothers (1990; 30), Matsuishi (1999; 39), Piven (1993; 34), Williams (2008; 69)	7/23	Larsson (2005; 50): 1.97 (1.15–3.36) Hultman (2002; 44): 3.2 (1.2–8.2)
Suboptimal Apgar 1	Finegan (1979; 21), Glasson (2004; 48), Williams (2008; 69)	Burd (1999; 38), Eaton (2001; 41), Torrey (1975; 20), Karmel (2010; 92), Levy (1988; 29), Maimburg (2008; 75), Mason-Brothers (1990; 30), Matsuishi (2005)	3/11	
Transfer to special care	Guillem (2006; 62), Deykin (1980; 23), Maimburg (2006; 61), Matsuishi (2005)	Glasson (2004; 48), Levy (1988; 29), Mason-Brothers (1990; 30)	4/7	Maimburg (2006; 61): 1.8 (1.3–2.7)
Markers of hypoxia				
Lack of first cry, breath or oxygen; blue baby	Brimacombe (2007; 63), Bryson (1988; 27), Glasson (2004; 48), Stein (2006; 57), Zhang (2010; 95)	Deykin (1980; 23), Laxer (1988; 28), Matsuishi (2005), Stein (2006; 57)	5/9	
Respiratory distress syndrome or assisted ventilation or asphyxia	Bryson (1988; 27), Buchmayer (2009; 83), Dodds (2011; 100), Finegan (1979; 21), Juul-Dam (2001; 40), Gillberg (1983; 12), Sugie (2005; 56)	Bilder (2009; 79), Laxer (1988; 29), Lord (1991; 32), Maimburg (2006; 61), Mason-Brothers (1990; 30), Matsuishi (1999; 39), Piven (1993; 34), Sugie (2005; 56), Williams (2008; 69)	7/17	
Other specific conditions at birth				
Hyperbilirubinemia	Buchmayer (2009; 83), Finegan (1979; 21), Juul-Dam (2001; 40), Maimburg (2008; 75), Maimburg (2010; 94), Sugie (2005; 56), Zhang (2010; 95)	Brimacombe (2007; 63), Bryson (1988; 27), Croen (2005; 52), Deykin (1980; 23), Laxer (1988; 28), Lobasher (1970; 18), Lord (1991; 32), Mason-Brothers (1990; 30), Matsuishi (2005), Williams (2008; 69), Piven (1993; 34)	7/18	Maimburg (2008; 75): 3.7 (1.3–10.5) 9 (1.14–71) ^f Buchmayer (2009; 83): 1.32 (1.01–1.72) ^c
Neonatal encephalopathy	Badawi (2006; 58), Buchmayer (2009; 83), Dodds (2011; 100), Maimburg (2008; 75)		4/4	Buchmayer (2009; 83): 3.06 (1.56–5.99) Dodds (2011; 100): 5.59 (2.32–13.51) Maimburg (2008; 75): 3.1 (1.1–8.7)
Birth defects	Buchmayer (2009; 83), Dawson (2009; 80), Dodds (2011; 100), Guillem (2006; 62), Hultman (2002; 44), Lauritsen (2002; 42), Maimburg (2006; 61), Schendel (2009; 82), Stein (2006; 57), Tripi (2008; 71), Wier (2006; 59), Links (1980; 25)	Bilder (2009; 79), Deykin (1980; 23), Juul-Dam (2001; 40), Mason-Brothers (1990; 30)	12/ 16	Dawson (2009; 80): 1.6 (1.1–2.4) Schendel (2009; 82): 1.7 (1.2–2.4) Wier (2006; 59): 1.7 (1.1–2.4) Hultman (2002; 44): 1.8 (1.1–3.1) Maimburg (2006; 61): 1.9 (1.1–3.5)

Table 3. Continued.

Risk factors	Positive studies (univariate analysis)	Negative or null studies (univariate analysis)	P/T ^a	Adjusted effect estimates ^b
Neonatal or congenital infections	Buchmayer (2009; 83), Atladottir (2010; 89)	Bryson (1988; 27), Dodds (2011; 100), Gillberg (1983; 12), Matsuishi (2005), Piven (1993; 34), Williams (2008; 69)	2/8	Atladottir (2010; 89): 1.5 (1.03–2.32) ^d

Neonatal risk factors for autism that resulted in one positive result (reference) are as follows: fetal hypoxia (91); gastrointestinal diseases (39,47,57); intracranial hemorrhage (30,39,83,100); hypoglycemia (39,75,83); and Apgar 1 < 5 (30,39,69).

Neonatal risk factors for autism that resulted in no positive results are as follows: hemolytic disease (21); elevated IgM (39,97); anemia (12,21,27,100); near-death situation (41); poor condition at birth (23); Apgar 5 < 5 (30,39,69,100) and resuscitation needed (18,27,69); aspiration (21); exchange transfusion (75); difficulties regulating temperature (34); and trauma (83).

^aStatistically significant positive results/total.

^bEffect estimates were Relative Risk and Odds Ratio.

^cAdjusted for mother/pregnancy characteristics only.

^dGirls.

^eMales only.

^fTerm birth.

Table 4. Pre-, peri- and neonatal risk factors in autism: summary of the most robust results.

Family factors	Parental age Parity Mother born abroad
Maternal pregnancy factors	Bleeding Pre-eclampsia
Delivery factors	Breech presentation Scheduled cesarean Small for gestational age
Baby with adverse conditions	Prematurity Low Apgar Hyperbilirubinemia Low birthweight/slow growth Encephalopathy Birth defects

consider preterm birth and LBW as truly being independent from other potential risks, such as being small for gestational age at birth. The phenomenon of being small for gestational age has heterogeneous etiologies, including genetic factors and placental insufficiency (44,101); therefore, the reason for an increased risk of autistic disorders among these children is not clear. In the study by Durkin *et al.*, the mean parental age of autistic children was correlated in unadjusted analyses with birthweight, gestational age and preterm birth (76). Buchmayer *et al.* showed that a possible association between autism and preterm birth was mediated by prenatal factors or neonatal complications (83). Mann *et al.* identified the following three main maternal characteristics that predisposed to LBW

and preterm birth (90): maternal genitourinary tract infection (102); antenatal tobacco use (102); and pre-eclampsia and eclampsia (103). Upon assessing these variables as potential risk factors, neither genitourinary tract infections nor antenatal tobacco was associated with an increased risk for autism. In contrast, the presence of pre-eclampsia or eclampsia appeared to be a strong risk factor, only partly mediated by birthweight. Schendel *et al.* observed that the risk associated with LBW newborns was even higher with other developmental disabilities than it is in autism. This finding led her to hypothesize that coexisting autism as a co-morbidity could be an unrecognized feature of very low birthweight infants and eventual developmental disability (82). Recently, Limperopoulos *et al.* followed a cohort of preterm children with severe preterm delivery and LBW (less than 1500 g) and found a high prevalence of children with a positive initial screening for autism at 20 months (104). In multivariate analyses, a higher score was associated with LBW and gestational age, male gender, prenatal infection, chorioamnionitis, illness severity on admission and abnormal MRI (cerebellar hemorrhagic injury, combined supra- and infratentorial parenchymal damage). These findings were consistent with a previous study in which an isolated cerebellar hemorrhagic injury was found in one-third of preterm children with a positive autism screening test (105). Another recent study noted that a low Apgar score at one minute was specifically associated with an increased risk for autism among LBW babies (106).

All of these recent studies highlight the possibility that the numerous factors associated with both preterm birth and autistic disorders may be interrelated through causal pathways. For a better appraisal of the magnitude of risk of

preterm birth and LBW for autism, future studies should include children with co-morbid congenital anomalies. Moreover, one likely outcome is that among certain subgroups of children, some of these risk factors may be masked when the study design includes both term and preterm children. As Buchmayer *et al.* found different contributing factors for preterm vs. term births, future studies should be designed to include stratified analysis of gestational age (83).

Finally, looking for an association between autism and hyperbilirubinemia is worthwhile for several reasons. Hyperbilirubinemia is thought to exert toxicity on the basal ganglia and cerebellum, two structural brain regions that have been identified as important in the development of autism. Also, a disparity exists concerning the management of this condition, and changes in disease management have led to less aggressive treatment. Such a trend could produce an increment of neurodevelopment sequelae that may account for the increasing prevalence of autism. Hyperbilirubinemia is a neonatal factor that is more distinctive from the prenatal period than are other neonatal factors. Recent, well-designed studies indicate that Rhesus incompatibility might not be the key to understanding hyperbilirubinemia in autism (66,70,73,107). Causes of hyperbilirubinemia in autism require further research.

Discrepancies across studies can partly be explained by the heterogeneity of study design and methodological limitations. Many studies had recruitment and design problems. Firstly, the population of PDD could either be recruited from a population-based registry or from clinical samples at medical centres, inducing unavoidable selection bias. The age at the time of inclusion in studies also led to some differences. Specifically, we may see a possible overestimation of risk factors in younger populations because parents could seek care more rapidly after pre-, peri- and neonatal complications compared with those for whom diagnosis came later (69). On the contrary, it is more difficult to gather valid obstetric information and obtain a reliable diagnosis for older children. The second problem concerns the case ascertainment that led to a great variability between the populations studied. The definition of autism varied according to diagnostic classification. A few studies described a standardized assessment with PDD diagnostic tools, somatic examination for children and new assessments. Most studies based the reliability of diagnosis on retrospective data supplied by parents or medical records. Moreover, depending on authors, the clinical group either constituted children with a broad diagnosis of PDD (52) or a diagnosis restricted to infantile autism (75). Finally, variability of the exclusion criteria (such as co-morbid genetic and neurodevelopmental disorders) may lead to a loss of information about the genesis of autism in some children, especially in those for whom a malformation could be a marker of prenatal insults. Moreover, some evidence points

towards a nonspecific impact of perinatal factors on autism and intellectual disability.

The data collection process also was accompanied by various problems. In some studies, data were derived from multiple pieces of clinical information. Other studies were based only on the parents' interview, which may introduce information bias (28,45). The format of the data itself was often problematic (definitions, a lack of cut-off scores, categorical vs. continuous variables). The lack of a standardized measure of exposure impeded comparison among studies. Several studies used an overall optimality score to implicate the role of obstetric complications in autism. Such a score incorporated an aggregated score of various perinatal and obstetric conditions. This approach may have resulted in nondifferential misclassification, preventing a normative definition of the different types of complications, and underestimated true associations with individual factors (12,35,36,108). Moreover, although an item-by-item analysis could be performed, these weighted scales do not cover certain categories of perinatal factors.

The definition of controls was another factor that contributed to variability among studies. The incidence of obstetric complication could be compared with healthy controls, non-autistic disabled patients, siblings or national statistics (63). Using unaffected siblings as controls may help to identify risk factors and to control for hereditary background, family environment and maternal predisposition to complications in pregnancy or birth (21,23,27,30,32,34,35,36,48,108). These studies excluded sporadic cases of autism for which obstetric factors could share the same etiology as autism itself and, generally, such studies were even more difficult to carry out with adequate sample sizes. A large proportion of studies did not use non-affected controls matched for IQ or corrected for sex ratio (21). However, using a group of non-autistic disabled patients may reduce the magnitude of nonspecific factors potentially involved in several neurodevelopmental disorders.

The final limitation in the comparison of these publications lay in the fact that distributions of children with PDD by factors such as birthweight, primiparous pregnancy and parental age have probably changed over the past 20 years. Therefore, statistically significant changes in risk factor prevalence may change the distribution among autistic children and increase the heterogeneity of results in recent studies and older ones. In the future, this concern should encourage the design of new, large, prospective studies rather than a homogenization of older data.

Conclusion

Despite recent advances in autism genetics, this present review indicates that several risk factors, such as those relevant to the pre-, peri- and neonatal period, may confer a small risk

for autism. Nonetheless, distinguishing whether these risks should be regarded as strictly environmental or related to a genetic vulnerability is not possible. Determining the contribution of these risk factors may improve detection, earlier treatment and better prevention of the disease.

Funding

The study was supported by Grant Agreement No. 2006127 from the European Commission, ENSACP.

References

- Autism and Developmental Disabilities Monitoring Network Surveillance Year 2006 Principal Investigators; Centers for Disease Control and Prevention (CDC). Prevalence of autism spectrum disorders – Autism and Developmental Disabilities Monitoring Network, United States, 2006. *MMWR Surveill Summ.* 2009;58: 1–20.
- Pasamanick B, Rogers ME, Lilienfeld AM. Pregnancy experience and the development of behavior disorders in children. *Am J Psychiatry.* 1956;112:613–18.
- Bailey A, Le Couteur A, Gottesman I, Bolton P, Simonoff E, Yuzda E, et al. Autism as a strongly genetic disorder: evidence from a British twin study. *Psychol Med.* 1995;25:63–77.
- Barbarese WJ, Katusic SK, Colligan RC, Weaver AL, Jacobsen SJ. The incidence of autism in Olmsted County, Minnesota, 1976–1997: results from a population-based study. *Arch Pediatr Adolesc Med.* 2005;159:37–44.
- Parner ET, Schendel DE, Thorsen P. Autism prevalence trends over time in Denmark: changes in prevalence and age at diagnosis. *Arch Pediatr Adolesc Med.* 2008;162:1150–6.
- Minshew NJ, Williams DL. The new neurobiology of autism: cortex, connectivity, and neuronal organization. *Arch Neurol.* 2007;64:945–50.
- Pardo CA, Eberhart CG. The neurobiology of autism. *Brain Pathol.* 2007;17:434–47.
- Zimmerman AW, Connors SL, Matteson KJ, Lee LC, Singer HS, Castaneda JA, et al. Maternal antibrain antibodies in autism. *Brain Behav Immun.* 2007;21:351–7.
- Cohen D, Pichard N, Tordjman S, Baumann C, Burglen L, Excoffier E, et al. Specific genetic disorders and autism: clinical contribution towards their identification. *J Autism Dev Disord.* 2005;35:103–16.
- Guilmatre A, Dubourg C, Mosca AL, Legallie S, Goldenberg A, Drouin-Garraud V, et al. Recurrent rearrangements in synaptic and neurodevelopmental genes and shared biologic pathways in schizophrenia, autism, and mental retardation. *Arch Gen Psychiatry.* 2009;66:947–56.
- Brasic JR, Holland JA. A qualitative and quantitative review of obstetric complications and autistic disorder. *J Dev Phys Disabil.* 2007;19:337–64.
- Gillberg C, Gillberg IC. Infantile autism: a total population study of reduced optimality in the pre-, peri-, and neonatal period. *J Autism Dev Disord.* 1983;13:153–66.
- Cryan E, Byrne M, O'Donovan A, O'Callaghan E. A case-control study of obstetric complications and later autistic disorder. *J Autism Dev Disord.* 1996;26: 453–60.
- Kolevzon A, Gross R, Reichenberg A. Prenatal and perinatal risk factors for autism: a review and integration of findings. *Arch Pediatr Adolesc Med.* 2007;161:326–33.
- Gardener H, Spiegelman D, Buka SL. Prenatal risk factors for autism: comprehensive meta-analysis. *Br J Psychiatry.* 2009;195:7–14.
- American Psychiatric Association. Diagnostic and statistical manual of mental disorders. DSM-IV (4th edn). Washington, DC: American Psychiatric Association, 1994.
- World Health Organization. The ICD-10 classification of mental and behavioural disorders. Diagnostic criteria for research. Geneva: World Health Organization, 1993.
- Lobascher ME, Klingerlee PE, Gubbay SS. Childhood autism: an investigation of aetiological factors in twenty-five cases. *Br J Psychiatry.* 1970;117:525–9.
- Knobloch H, Pasamanick B. Some etiologic and prognostic factors in early infantile autism and psychosis. *Pediatrics.* 1975;55:182–91.
- Torrey EF, Hersch SP, McCabe KD. Early childhood psychosis and bleeding during pregnancy. A prospective study of gravid women and their offspring. *J Autism Child Schizophr.* 1975;5:287–97.
- Finegan JA, Quarrington B. Pre-, peri-, and neonatal factors and infantile autism. *J Child Psychol Psychiatry.* 1979;20:119–28.
- Deykin EY, MacMahon B. Viral exposure and autism. *Am J Epidemiol.* 1979;109:628–38.
- Deykin EY, MacMahon B. Pregnancy, delivery, and neonatal complications among autistic children. *Am J Dis Child.* 1980;134:860–4.
- Gillberg C. Maternal age and infantile autism. *J Autism Dev Disord.* 1980;10:293–7.
- Links PS, Stockwell M, Abichandani F, Simeon J. Minor physical anomalies in childhood autism. Part I. Their relationship to pre- and perinatal complications. *J Autism Dev Disord.* 1980;10:273–85.
- Tsai LY, Stewart MA. Etiological implication of maternal age and birth order in infantile autism. *J Autism Dev Disord.* 1983;13:57–65.
- Bryson SE, Smith IM, Eastwood D. Obstetrical suboptimality in autistic children. *J Am Acad Child Adolesc Psychiatry.* 1988;27:418–22.
- Laxer G, Rey M, Ritvo ER. A comparison of potentially pathologic factors in European children with autism, Down's syndrome, and multiple physical handicaps. *J Autism Dev Disord.* 1988;18:309–13.

29. Levy S, Zoltak B, Saelens T. A comparison of obstetrical records of autistic and nonautistic referrals for psychoeducational evaluations. *J Autism Dev Disord*. 1988;18:573–81.
30. Mason-Brothers A, Ritvo ER, Pingree C, Petersen PB, Jensen WR, McMahon WM, et al. The UCLA-University of Utah epidemiologic survey of autism: prenatal, perinatal, and postnatal factors. *Pediatrics*. 1990;86:514–19.
31. Ward AJ. A comparison and analysis of the presence of family problems during pregnancy of mothers of “autistic” children and mothers of normal children. *Child Psychiatry Hum Dev*. 1990;20:279–88.
32. Lord C, Mulloy C, Wendelboe M, Schopler E. Pre- and perinatal factors in high-functioning females and males with autism. *J Autism Dev Disord*. 1991;21:197–209.
33. Mouridsen SE, Rich B, Isager T. Brief report: parental age in infantile autism, autistic-like conditions, and borderline childhood psychosis. *J Autism Dev Disord*. 1993;23:387–96.
34. Piven J, Simon J, Chase GA, Wzorek M, Landa R, Gayle J, et al. The aetiology of autism: pre-, peri- and neonatal factors. *J Am Acad Child Adolesc Psychiatry*. 1993;32:1256–63.
35. Bolton PF, Murphy M, Macdonald H, Whitlock B, Pickles A, Rutter M. Obstetric complications in autism: consequences or causes of the condition? *J Am Acad Child Adolesc Psychiatry*. 1997;36:272–81.
36. Deb S, Prasad KB, Seth H, Eagles JM. A comparison of obstetric and neonatal complications between children with autistic disorder and their siblings. *J Intellect Disabil Res*. 1997;41:81–6.
37. Zambrino CA, Balottin U, Bettaglio E, Gerardo A, Lanzi G. Obstetrical suboptimality in autistic children: an Italian sample. *J Autism Dev Disord*. 1997;27:493–4.
38. Burd L, Severud R, Kerbeshian J, Klug MG. Prenatal and perinatal risk factors for autism. *J Perinat Med*. 1999;27:441–50.
39. Matsushita T, Yamashita Y, Ohtani Y, Ornitz E, Kuriya N, Murakami Y, et al. Brief report: incidence of and risk factors for autistic disorder in neonatal intensive care unit survivors. *J Autism Dev Disord*. 1999;29:161–6.
40. Juul-Dam N, Townsend J, Courchesne E. Prenatal, perinatal, and neonatal factors in autism, pervasive developmental disorder-not otherwise specified, and the general population. *Pediatrics*. 2001;107:E63.
41. Eaton WW, Mortensen PB, Thomsen PH, Frydenberg M. Obstetric complications and risk for severe psychopathology in childhood. *J Autism Dev Disord*. 2001;31:279–85.
42. Lauritsen MB, Mors O, Mortensen PB, Ewald H. Medical disorders among inpatients with autism in Denmark according to ICD-8: a nationwide register-based study. *J Autism Dev Disord*. 2002;32:115–19.
43. Croen LA, Grether JK, Selvin S. Descriptive epidemiology of autism in a California population: who is at risk? *J Autism Dev Disord*. 2002;32:217–24.
44. Hultman CM, Sparén P, Cnattingius S. Perinatal risk factors for infantile autism. *Epidemiology*. 2002;13:417–23.
45. Wilkerson DS, Volpe AG, Dean RS, Titus JB. Perinatal complications as predictors of infantile autism. *Int J Neurosci*. 2002;112:1085–98.
46. Courchesne E, Carper R, Akshoomoff N. Evidence of brain overgrowth in the first year of life in autism. *JAMA*. 2003;290:337–44.
47. Williams G, Oliver JM, Allard A, Sears L. Autism and associated medical and familial factors: a case control study. *Journal of Developmental and Physical Disabilities*. 2003;15:335–49.
48. Glasson EJ, Bower C, Petterson B, de Klerk N, Chaney G, Hallmayer JF. Perinatal factors and the development of autism: a population study. *Arch Gen Psychiatry*. 2004;61:618–27.
49. Torrey E, Dhavale D, Lawlor J, Yolken R. Autism and head circumference in the first year of life. *Biol Psychiatry*. 2004;56:892–4.
50. Larsson HJ, Eaton WW, Madsen KM, Vestergaard M, Olesen AV, Agerbo E, et al. Risk factors for autism: perinatal factors, parental psychiatric history, and socioeconomic status. *Am J Epidemiol*. 2005;161:916–25; discussion 26–8.
51. Lauritsen MB, Pedersen CB, Mortensen PB. Effects of familial risk factors and place of birth on the risk of autism: a nationwide register-based study. *J Child Psychol Psychiatry*. 2005;46:963–71.
52. Croen LA, Yoshida CK, Odouli R, Newman TB. Neonatal hyperbilirubinemia and risk of autism spectrum disorders. *Pediatrics*. 2005;115:e135–8.
53. Croen LA, Grether JK, Yoshida CK, Odouli R, Van de Water J. Maternal autoimmune diseases, asthma and allergies, and childhood autism spectrum disorders: a case-control study. *Arch Pediatr Adolesc Med*. 2005;159:151–7.
54. Beversdorf DQ, Manning SE, Hillier A, Anderson SL, Nordgren RE, Walters SE, et al. Timing of prenatal stressors and autism. *J Autism Dev Disord*. 2005;35:471–8.
55. Hazlett HC, Poe M, Gerig G, Smith RG, Provenzale J, Ross A, et al. Magnetic resonance imaging and head circumference study of brain size in autism: birth through age 2 years. *Arch Gen Psychiatry*. 2005;62:1366–76.
56. Sugie Y, Sugie H, Fukuda T, Ito M. Neonatal factors in infants with Autistic Disorder and typically developing infants. *Autism*. 2005;9:487–94.
57. Stein D, Weizman A, Ring A, Barak Y. Obstetric complications in individuals diagnosed with autism and in healthy controls. *Compr Psychiatry*. 2006;47:69–75.
58. Badawi N, Dixon G, Felix JF, Keogh JM, Petterson B, Stanley FJ, et al. Autism following a history of newborn encephalopathy: more than a coincidence? *Dev Med Child Neurol*. 2006;48:85–9.
59. Wier ML, Yoshida CK, Odouli R, Grether JK, Croen LA. Congenital anomalies associated with autism spectrum disorders. *Dev Med Child Neurol*. 2006;48:500–7.

60. Reichenberg A, Gross R, Weiser M, Bresnahan M, Silverman J, Harlap S, et al. Advancing paternal age and autism. *Arch Gen Psychiatry*. 2006;63:1026–32.
61. Maimburg RD, Vaeth M. Perinatal risk factors and infantile autism. *Acta Psychiatr Scand*. 2006;114:257–64.
62. Guillem P, Cans C, Guinchat V, Ratel M, Jouk PS. Trends, perinatal characteristics, and medical conditions in pervasive developmental disorders. *Dev Med Child Neurol*. 2006;48:896–900.
63. Brimacombe M, Ming X, Lamendola M. Prenatal and birth complications in autism. *Matern Child Health J*. 2007;11:73–9.
64. Croen LA, Najjar DV, Fireman B, Grether JK. Maternal and paternal age and risk of autism spectrum disorders. *Arch Pediatr Adolesc Med*. 2007;161:334–40.
65. Maimburg RD, Vaeth M. Do children born after assisted conception have less risk of developing infantile autism? *Hum Reprod*. 2007;22:1841–3.
66. Miles JH, Takahashi TN. Lack of association between Rh status, Rh immune globulin in pregnancy and autism. *Am J Med Genet A*. 2007;143A:1397–407.
67. Brown GE, Jones SD, MacKewn AS, Plank EJ. An exploration of possible pre- and postnatal correlates of autism: a pilot survey. *Psychol Rep*. 2008;102:273–82.
68. Kinney DK, Miller AM, Crowley DJ, Huang E, Gerber E. Autism prevalence following prenatal exposure to hurricanes and tropical storms in Louisiana. *J Autism Dev Disord*. 2008;38:481–8.
69. Williams K, Helmer M, Duncan GW, Peat JK, Mellis CM. Perinatal and maternal risk factors for autism spectrum disorders in New South Wales, Australia. *Child Care Health Dev*. 2008;34:249–56.
70. Geier DA, Mumper E, Gladfelter B, Coleman L, Geier MR. Neurodevelopmental disorders, maternal Rh-negativity, and Rho(D) immune globulins: a multi-center assessment. *Neuro Endocrinol Lett*. 2008;29:272–80.
71. Tripi G, Roux S, Canziani T, Bonnet Brilhault F, Barthélemy C, Canziani F. Minor physical anomalies in children with autism spectrum disorder. *Early Hum Dev*. 2008;84:217–23.
72. Schendel D, Bhasin TK. Birth weight and gestational age characteristics of children with autism, including a comparison with other developmental disabilities. *Pediatrics*. 2008;121:1155–64.
73. Croen LA, Matevia M, Yoshida CK, Grether JK. Maternal Rh D status, anti-D immune globulin exposure during pregnancy, and risk of autism spectrum disorders. *Am J Obstet Gynecol*. 2008;199:234.e1–6.
74. Tsuchiya KJ, Matsumoto K, Miyachi T, Tsujii M, Nakamura K, Takagai S, et al. Paternal age at birth and high-functioning autistic-spectrum disorder in offspring. *Br J Psychiatry*. 2008;193:316–21.
75. Maimburg RD, Vaeth M, Schendel DE, Bech BH, Olsen J, Thorsen P. Neonatal jaundice: a risk factor for infantile autism? *Paediatr Perinat Epidemiol*. 2008;22:562–8.
76. Durkin MS, Maenner MJ, Newschaffer CJ, Lee LC, Cunniff CM, Daniels JL, et al. Advanced parental age and the risk of autism spectrum disorder. *Am J Epidemiol*. 2008;168:1268–76.
77. Hack M, Taylor HG, Schluchter M, Andreias L, Drotar D, Klein N. Behavioral outcomes of extremely low birth weight children at age 8 years. *J Dev Behav Pediatr*. 2009;30:122–30.
78. Li J, Vestergaard M, Obel C, Christensen J, Precht DH, Lu M, et al. A nationwide study on the risk of autism after prenatal stress exposure to maternal bereavement. *Pediatrics*. 2009;123:1102–7.
79. Bilder D, Pinborough-Zimmerman J, Miller J, McMahon W. Prenatal, perinatal, and neonatal factors associated with autism spectrum disorders. *Pediatrics*. 2009;123:1293–300.
80. Dawson S, Glasson EJ, Dixon G, Bower C. Birth defects in children with autism spectrum disorders: a population-based, nested case-control study. *Am J Epidemiol*. 2009;169:1296–303.
81. King MD, Fountain C, Dakhallah D, Bearman PS. Estimated autism risk and older reproductive age. *Am J Public Health*. 2009;99:1673–9.
82. Schendel DE, Autry A, Wines R, Moore C. The co-occurrence of autism and birth defects: prevalence and risk in a population-based cohort. *Dev Med Child Neurol*. 2009;51:779–86.
83. Buchmayer S, Johansson S, Johansson A, Hultman CM, Sparen P, Cnattingius S. Can association between preterm birth and autism be explained by maternal or neonatal morbidity? *Pediatrics*. 2009;124:e817–25.
84. Grether JK, Anderson MC, Croen LA, Smith D, Windham GC. Risk of autism and increasing maternal and paternal age in a large North American population. *Am J Epidemiol*. 2009;170:1118–26.
85. Grether JK, Li SX, Yoshida CK, Croen LA. Antenatal ultrasound and risk of autism spectrum disorders. *J Autism Dev Disord*. 2010;40:238–45.
86. Sasanfar R, Haddad SA, Tolouei A, Ghadami M, Yu D, Santangelo SM. Paternal age increases the risk for autism in an Iranian population sample. *Mol Autism*. 2010;1:2.
87. Shelton JF, Tancredi DJ, Hertz-Picciotto I. Independent and dependent contributions of advanced maternal and paternal ages to autism risk. *Autism Res*. 2010;3:30–9.
88. Johnson S, Hollis C, Kochhar P, Hennessy E, Wolke D, Marlow N. Autism spectrum disorders in extremely preterm children. *J Pediatr*. 2010;156:525–31.
89. Atladottir HO, Thorsen P, Schendel DE, Ostergaard L, Lemcke S, Parner ET. Association of hospitalization for infection in childhood with diagnosis of autism spectrum disorders: a Danish cohort study. *Arch Pediatr Adolesc Med*. 2010;164:470–7.
90. Mann JR, McDermott S, Bao H, Hardin J, Gregg A. Pre-eclampsia, birth weight, and autism spectrum disorders. *J Autism Dev Disord*. 2010;40:548–54.

91. Burstyn I, Sithole F, Zwaigenbaum L. Autism spectrum disorders, maternal characteristics and obstetric complications among singletons born in Alberta, Canada. *Chronic Dis Can.* 2010;30:125–34.
92. Karmel BZ, Gardner JM, Meade LS, Cohen IL, London E, Flory MJ, et al. Early medical and behavioral characteristics of NICU infants later classified with ASD. *Pediatrics.* 2010;126:457–67.
93. Hultman CM, Sandin S, Levine SZ, Lichtenstein P, Reichenberg A. Advancing paternal age and risk of autism: new evidence from a population-based study and a meta-analysis of epidemiological studies. *Mol Psychiatry.* 2011;16:1203–12.
94. Maimburg RD, Bech BH, Væth M, Møller-Madsen B, Olsen J. Neonatal jaundice, autism, and other disorders of psychological development. *Pediatrics.* 2010;126:872–8.
95. Zhang X, Lv CC, Tian J, Miao RJ, Xi W, Hertz-Picciotto I, et al. Prenatal and perinatal risk factors for autism in China. *J Autism Dev Disord.* 2010;40:1311–21.
96. Atladóttir HO, Thorsen P, Østergaard L, Schendel DE, Lemcke S, Abdallah M, et al. Maternal infection requiring hospitalization during pregnancy and autism spectrum disorders. *J Autism Dev Disord.* 2010;40:1423–30.
97. Grether JK, Croen LA, Anderson MC, Nelson KB, Yolken RH. Neonatally measured immunoglobulins and risk of autism. *Autism Res.* 2010;3:323–32.
98. Burstyn I, Wang X, Yasui Y, Sithole F, Zwaigenbaum L. Autism spectrum disorders and fetal hypoxia in a population-based cohort: accounting for missing exposures via Estimation-Maximization algorithm. *BMC Med Res Methodol.* 2011;11:2.
99. Haglund NG, Källén KB. Risk factors for autism and Asperger syndrome: perinatal factors and migration. *Autism.* 2011;15:163–83.
100. Dodds L, Fell DB, Shea S, Armson BA, Allen AC, Bryson S. The role of prenatal, obstetric and neonatal factors in the development of autism. *J Autism Dev Disord.* 2011;41:891–902.
101. Palo P, Erkkola R. Risk factors and deliveries associated with preterm, severely small for gestational age fetuses. *Am J Perinatol.* 1993;10:88–91.
102. Goldenberg RL, Andrews WW, Goepfert AR, Faye-Petersen O, Cliver SP, Carlo WA, et al. The Alabama Preterm Birth Study: umbilical cord blood *Ureaplasma urealyticum* and *Mycoplasma hominis* cultures in very preterm newborn infants. *Am J Obstet Gynecol.* 2008;198:43 e1–5.
103. Sibai B, Dekker G, Kupferminc M. Pre-eclampsia. *Lancet.* 2005;365:785–99.
104. Limperopoulos C, Bassan H, Sullivan NR, Soul JS, Robertson RL Jr, Moore M, et al. Positive screening for autism in ex-preterm infants: prevalence and risk factors. *Pediatrics.* 2008;121:758–65.
105. Limperopoulos C, Bassan H, Gauvreau K, Robertson RL Jr, Sullivan NR, Benson CB, et al. Does cerebellar injury in premature infants contribute to the high prevalence of long-term cognitive, learning, and behavioral disability in survivors? *Pediatrics.* 2007;120:584–93.
106. Indredavik MS, Vik T, Evensen KA, Skranes J, Taraldsen G, Brubakk AM. Perinatal risk and psychiatric outcome in adolescents born preterm with very low birth weight or term small for gestational age. *J Dev Behav Pediatr.* 2010;31:286–94.
107. Zandi PP, Kalaydjian A, Avramopoulos D, Shao H, Fallin MD, Newschaffer CJ. Rh and ABO maternal-fetal incompatibility and risk of autism. *Am J Med Genet B Neuropsychiatr Genet.* 2006;141B:643–7.
108. Zwaigenbaum L, Szatmari P, Jones MB, Bryson SE, MacLean JE, Mahoney WJ, et al. Pregnancy and birth complications in autism and liability to the broader autism phenotype. *J Am Acad Child Adolesc Psychiatry.* 2002;41:572–9.

Supporting information

Additional Supporting Information may be found in the on-line version of this article:

Table S1. Description of the studies included in the reviews according to the following characteristics: country where the study was conducted, first author's name and year of publication, number of subjects (*N*), origins of the obstetrical data, psychiatric diagnosis and classifications used, control population characteristics, intellectual quotient included in the analysis (IQ), neurodevelopmental comorbidities excluded or not, adjusted odd ratio calculated (AOC).

Please note: Wiley-Blackwell are not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.