

Persistent primitive reflexes and childhood neurobehavioral disorders.

 EXPORT  Add To My List   

Database: APA PsycInfo Chapter

[Melillo, Robert \(/search/results?term=Melillo,%20Robert&latSearchType=a\)](#)

Citation

Melillo, R. (2016). Persistent primitive reflexes and childhood neurobehavioral disorders. In G. Leisman & J. Merrick (Eds.), *Neuroplasticity in learning and rehabilitation* (pp. 65–99). Nova Biomedical Books.

Abstract

Persistent primitive reflexes have been noted in a number of neurobehavioral disorders and are thought to be related to delayed or absent developmental milestones in this population of children. This is also associated with the presence of clumsiness, incoordination, awkward posture, gait and other motor disturbances. The degree of motor incoordination seems to be related to cognitive dysfunction as well. ADHD, autism, dyslexia as well as almost all neurodevelopmental disorders have been associated with anatomical and functional effects that correlate with the motor incoordination, motor disturbance, cognitive delays and the presence of persistent primitive reflexes. For some time researchers have debated if the structural anatomic and volumetric differences in disorders such as ADHD and autism represent deviant developmental changes or whether they reflect a maturational delay. We review the literature that clearly demonstrates that these disorders and the structural differences represent cortical maturational delays not deviant development. We also note that persistent primitive reflexes are the earliest markers for this delay and that this delayed maturation will eventually lead to the presence of autism, ADHD, and other neurobehavioral disorders. We also note that these disorders and their recent reported increased incidence is related to a combination of genetic and epigenetic factors mostly driven by environmental and lifestyle changes affecting early motor development, sensory stimulation and activity dependent synaptogenesis and neuroplasticity. Symptom variations between these neurobehavioral disorders may be related to asymmetrical maturational differences resulting from different rates of maturation of the right and left hemisphere. Asymmetric persistent primitive reflexes may also be an early marker related to this maturational imbalance. This abnormal pattern of hemispheric asymmetry may lead to desynchronization, underconnectivity, and ultimately a functional disconnection between regions of the brain and cortex. Lastly we comment of the possibility of exercises that can inhibit and remediate persistent primitive reflexes as one possible target for early treatment of these disorders. (PsycINFO Database Record (c) 2016 APA, all rights reserved)



(<https://fastbooks.com/apa-databases/psycinfo/>)

Contact APA (<https://www.apa.org/pubs/databases/contact>) | Give Feedback (</general/feedback>) | Help (<https://help.psycnet.org>)

© 2024 American Psychological Association. All rights, including for text and data mining, AI training, and similar technologies, are reserved. (<https://www.apa.org/about/contact/copyright>)