

Why Do We Find it so Hard to Calculate the Burden of Neurodevelopmental Disorders?

Mary Ann Megan Cleaton
and Amanda Kirby*

Dyscovery Centre, Innovation House,
University of South Wales, William Price
Business Park, Treforest, Pontypridd, Wales,
UK

*Corresponding author: Amanda Kirby

 amanda.kirby@southwales.ac.uk

MBBS, MRCP, PhD, Professor of
Developmental Disorders, Dyscovery
Centre, Innovation House, University of
South Wales, William Price Business Park,
Treforest, Pontypridd, Wales, UK.

Citation: Cleaton MAM, Kirby A (2018)
Why Do We Find it so Hard to Calculate the
Burden of Neurodevelopmental Disorders? J
Child Dev Disord. Vol.4 No.3:10

Abstract

Context: Neurodevelopmental Disorders (NDDs) comprise a spectrum of prevalent disorders affecting social, communication, activity, attention, motor coordination and literacy and numeracy skills. NDDs are highly heritable, frequently co-occur, predominantly persist throughout life and are commonly associated with adverse psychological and psychosocial outcomes for both the individual and their family. However, these features of NDDs are rarely acknowledged or considered within research (including burden-of-illness studies) or by healthcare and educational professionals.

Review: We summarise up-to-date UK and USA prevalence information and compare the relative co-occurrence rates both among NDDs and between NDDs and other psychiatric and medical conditions. We then catalogue the known psychosocial outcomes for individuals with NDDs, their parents and siblings and their children, including considering the heritability of NDDs. We also discuss population groups where NDDs are currently under-diagnosed and consider cases where NDDs may be mis-diagnosed.

Conclusions & Recommendations: We offer suggestions to advance our knowledge of the burden of NDDs and shape the future research agenda to develop interventions, management plans and policies to address the burden. This is of vital importance to ensure individuals with NDDs and their families receive the best possible care.

Keywords: Neurodevelopmental disorders; Prevalence; Co-occurrence; Psychosocial outcome; Burden-of-illness

Received: July 31, 2018; **Accepted:** August 31, 2018; **Published:** September 07, 2018

Introduction

Neurodevelopmental Disorders (NDDs) comprise a spectrum of disorders that include Autism Spectrum Disorder (ASD), Attention Deficit/Hyperactivity Disorder (ADHD), Developmental Coordination Disorder (DCD, also called Dyspraxia), Developmental Language Disorder (DLD, also called Dysphasia and previously called Specific Language Impairment), Dyscalculia (also called Mathematical or Arithmetic Disability), Dyslexia (also called Reading Disability), Intellectual Disability (ID, also called cognitive disability, previously called mental retardation) and Tic Disorders (including Tourette Syndrome and chronic tic disorder). Some definitions of NDDs also include Foetal Alcohol Syndrome, although we have not included it.

The prevalence of NDDs varies globally; however, the burden associated with NDDs is considerable, yet arguably underestimated. This paper provides a comprehensive review of the literature on NDDs, including the scale of the effect, and posits that both missing diagnoses and mis-diagnosis as well as the general 'messiness' of NDDs considerably limits our understanding of this spectrum of disorders. We offer three recommendations to advance our knowledge of the burden of NDDs and shape the future research agenda to develop interventions, management plans and policies to address this burden.

Prevalence

The exact prevalence rates of NDDs vary considerably between

countries and are generally higher in the United States of America (USA) than in the United Kingdom (UK) (**Table 1**). Although it was historically considered that children 'grew out of' most NDDs excepting ASD and ID, there is now good evidence that the impact of NDDs is lifelong for many individuals. In ASD, it has controversially been claimed that 'optimal outcome' (i.e. no or sub-clinical ASD symptoms) may occur in 3-20% of individuals, dependent on symptom severity, age of diagnosis and age of treatment [1]. In any case, most if not all individuals will have persisting challenges to a greater or lesser degree throughout life. The majority of studies indicate ADHD also predominantly persists into late adolescence and adulthood, with persistence occurring in between 50% and 80% of cases [2-7]. However, a few studies indicate much lower persistence of childhood ADHD and an independent cohort of young adult-onset ADHD [8,9]. Approximately 70% of children with DCD continue to exhibit significant motor difficulties through adolescence and

into adulthood [10] and around 73% of 5-year-old children with language impairments continue to have these impairments when aged 18-20 years [11]. Dyslexia and Dyscalculia show similarly high rates of persistence. Jacobson found that among Dyslexic children aged 8-9 years, 83% failed to achieve reading standards expected for their age and experience at age 15-16 years despite the majority of these pupils receiving remedial instruction [12]. Additionally, 85% of these children had persisting phonological deficits and 60 % had persisting decoding deficits at age 18-19 years [13]. Mussolin et al. found that adults aged 18-50 years who had been diagnosed with Dyscalculia as children were slower and more error-prone when completing a mathematics battery [14]. The persistence of Tic Disorders depends on the diagnosis: Provisional Tic Disorder (formerly Transient Tic Disorder) represents the bulk of childhood cases and is classified as a tic disorder that resolves within a year [15]; however, Chronic Tic Disorder (by definition) and Tourette Syndrome are more

Table 1 Estimated childhood prevalence of NDDs.

NDD	UK		USA	
	Est. childhood prevalence (%)	References	Est. childhood prevalence (%)	References
ASD ^a	0.6-3.5	[203-208]	0.5-2.5	[209-216]
ADHD	0.5-2.2	[207,217-220]	2.0-16.1	[56,175,221-246]
DCD	1.8	[247]	<10.1 ^b	[248]
DLD	possibly >0.5, <2.2 ^c	[249,250]	7.4	[251]
Dyscalculia	3.6	[252]	<5.7-6.4 ^d	[253,254]
Dyslexia	2.3-6.2	[252,255,256]	8.3-12.0	[254,257,258]
ID ^e	0.3-0.5 (severe) 2.2-2.5 (mild) 2.7-10.6 (MLD) 0.03 (SLD)	[255,259-261]	1.14-3.6 (all) 0.3-0.4 (severe) 3.7-7.9 (mild)	[259,262-267]
Tic Disorders	6.0 (all) 0.7-3.0 (TS) 1.1 (CTD)	[268-270]	2.7-18.5 (all) 0.3-3.8 (TS)	[239,271-277]

Abbreviations: ADHD, attention-deficit/hyperactivity disorder; ASD, autism spectrum disorder; CTD, chronic tic disorder; DCD, developmental coordination disorder; DLD, developmental language disorder; ID, intellectual disability; MLD, moderate learning difficulty; SLD, severe learning difficulty; TS, Tourette's Syndrome; UK, United Kingdom; USA, United States of America.

- As the prevalence of ASD has increased markedly in the last thirty years, only studies whose data was collected within the last 15 years are included in this estimate.
- No estimate of the prevalence of DCD using DSM diagnostic criteria has been made for the USA. However, 10.1 % of American children scored <5th centile on the Movement Assessment Battery for Children, a test often used to determine Criterion A of a DSM-IV or -V diagnosis of DCD [248]. In contrast, an estimated 4.2-4.5 of UK children score this [27,247].
- The prevalence of DLD among UK 3-year-olds is estimated to be 0.5-0.7% but excluded children of normal intelligence [250]. Lindsay and Strand found that 2.2% of all students aged 5-16 in English schools had Speech, Language or Communication Needs at School Action Plus or above, indicating special educational needs requiring external services [249]. This figure includes children with speech, language and/or communication difficulties other than DLD but excludes children who only require within-school support, children whose parents refuse to have them assessed for special educational needs, children who are electively home-educated, etc. Thus, it is hard to tell if this is an over- or underestimate.
- No estimate of the prevalence of Dyscalculia has been made for the USA. However, Badian (1983) found 6.4% of American children in Grades 1-8 scored <20th centile on the Stanford Achievement Test (SAT) Mathematics Computation Subtest and later [254] found the prevalence of this score was 5.7%. No control was made for IQ, so this figure includes children with general learning disorders as well as those with a specific mathematics learning disorder (i.e. Dyscalculia). Thus, it is an overestimate.
- Prevalence rate estimates of ID depend greatly on diagnostic criteria. In this table, we list the prevalence of severe (IQ < 50), mild (IQ 50-70) and all ID (IQ ≤ 70). Prevalence rates for all ID are frequently estimated to be lower than would be expected by combining prevalence estimates for severe ID and mild ID as mild ID is frequently under-counted in overall estimates. Within the UK, there are few ID prevalence rate estimates using IQ cut-offs. However, there are prevalence rates for moderate and severe learning difficulties (MLD and SLD) – somewhat vague descriptions used to assess special educational need.

persistent. For example, in between 50% and 90% of individuals with Tourette Syndrome at least some tic symptoms persist into adulthood [16-18].

Co-occurrence

The impact of NDDs on the individual is not only lifelong but also dependent on the degree of co-occurrence. Relatively few individuals exhibit a single 'pure' disorder and many exhibits more than two co-occurring disorders (**Table 2**). Even among those with a 'pure' NDD, subclinical co-occurrence is common, for example autism symptoms in children with ADHD [19] or with DLD [20,21]. Indeed, it has been argued that the term 'comorbidity' is of questionable value in the context of NDDs [22]. This is certainly the case for Tourette Syndrome, where 88 % of individuals have at least one co-occurring condition [23], and ASD, where 70% of individuals have at least one co-occurring NDD or psychiatric disorder [24]. The specific pattern and degree of co-occurrence among NDDs has clinical relevance: for example, co-occurrence has a relationship with severity of symptoms and

is, unsurprisingly, associated with poorer outcomes as there are complex cumulative and interactional impacts between NDDs [19,25-27].

In Sweden, the reality of overlapping symptoms and multiple disorders has been recognised by the terms MBD (Minimal Brain Dysfunction, covering a spectrum of deficits, including attention, learning and/or motor deficits, despite normal intelligence) [28], DAMP (Deficits in Attention, Motor Control and Perception, broadly correlating to ADHD plus DCD) [29] and, more recently, ESSENCE (Early Symptomatic Syndromes Eliciting Neurodevelopmental Clinical Examinations, describing symptoms observed in very early childhood that may resolve into ≥ 1 NDDs) [30]. Recognition of the NDD profile has led to clinical pathways and research teams working in a multi-disciplinary manner. Examples of this include a multi-professional, multi-agency referral pathway model in Swansea, UK [31] and the Province of Ontario Neurodevelopmental Disorders (POND) Network in Canada, which has brought together a multi-disciplinary team to research NDDs as a whole [32]. In the United States, similar

Table 2 Estimated co-occurrence of NDDs.

		Proportion with secondary diagnosis of... (%)							
		ASD	ADHD	DCD	DLD	Dyscalculia	Dyslexia	ID ^a	Tic Disorders
Primary diagnosis	ASD		3-78 ^b	25-85 ^c	21 ^d	NE	14 ^e	15-51 ^f	8-60 ^g
	ADHD	6 ^h		18-53 ⁱ	24 ^j	7-18 ^k	18-45 ^l	11-24 ^m	9-33 ⁿ
	DCD	6 ^o	19-53 ^p		$\geq 45q$	31-51 ^r	24-56 ^s	NA ^t	$\leq 34u$
	DLD	4-8 ^v	18-61 ^w	30-71 ^x		62 ^y	48-87 ^z	$\leq 27aa$	NE
	Dyscalculia	NE	39 ^{ab}	25-44 ^{ac}	NE		26-48 ^{ad}	NE ^{ae}	NE
	Dyslexia	NE	18-50 ^{af}	16-53 ^{ag}	NE	39-48 ^{ah}		NE ^{ae}	8 ^{ai}
	ID ^a	10-28 ^{aj}	18-55 ^{ak}	NA ^t	54-79 ^{al}	5 ^{am}	14-17 ^{an}		NE
	Tic Disorders	3-22 ^{ao}	26-82 ^{ap}	13-24 ^{aq}	18 ^{ar}	22-23 ^{as}	22-36 ^{at}	3 ^{au}	

NA, not applicable; NE, no estimates available.

- a. Note that most of the available co-occurrence data regarding ID and other NDDs is from individuals with mild ID (IQ 50-70).
- b. [24,180,278-287].
- c. [288-292].
- d. [293].
- e. [35].
- f. [210-212,214,267,294-297].
- g. [35,298,299].
- h. [300].
- i. [22,181,289,290,301-306].
- j. [307].
- k. [308-310].
- l. [22,302,308,309,311-313].
- m. [302,314-316].
- n. [302,306,313].
- o. [317].
- p. [301,318].
- q. [319].
- r. [320,321].
- s. [27,320].
- t. Children with an IQ<70 are specifically excluded from being given a diagnosis of DCD [15,151].
- u. [26].
- v. [20,322].
- w. [280,323-325].

- x. [185,290,323,326-332].
- y. [322].
- z. [322,333-335].
- aa. [324].
- ab. [336].
- ac. [290,337].
- ad. [336,338-340].
- ae. There are no co-occurrence rate estimates for individuals with a primary diagnosis of Dyslexia or Dyscalculia and a secondary diagnosis of ID, likely because when these conditions co-occur the ID is usually diagnosed first and considered the primary diagnosis.
- af. [22,311-313].
- ag. [22,290,341].
- ah. [338-340].
- ai. [313].
- aj. [267,342,343].
- ak. [342-344].
- al. [342,344].
- am. [342].
- an. [342,343].
- ao. [23,345-350].
- ap. [23,307,313,345,348-354].
- aq. [350,353].
- ar. [307].
- as. [23,349,355].
- at. [23,313,349,350,352,355].
- au. [23,349].

arguments about co-occurrence have been made regarding psychiatric disorders, resulting in the Research Domain Criteria (RDoC) project [33]. Although this project is creating a research classification system and not considering diagnostic criteria or pathways, it marks a paradigm shift by considering psychiatric disorders in terms of affected neurobiological and behavioural systems rather than traditional categories or diagnostic criteria.

Accumulation of Adversity

There is increasing evidence that many individuals with NDDs accumulate problems over time in a cascading and cumulative manner. Cairney et al. proposed the 'Environmental Stress Hypothesis' to explain the developmental cascade from childhood DCD to childhood internalising problems to adult clinical depression and anxiety [34]. Other NDDs also show very high rates of co-occurring and/or secondary health problems, particularly psychiatric conditions (**Table 3**). For example, Hofvander et al. found that 80% of adults with Asperger's disorder and 100% of adults with autism or PDD-NOS had at least one co-occurring DSM-IV Axis I diagnosis [35]. Kirby et al. found that 20% of employed and 33% of unemployed adults with DCD had clinical depression and 79% of employed and 73% of unemployed adults with DCD had borderline or abnormal levels of anxiety [36]. The long-term impact is not always psychological. DCD, for example, has been shown to have an impact on physical health: children who scored $\leq 5^{\text{th}}$ percentile on the Bruininks-Oseretsky

Test of Motor Proficiency, and thus had possible DCD, were at greater risk of being overweight or obese [37] and had lower cardiorespiratory fitness [38] than their typically developing peers.

As well as being associated with an accumulation of medical problems over time, NDDs have also been shown to be associated with adverse psychosocial outcomes. ASD is associated with: lower quality of life [39]; poor social and romantic functioning [21,40]; school bullying [41]; physical and sexual abuse [42,43]; inappropriate sexual behaviours [44]; risky driving behaviour [45]; unemployment [21,46-48]; and, among the employed, low earnings [47]. ADHD has been found to be the most powerful predictor of violent offending among imprisoned male offenders [49]. It has also been associated with: lower quality of life [50,51]; poor academic achievement [52,53]; not being in education, employment or training (NEET) [54]; unemployment [55,56]; problem gambling [57,58]; risky sexual behaviour [59,60]; teenage pregnancy [61]; marriage breakdown and divorce [56,62-64]; multiple marriages [64]; violent and non-violent offending [65]; accident and injury [66,67]; suicide [68]; and all-cause mortality [69]. DLD has also been associated with similar difficulties: lower self-esteem [70-72], shyness [72], poor educational achievement [73-76], early parenthood [76], low socio-economic status (SES) of occupation [75,76] and being NEET [74]. DCD has been associated with lower quality of life [77], poorer self-concept [78], social isolation [79], poor academic achievement [80],

Table 3 Other mental and physical health conditions commonly co-occurring with NDDs.

NDD	Other commonly co-occurring condition(s)	
	Mental	Physical
ASD	Anxiety disorders, eating disorders, gender dysphoria, mood disorders, obsessive-compulsive disorder, personality disorders, schizophrenia, substance use disorders [35,281,283,286,356-360]	Allergies, ear infections, epilepsy or seizures, gastrointestinal disorders, hearing impairment, immune disorders, generalised joint hypermobility, metabolic disorders, neurotransmitter disorders, overweight and obesity, sleep disorders [267,361-372]
ADHD	Anxiety disorders, gender dysphoria, mood disorders, obsessive-compulsive disorder, personality disorders, schizophrenia, substance use disorders [302,314,360,373-376]	Allergies, asthma, epilepsy or seizures, gastrointestinal disorders, headaches or migraine, hearing impairment, generalised joint hypermobility, overweight and obesity, vision impairment, sleep disorders [300,362,365,373,377,378]
DCD	Anxiety disorders, mood disorders, personality disorders, substance use disorders [26,36,81,379]	Epilepsy or seizures, generalised joint hypermobility, overweight and obesity, sleep disorders [82,195,362,364,380-383]
DLD	Anxiety disorders, mood disorders, obsessive-compulsive disorder, personality disorders, schizophrenia [384-386]	Epilepsy or seizures [387]
Dyscalculia	Mood disorders, schizophrenia [388-390]	Epilepsy or seizures [196,391]
Dyslexia	Anxiety disorders, mood disorders, schizophrenia [388-390,392,393]	Epilepsy or seizures [196,198]
ID	Anxiety disorders, eating disorders, mood disorders, personality disorders, schizophrenia, substance use disorders [394-402]	Various, depending on aetiology of ID
Tic Disorders	Anxiety disorders, mood disorders, obsessive-compulsive disorder, personality disorders [23,271,307,348-350,403-405]	Allergies, epilepsy or seizures, sleep disorders [23,349,406,407]

lower physical activity levels [81-83], difficulties with learning to drive [84,85], falls [86,87] and potentially accidents, particularly those associated with crossing roads [88]. Little research has been carried out into the psychosocial outcomes of Dyscalculia. However, Dyslexia has been associated with a reduced quality of life [89] and lower self-esteem [90,91] as well as with poor self-concept when co-occurring with attention or speech problems [92]. ID has been associated with: lower quality of life [93]; stigmatisation [94]; school bullying [95]; physical and sexual abuse and neglect [96,97]; risky sexual behaviour [98]; violent and sexual victimisation and offending [99]; and all-cause mortality [100]. Tic Disorders have been associated with poor psychosocial outcome: lower quality of life [101]; stigmatisation [102]; poorer self-concept [103]; social difficulties [104]; school bullying [105,106]; reduced competence at school [107]; poor academic achievement [108,109]; suicide [110]; and premature mortality [111].

Familial Impact

The effects of an NDD 'ripple out', reaching beyond the affected individual. Most research into these 'ripple effects' has been focused on the families and, particularly, the parents of individuals with NDDs. Parents of children with NDDs have high levels of parenting stress [112]. In particular, ASD is associated with lower family and maternal quality of life [113], increased parental depression and anxiety [114] and affiliate stigma [115]. ADHD is associated with similar issues: lower parental quality of life [116], poorer family functioning [117] and higher levels of parental depressive disorders and depressive personality disorders [118]. Even Dyslexia, which is generally considered a comparatively mild NDD, can have 'ripple effects'. Mothers of children with Dyslexia have higher levels of stress and both anxious and depressive symptoms [112,119,120] although family functioning is not impaired by the presence of Dyslexia and being a parent of a child with Dyslexia is not a significant risk factor for a mental health disorder [121]. ID is generally associated with a good family quality of life [122,123] but is associated with affiliate stigma [115] and higher rates of maternal psychiatric disorder [124,125] and maternal poor health [126]. Tic Disorders are associated with lower parental health-related quality of life (particularly for mothers), increased maternal depression and increased paternal anxiety [127].

Although some of the associations between child NDD and parental psychiatric difficulties may be attributable to shared genetics, at least some are likely to be causal. This is suggested by various studies showing dose-responses, for example between child DCD symptom severity and parenting stress [128], or changes in parental symptoms following child treatment, for example improvement in parental quality of life and depressive mood following methylphenidate treatment of children with ADHD [129]. Although the majority of research has focused on the parents of children with NDDs, it is possible that siblings and other relatives are also affected. Evidence from ASD suggests that effects on siblings are positive as well as negative, may

be complex and are dependent on and/or modulated by other risk factors such as gender, birth order, family size and parental mental health [130].

The impact of NDDs is likely to extend not only to parents and siblings but also to the affected individual's children. In part, this is because NDDs are highly heritable (**Table 4**). The little research regarding the children of individuals with NDDs has predominantly focused on ADHD. Independent of parental psychopathology or offspring ADHD, parental ADHD is associated with high levels of family conflict and low levels of family cohesion [131] whilst maternal ADHD is associated with less positive parenting and consistent discipline by mothers, poorer maternal problem solving and less maternal monitoring of children's activities [132]. Over-reactive and inconsistent discipline has also been reported for mothers and fathers with ADHD both in clinical [133-135] and community samples [136,137]. Overall, the literature suggests that although ADHD is highly heritable, its severity and persistence and the prevalence of co-occurring conditions may be moderated by the quality of parenting and the ability of parents to obtain and consistently deliver treatment to their offspring, which themselves may be affected by parental ADHD and co-occurring conditions [138-143]. It is likely that the effects of other NDDs also extend to the individuals' children, given the known associations between NDDs and psychiatric illness (**Table 3**) and between parental psychiatric illness and offspring mental health [144].

Issues with Estimating the Burden of Disease

Having considered the scale of the effect NDDs have on affected individuals, their families, their communities and society as a whole, one can begin to calculate the burden of illness that NDDs represent. However, the literature is affected by several major issues which produce biases whose effects are difficult to predict. Thus, anyone attempting to compare the burden of NDDs, either between disorders, between countries and/or over time quickly ends up comparing not only apples to oranges but to a whole fruit bowl of inconsistency and inaccuracy.

Table 4 Heritability of neurodevelopmental disorders, as determined using twin studies^a.

NDD	Heritability estimate (%)	References
ASD	64-91	[408]
ADHD	70-80	[409]
DCD	approx. 70	[410]
DLD	>50	[411]
Dyscalculia	38-43	[412,413]
Dyslexia	40-75	[413-417]
ID	Various (dependent on aetiology)	NA
Tic disorders	25-37	[418]

NA, not applicable.

a. Although imperfect, twin studies provide a useful estimate of heritability [419].

Inconsistency in terminology

One of the primary issues which dog the field of NDD research, and thus impede the calculation of the burden of NDDs, is inconsistency of language. This is perhaps best illustrated by DCD, which has been known by a huge variety of other monikers – developmental apraxia, ‘clumsy child syndrome’, motor learning difficulty, perceptuo-motor dysfunction, developmental dyspraxia, sensory integration disorder and DCD [145]. The terms MBD [28], DAMP [29] and ESSENCE [30] also include motor difficulties. Similar issues with terminology have been documented for DLD [146]. As well as inconsistency of language, there are inconsistencies in definition. Again, using DCD as an example – do you exclude individuals diagnosed with ASD [147] or include them [15]? Should you use the Motor Assessment Battery for Children-2 [148]; the Bruininks-Oseretsky Test of Motor Proficiency, Second Edition [149], which measures slightly different constructs; the DCD Questionnaire [150]; or another screening or assessment method? And for these tests do you use a cut-off of $\leq 5^{\text{th}}$ percentile or $\leq 15^{\text{th}}$ percentile [151]? Many studies have used different tools and different cut-offs. Such tools and definitions continue to evolve as we gain greater biological insight into NDDs. However, until standardised definitions, exclusions, diagnostic tools and cut-off points are chosen, validated and consistently used, it will remain exceedingly difficult to compare research findings, particularly across countries and over time.

Transferability

Another major issue affecting our ability to calculate the burden of NDDs is geography. This is a limitation of all health economics studies: they are specific to a given country and its characteristics (income/development level, demographics, structure and funding of healthcare, etc.). However, geography has a particularly strong effect on NDD burden-of-illness studies. Partly, this is due to the marked differences noted in diagnosed prevalence between countries (Table 1), which may themselves be driven by geographic differences in service provision and benefits of a diagnosis (see below). However, NDDs are expensive diseases to manage which have an economic impact on a wide variety of public sectors including education, supported housing and employment, criminal justice and loss of tax revenue due to underemployment. Thus, direct medical costs for NDDs may be lower in countries with a healthcare system based primarily on private healthcare and thus less incentive to pay for early medical interventions to offset long-term costs to the public purse. The USA provides an example of this: its private healthcare system has a history of denying timely, effective treatment to individuals with ASD, resulting in the introduction of state-mandated waiver programmes to enforce private insurance coverage [152]. Subsequently, measurable differences in service use and direct medical costs have been observed between mandate-eligible and ineligible children with ASD [153]. Similar differences in healthcare expenditure are likely to exist between countries with primarily public healthcare, such as the UK, and primarily private healthcare, such as the USA.

Missed diagnoses and mis-diagnoses

To ensure calculations of NDD burden are accurate, robust national estimates of prevalence are required. This requires not only consistent naming of a disorder and use of diagnostic criteria, but also that all individuals requiring support are diagnosed in an accurate and timely manner. However, NDDs are frequently under-diagnosed and this differentially affects various demographic groups. There is increasing recognition that ASD may be under-diagnosed in girls [154,155], in part because the female social landscape may camouflage their ASD [156]. However, ASD diagnoses are also missed or delayed in ethnic minorities [157,158], including Gypsy/Romani and Irish Traveller children [159]; individuals living below the poverty line [157]; and those with co-occurring ADHD [157] and/or with psychiatric conditions [160]. Conversely, co-occurring NDDs are under-diagnosed among individuals with ASD [161]. Under-diagnosis of ADHD occurs in girls [162], children looked after by the state or adopted [163] and the prison population [164] whereas DLD is under-diagnosed in D/deaf individuals [165].

Mis-diagnosis is also an issue. ASD may be incorrectly diagnosed as DLD [166] or as a psychiatric condition, particularly among adolescents and adults [167]. Traumatic brain injury may be mis-diagnosed as ADHD [168]. There is concern that individuals may occasionally feign NDDs, notably Dyslexia [169] and ADHD [170], where there is financial or other gain to be made. It is also suspected that doctors sometimes exaggerate or invent symptoms (so-called ‘upcoding’) in order to gain support for their patients, for example if an individual has significant impairment and meets some criteria of several NDDs but does not pass the diagnostic thresholds for any individual NDD [171]. The introduction of ASD-specific state-mandated healthcare insurance waivers in the USA and other analogous initiatives elsewhere is likely to only exacerbate this by ensuring that those with ASD, but not necessarily those with other, similarly or more severely disabling conditions, receive support [172]. However, notably when considering burden-of-illness studies, under-diagnosis and mis-diagnosis mean that both the prevalence estimates and service use inventories for NDDs may be inaccurate.

Health professionals’ NDD literacy

A final notable issue affecting prevalence estimates of NDDs and thus burden-of-illness calculations is the low level of awareness among healthcare and education professionals and the general public. Without sufficient awareness, many diagnoses will be missed and thus prevalence estimates will be inaccurate. The rising global prevalence of ASD has been attributed in part to rising awareness [173,174]. A similar pattern has been observed for ADHD [175,176]. However, this does not necessarily extend to other NDDs. For example, only 29 % of Canadian, UK and USA family/general physicians and 23 % of teachers are familiar with DCD (Wilson, Neil, Kamps, & Babcock, 2013) whereas even within academic research Dyscalculia plays the poor relation of Dyslexia, which is far more widely recognised [178]. Awareness of NDDs both among professionals and the public is essential to ensure that people with NDDs get the help they need at the time when they need it.

A particular issue is a lack of awareness that NDDs often co-occur. This is true even within academic circles. Certain combinations of NDD are sometimes considered in research, such as ADHD and ASD [179,180] or ADHD and DCD [181,182]. However, studies in general and economic studies in particular rarely consider the co-occurrence of NDDs as a whole and nearly never consider cases where three or more NDDs are present. There is also a disconnect between the literature regarding social, attention and motor difficulties (ASD, ADHD, DCD) and the literature regarding learning difficulties (Dyscalculia, Dyslexia, ID) with very few studies investigating combinations of disorders spanning this divide. ASD research suffers especially in this respect as it has been the habit of researchers to consider associated symptoms such as motor, attention or learning deficits as just 'part of the syndrome' [161] or even a diagnostic feature [183,184] rather than as co-occurring condition(s), an issue potentially driven by the historic exclusion of ADHD and/or DCD diagnoses in the case of ASD using DSM-IV criteria [147]. Within NDD research as a whole, there is often an obsession with identifying cohorts of 'pure' NDDs or restricting cohorts in other respects such as excluding those with low IQ or co-occurring psychiatric conditions. This occurs despite the fact that all evidence points to 'pure' disorders being in the minority (**Table 2**). For example, a study of Canadian children with ADHD, DCD, Dyslexia, Conduct Disorder, Oppositional Defiant Disorder, anxiety and/or depression found that 52% of these children had two or more disorders and among those with ADHD this figure was 80% [22]. The artificial siloing of NDD research helps no one and means results are not representative of most individuals with NDDs.

Recognising the 'messiness' of NDDs

Ignoring the messy reality of NDDs means cohorts are often exceedingly unrepresentative of the population, rendering research at best limited and at worst flawed and potentially meaningless. Additionally, by failing to recognise the prevalence of co-occurrence ourselves, we make it even less likely that healthcare professionals, education professionals, governments and funding bodies will also recognise this. This has knock-on effects, both for patients and for economic analyses. If family/general practitioners and teachers are unaware of the rates of co-occurrence, they are more likely to consider the patient with multiple NDDs as a hypochondriac or 'difficult'. However, the reality is that NDDs are associated with a dose-response of cumulative adversity: individuals with multiple co-occurring NDDs are in most need of support. Moreover, some combinations of NDDs and/or psychiatric conditions are associated with worse outcomes than others. For example: presence of a co-occurring psychiatric condition is associated with persistence of ADHD and greater functional impairment [2,6]; co-occurrence of DCD and DLD is associated with lower quality of life than DLD alone [185]; and presence of multiple NDDs is associated with increasingly impaired language and academic skills in a dose-responsive manner [186]. By ignoring the issue of co-occurrence in NDDs and ignoring the fact that not all co-occurring combinations of NDDs are equal, we risk leaving the most vulnerable inadequately supported and trapped in an endless cycle of cumulative adversity. This represents not only an ethical problem but also a financial

one as individuals accumulate psychiatric conditions, and/or poorer outcomes such as entering the criminal justice system or becoming or remaining unemployed or homeless [26]. A better understanding of the prevalence, service usage and outcome of various combinations of co-occurring NDDs is sorely needed in order to accurately account for the effects of co-occurrence on the cost of illness.

What Must We do to Capture the True Burden of NDDs?

We have catalogued the major issues preventing us from capturing and comparing the true burden of NDDs. Rectifying these represents a significant challenge. However, three key recommendations can be made.

Firstly, the issue of co-occurrence needs to be confronted in order that services be targeted appropriately and the cost of these be accurately assessed. One model for this is ESSENCE [30], which proposes that children aged less than three years presenting with any issues in general development, communication and language, social inter-relatedness, motor coordination, attention, activity, behaviour, mood and/or sleep not be compartmentalised off to 'ASD only', 'ADHD only', etc. centres. Instead, they should initially be assessed in a holistic manner for all ESSENCE symptoms in order to consider the overall impairment of the child and thus provide all necessary services in a timely manner. Subsequently, specialist/targeted (e.g. ASD-specific, ADHD-specific) support can be provided on the basis of need. A similar model is advocated by the POND Network [32]. Given the prevalence of co-occurrence among NDDs (**Table 2**) and the demonstrated importance of early treatment for example for ASD [187] or ADHD [188], these seem eminently sensible suggestions. Indeed, we would take it further and suggest that the ESSENCE/POND model should be followed irrespective of age at initial diagnosis and that, particularly among individuals whose diagnosis was delayed until adulthood, assessment for psychiatric conditions should also be automatically considered.

Secondly, we need to address the issue of under-diagnosis of NDDs in certain sociodemographic groups. This is necessary to generate accurate prevalence estimates on which to base burden-of-illness calculations but also in order to improve equity in service delivery. We do not endorse the wholesale screening of all children. However, there is an argument for the targeted use of routine screening following the ESSENCE/POND model within at-risk populations. These populations include those known to be more likely to have NDDs such as looked-after and adopted children [189], children excluded from school [190] and prison populations [49,191]. It may also be beneficial to target populations with certain medical conditions based on their known co-occurrence with NDDs (**Table 4**). An example is Rolandic epilepsy (also known as Benign Childhood Epilepsy with Centrotemporal Spikes (BECCTS)), which frequently co-occurs with ADHD [192,193], DLD [194], DCD [195], Dyscalculia [196,197] and Dyslexia [198]. In addition, given the heritable nature of NDDs, screening should be considered where there is a strong family history of NDDs.

However, routine screening and increased professional awareness is only part of the process. We as a society need to ensure that diagnosis is a worthwhile process that results in benefits that outweigh the downsides such as stigma [199] or the fear that a 'label' may become a self-fulfilling prophecy [200]. Unless this is done, individuals and/or their families may opt out of gaining a diagnosis [200] as they fear over-diagnosis, i.e. correct diagnosis that nonetheless does not benefit the individual as it comes with stigma and/or little or no medical or educational assistance [201]. This is a valid fear. For example, among American adolescents and young adults with Asperger's Syndrome, symptom severity is *inversely* associated with risk of victimization [202].

Thirdly, we need to base burden-of-illness studies on all aspects of NDDs not just the direct medical costs paid by public or private healthcare insurance. This is a particular issue with NDDs as many of the costs are borne by other public services such as education, supported housing, training and employment. Other costs of NDDs are also hidden, for example under the costs of co-occurring, potentially secondary, psychiatric conditions (**Table 3**) or the costs of unemployment, homelessness or being in the criminal justice system. There is a clear need for society to invest in early diagnosis and timely provision of appropriate medical and educational services for individuals with NDDs in order to minimise the economic and psychosocial burden of these hidden

costs. Until robust burden-of-illness calculations are carried out for NDDs that consider all aspects of economic impact, we cannot conduct comprehensive economic evaluations that accurately reflect the population. Without these, we will be unable to ensure that individuals with NDDs receive the best care possible in a timely manner.

Acknowledgement

MAMC and AK thank Prof Paula Lorgelly and two anonymous reviewers for proof-reading this review.

Conflicts of Interest

MAMC declares she has no potential conflicts of interest to disclose. AK is the parent of an adult with NDDs, the chair of Movement Matters UK, patron of the Dyspraxia Association of New Zealand, Advisor to the Dyspraxia Association in the Republic of Ireland, Medical Advisor to the Dyspraxia Foundation in the UK and is on the Hidden Impairment National Group for the Department for Work and Pensions, UK.

Compliance with Ethical Standards

MAMC was supported by a grant from the Waterloo Foundation (grant no. 1944/3218) whilst writing this paper.

References

- Helt M, Kelley E, Kinsbourne M, Pandey J, Boorstein H, et al. (2008) Can children with autism recover? If so, how?. *Neuropsychol Rev* 18: 339-366.
- Biederman J, Petty CR, Clarke A, Lomedico A, Faraone SV (2011) Predictors of persistent ADHD: an 11-year follow-up study. *J Psychiatr Res* 45: 150-155.
- Cadman T, Findon J, Eklund H, Hayward H, Howley D, et al. (2016) Six-year follow-up study of combined type ADHD from childhood to young adulthood: predictors of functional impairment and comorbid symptoms. *Eur Psychiatry* 35: 47-54.
- Guelzow BT, Loya F, Hinshaw SP (2017) How Persistent is ADHD into Adulthood? Informant Report and Diagnostic Thresholds in a Female Sample. *J Abnorm Child Psychol* 45: 301-312.
- Hechtman L, Swanson JM, Sibley MH, Stehli A, Owens EB, et al. (2016) Functional adult outcomes 16 years after childhood diagnosis of attention-deficit/hyperactivity disorder: MTA results. *J Am Acad Child Adolesc Psychiatry* 55: 945-952.
- Mick E, Byrne D, Fried R, Monuteaux M, Faraone SV, et al. (2011) Predictors of ADHD persistence in girls at 5-year follow-up. *J Atten Disord* 15: 183-192.
- Tuithof M, Ten Have M, Van Dorsselaer S, De Graaf R (2014) Prevalentie, persistentie en gevolgen van ADHD in de Nederlandse volwassen bevolking. *Tijdschr Psychiatr* 56: 10-19.
- Agnew-Blais JC, Polanczyk GV, Danese A, Wertz J, Moffitt TE, et al. (2016) Evaluation of the persistence, remission, and emergence of attention-deficit/hyperactivity disorder in young adulthood. *JAMA Psychiatry* 73: 713-720.
- Moffitt TE, Houts R, Asherson P, Belsky DW, Corcoran DL, et al. (2015) Is adult ADHD a childhood-onset neurodevelopmental disorder? evidence from a four-decade longitudinal cohort study. *Am J Psychiatry* 172: 967-977.
- Losse A, Henderson SE, Elliman D, Hall D, Knight E, et al. (1991) Clumsiness in children-do they grow out of it? A 10-year follow-up study. *Dev Med Child Neurol* 33: 55-68.
- Johnson CJ, Beitchman JH, Young A, Escobar M, Atkinson L, et al. (1999) Fourteen-year follow-up of children with and without speech/language impairments. *J Speech Lang Hear Res* 42: 744.
- Jacobson C (1999) How persistent is reading disability? Individual growth curves in reading. *Dyslexia* 5: 78-93.
- Svensson I, Jacobson C (2006) How persistent are phonological difficulties? A longitudinal study of reading retarded children. *Dyslexia* 12: 3-20.
- Mussolin C, Martin R, Schiltz C (2011) Relationships between number and space processing in adults with and without dyscalculia. *Acta Psychol (Amst)* 138: 193-203.
- American Psychiatric Association (2013) Diagnostic and statistical manual of mental disorders (5th Edn). Washington, DC: American Psychiatric Publishing.
- Leckman JF, Zhang H, Vitale A, Lahnin F, Lynch K, et al. (1998) Course of tic severity in tourette syndrome: the first two decades. *Pediatrics* 102: 14-19.
- Pappert EJ, Goetz CG, Louis ED, Leurgans S (2003) Objective assessments of longitudinal outcome in Gilles de la Tourette's syndrome. *Neurology* 61: 936-940.
- Burd L, Kerbeshian J, Klug MG, Burd L, Avery K, et al. (2001) Long-

- term follow-up of an epidemiologically defined cohort of patients with Tourette syndrome. *J Child Neurol* 16: 431-437.
- 19 Green JL, Rinehart N, Anderson V, Nicholson JM, Jongeling B, et al. (2015) Autism spectrum disorder symptoms in children with ADHD: a community-based study. *Res Dev Disabil* 47: 175-184.
- 20 Conti-Ramsden G, Simkin Z, Botting N (2006) The prevalence of autistic spectrum disorders in adolescents with a history of specific language impairment (SLI). *J Child Psychol Psychiatry Allied Discip* 47: 621-628.
- 21 Howlin P, Mawhood L, Rutter M (2000) Autism and Developmental Receptive Language Disorder-a follow-up comparison in early adult life. II: social, behavioural, and psychiatric Outcomes. *J Child Psychol Psychiatry* 41 : 561-578.
- 22 Kaplan BJ, Dewey DM, Crawford SG, Wilson BN (2001) The term comorbidity is of questionable value in reference to developmental disorders: data and theory. *J Learn Disabil* 34: 555-565.
- 23 Freeman RD, Fast DK, Burd L, Kerbeshian J, Robertson MM, et al. (2000) An international perspective on Tourette syndrome: selected findings from 3500 individuals in 22 countries. *Dev Med Child Neurol* 42: 436-447.
- 24 Simonoff E, Pickles A, Charman T, Chandler S, Loucas T, et al. (2008) Psychiatric disorders in children with autism spectrum disorders: prevalence, comorbidity, and associated factors in a population-derived sample. *J Am Acad Child Adolesc Psychiatry* 47: 921-929.
- 25 Gadke DL, McKinney C, Oliveros A (2016) Autism Spectrum Disorder Symptoms and Comorbidity in Emerging Adults. *Child Psychiatry Hum Dev* 47: 194-201.
- 26 Rasmussen P, Gillberg C (2000) Natural outcome of ADHD with developmental coordination disorder at age 22 years: a controlled, longitudinal, community-based study. *J Am Acad Child Adolesc Psychiatry* 39: 1424-1431.
- 27 Schoemaker MM, Lingam R, Jongmans MJ, van Heuvelen MJG, Emond A (2013) Is severity of motor coordination difficulties related to co-morbidity in children at risk for developmental coordination disorder?. *Res Dev Disabil* 34 : 3084-3091.
- 28 Köhler L (1973) Physical examination of four-year-old children. *Acta Paediatr Scand* 62: 181-192.
- 29 Gillberg C (1986) Attention deficit disorder: diagnosis, prevalence, management and outcome. *Paediatrician* 3: 108-118.
- 30 Gillberg C (2010) The ESSENCE in child psychiatry: early symptomatic syndromes eliciting neurodevelopmental clinical examinations. *Res Dev Disabil* 31: 1543-1551.
- 31 Salmon G, Cleave H, Samuel C (2006) Development of multi-agency referral pathways for attention-deficit hyperactivity disorder, developmental coordination disorder and autistic spectrum disorders: reflections on the process and suggestions for new ways of working. *Clin Child Psychol Psychiatry* 11: 63-81.
- 32 <https://pond-network.ca/>
- 33 Cuthbert BN, Insel TR (2013) Toward the future of psychiatric diagnosis: the seven pillars of RDoC. *BMC Med* 11: 126.
- 34 Cairney J, Rigoli D, Piek J (2013) Developmental coordination disorder and internalizing problems in children: the environmental stress hypothesis elaborated. *Dev Rev* 33: 224-238.
- 35 Hofvander B, Delorme R, Chaste P, Nydén A, Wentz E, et al. (2009) Psychiatric and psychosocial problems in adults with normal-intelligence autism spectrum disorders. *BMC Psychiatry* 9: 1-9.
- 36 Kirby A, Williams N, Thomas M, Hill EL (2013) Self-reported mood, general health, wellbeing and employment status in adults with suspected DCD. *Res Dev Disabil* 34: 1357-1364.
- 37 Cairney J, Hay J, Veldhuizen S, Missiuna C, Mahlberg N, et al. (2010) Trajectories of relative weight and waist circumference among children with and without developmental coordination disorder. *Cmaj* 182: 1167-1172.
- 38 Cairney J, Veldhuizen S, King-Dowling S, Faught BE, Hay J (2017) Tracking cardiorespiratory fitness and physical activity in children with and without motor coordination problems. *J Sci Med Sport* 20: 380-385.
- 39 Ikeda E, Hinckson E, Krägeloh C (2014) Assessment of quality of life in children and youth with autism spectrum disorder: a critical review. *Qual Life Res* 23: 1069-1085.
- 40 Stokes M, Newton N, Kaur A (2007) Stalking, and social and romantic functioning among adolescents and adults with autism spectrum disorder. *J Autism Dev Disord* 37: 1969-1986.
- 41 Hwang S, Kim YS, Koh YJ, Leventhal BL (2018) Autism spectrum disorder and school bullying: who is the victim? who is the perpetrator?. *J Autism Dev Disord* 48: 225-238.
- 42 Brown-Lavoie SM, Viecili MA, Weiss JA (2014) Sexual knowledge and victimization in adults with autism spectrum disorders. *J Autism Dev Disord* 44: 2185-2196.
- 43 Mandell DS, Walrath CM, Manteuffel B, Sgro G, Pinto-Martin J (2005) The prevalence and correlates of abuse among children with autism served in comprehensive community-based mental health settings. *Child Abuse Negl* 29: 1359-1372.
- 44 Fernandes LC, Gillberg CI, Cederlund M, Hagberg B, Gillberg C, et al. (2016) Aspects of sexuality in adolescents and adults diagnosed with autism spectrum disorders in childhood. *J Autism Dev Disord* 46: 3155-3165.
- 45 Daly BP, Nicholls EG, Patrick KE, Brinckman DD, Schultheis MT (2014) Driving behaviors in adults with autism spectrum disorders. *J Autism Dev Disord* 44: 3119-3128.
- 46 Bush KL, Tassé MJ (2017) Employment and choice-making for adults with intellectual disability, autism, and down syndrome. *Res Dev Disabil* 65: 23-34.
- 47 Roux AM, Shattuck PT, Cooper BP, Anderson KA, Wagner M, et al. (2013) Postsecondary employment experiences among young adults with an autism spectrum disorder. *J Am Acad Child Adolesc Psychiatry* 52: 931-939.
- 48 Shattuck PT, Narendorf SC, Cooper BP, Sterzing PR, Wagner M, et al. (2012) Postsecondary education and employment among youth with an autism spectrum disorder. *Pediatrics* 129: 1042-1049.
- 49 Young S, Thome J (2011) ADHD and offenders. *World J Biol Psychiatry* 12: 124-128.
- 50 Agarwal R, Goldenberg M, Perry R, Ishak WW (2012) The quality of life of adults with attention deficit hyperactivity disorder: a systematic review. *Innov Clin Neurosci* 9: 10-21.
- 51 Danckaerts M, Sonuga-Barke EJS, Banaschewski T, Buitelaar JK, Döpfner M, et al. (2010) The quality of life of children with attention deficit/hyperactivity disorder: a systematic review. *Eur Child Adolesc Psychiatry* 19: 83-105.
- 52 Erskine HE, Norman RE, Ferrari AJ, Chan GC, Copeland WE, et al.

- (2016) Long-term outcomes of attention-deficit/hyperactivity disorder and conduct disorder: a systematic review and meta-analysis. *J Am Acad Child Adolesc Psychiatry* 55: 841-850.
- 53 Fleming M, Fitton CA, Steiner MFC, McLay JS, Clark D, et al. (2017) Educational and health outcomes of children treated for attention-deficit/hyperactivity disorder. *JAMA Pediatr* 171: e170691.
 - 54 Goldman-Mellor S, Caspi A, Arseneault L, Ajala N, Ambler A, et al. (2016) Committed to work but vulnerable: self-perceptions and mental health in NEET 18-year-olds from a contemporary British cohort. *J Child Psychol Psychiatry* 57: 196-203.
 - 55 Halmøy A, Fasmer OB, Gillberg C, Haavik J (2009) Occupational outcome in adult ADHD: impact of symptom profile, comorbid psychiatric problems, and treatment. *J Atten Disord* 13: 175-187.
 - 56 Kessler RC, Adler L, Barkley R, Biederman J, Conners CK, et al. (2006) The prevalence and correlates of adult ADHD in the United States: results from the national comorbidity survey replication. *Am J Psychiatry* 163: 716-723.
 - 57 Retz W, Ringling J, Retz-Junginger P, Vogelgesang M, Rösler M (2016) Association of attention-deficit/hyperactivity disorder with gambling disorder. *J Neural Transm* 123: 1013-1019.
 - 58 Waluk OR, Youssef GJ, Dowling NA (2016) The relationship between problem gambling and attention deficit hyperactivity disorder. *J Gambl. Stud* 32: 591-604.
 - 59 Chen MH, Hsu JW, Huang KL, Bai YM, Ko NY, et al. (2018) Sexually transmitted infection among adolescents and young adults with attention-deficit/hyperactivity disorder: a nationwide longitudinal study. *J Am Acad Child Adolesc Psychiatry* 57: 48-53.
 - 60 Olazagasti MAR, Klein RG, Mannuzza S, Belsky ER, Hutchison JA, et al. (2013) Does childhood attention-deficit/hyperactivity disorder predict risk-taking and medical illnesses in adulthood? *J Am Acad Child Adolesc Psychiatry* 52: 153-162.
 - 61 Lehti V, Niemelö S, Heinze M, Sillanmaki L, Helenius H, et al. (2012) Childhood predictors of becoming a teenage mother among Finnish girls. *Acta Obstet Gynecol Scand* 91: 1319-1325.
 - 62 Biederman J, Faraone SV, Spencer T, Wilens T, Norman D, et al. (1993) Patterns of psychiatric comorbidity, cognition, and psychosocial functioning in adults with attention deficit hyperactivity disorder. *Am J Psychiatry* 150: 1792-1798.
 - 63 Biederman J, Faraone SV, Spencer T, Wilens T, Mick E, et al. (1994) Gender differences in a sample of adults with attention deficit hyperactivity disorder. *Psychiatry Res* 53: 13-29.
 - 64 Murphy K, Barkley RA (1996) Attention deficit hyperactivity disorder adults: comorbidities and adaptive impairments. *Compr Psychiatry* 37: 393-401.
 - 65 Knecht C, De Alvaro R, Martinez-Raga J, Balanza-Martinez V (2015) Attention-deficit hyperactivity disorder (ADHD), substance use disorders, and criminality: a difficult problem with complex solutions. *Int J Adolesc Med Health* 27: 163-175.
 - 66 Shilon Y, Pollak Y, Aran A, Shaked S, Gross-Tsur V (2012) Accidental injuries are more common in children with attention deficit hyperactivity disorder compared with their non-affected siblings. *Child Care Health Dev* 38: 366-370.
 - 67 Vaa T (2014) ADHD and relative risk of accidents in road traffic: a meta-analysis. *Accid Anal Prev* 62: 415-425.
 - 68 Impey M, Heun R (2012) Completed suicide, ideation and attempt in attention deficit hyperactivity disorder. *Acta Psychiatr Scand* 125: 93-102.
 - 69 London AS, Landes SD (2016) Attention deficit hyperactivity disorder and adult mortality. *Prev Med (Baltim)* 90: 8-10.
 - 70 Jerome AC, Fujiki M, Brinton B, James SL (2002) Self-esteem in children with specific language impairment. *J Speech Lang Hear Res* 45: 700.
 - 71 Lindsay G, Dockrell J, Palikara O (2010) Self-esteem of adolescents with specific language impairment as they move from compulsory education. *Int J Lang Commun Disord* 45: 561-571.
 - 72 Wadman R, Durkin K, Conti-Ramsden G (2008) Self-esteem, shyness, and sociability in adolescents with specific language impairment (SLI). *J Speech Lang Hear Res* 51: 938.
 - 73 Conti-Ramsden G, Durkin K, Simkin Z, Knox E (2009) Specific language impairment and school outcomes. I: identifying and explaining variability at the end of compulsory education. *Int J Lang Commun Disord* 44: 15-35.
 - 74 Conti-Ramsden G, Durkin K (2012) Postschool educational and employment experiences of young people with specific language impairment. *Lang Speech Hear Serv Sch* 43: 507.
 - 75 Conti-Ramsden G, Durkin K, Toseeb U, Botting N, Pickles A (2017) Education and employment outcomes of young adults with a history of developmental language disorder. *Int J Lang Commun Disord* 53: 237-255.
 - 76 Johnson CJ, Beitchman JH, Brownlie EB (2010) Twenty-year follow-up of children with and without speech-language impairments: family, educational, occupational, and quality of life outcomes. *Am J Speech-Language Pathol* 19: 51-65.
 - 77 Zwicker JG, Harris SR, Klassen AF (2013) Quality of life domains affected in children with developmental coordination disorder: a systematic review. *Child Care Health Dev* 39: 562-580.
 - 78 Cocks N, Barton B, Donnelly M (2009) Self-concept of boys with developmental coordination disorder. *Phys Occup Ther Pediatr* 29: 6-22.
 - 79 Kirby A, Edwards L, Sugden D (2011) Emerging adulthood in developmental co-ordination disorder: parent and young adult perspectives. *Res Dev Disabil* 32: 1351-1360.
 - 80 Harrowell I, Hollén L, Lingam R, Emond A (2018) The impact of developmental coordination disorder on educational achievement in secondary school. *Res Dev Disabil* 72: 13-22.
 - 81 Hill EL, Brown D (2013) Mood impairments in adults previously diagnosed with developmental coordination disorder. *J Ment Heal* 22: 334-340.
 - 82 Joshi D, Missiuna C, Hanna S, Hay J, Faught BE, et al. (2015) Relationship between BMI, waist circumference, physical activity and probable developmental coordination disorder over time. *Hum Mov Sci* 40: 237-247.
 - 83 Rivlis I, Hay J, Cairney J, Klentrou P, Liu J, et al. (2011) Physical activity and fitness in children with developmental coordination disorder: a systematic review. *Res Dev Disabil* 32: 894-910.
 - 84 Cousins M, Smyth MM (2003) Developmental coordination impairments in adulthood. *Hum Mov Sci* 22: 433-459.
 - 85 Kirby A, Sugden D, Beveridge S, Edwards L, Edwards R (2008) Dyslexia and developmental co-ordination disorder in further and higher education - similarities and differences. Does the 'label' influence the support given? *Dyslexia* 14: 197-213.
 - 86 Scott-Roberts S, Purcell C (2015) Incidence and impact of falls in

- adults with significant motor coordination difficulties. *J Comorbidity* 5: 44.
- 87 Scott-Roberts S, Purcell C (2018) Understanding the functional mobility of adults with developmental coordination disorder (DCD) through the international classification of functioning (icf). *Curr Dev Disord Reports* 5: 26-33.
- 88 Purcell C, Wilmut K, Wann JP (2017) The use of visually guided behaviour in children with Developmental Coordination Disorder (DCD) when crossing a virtual road. *Hum Mov Sci* 53: 37-44.
- 89 Balazs J, Miklosi M, Toro KT, Nagy-Varga D (2016) Reading disability and quality of life based on both self- and parent-reports: importance of gender differences. *Front Psychol* 7: 1-9.
- 90 Burden R (2008) Is dyslexia necessarily associated with negative feelings of self-worth? a review and implications for future research. *Dyslexia* 14: 188-196.
- 91 Carawan LW, Nalavany BA, Jenkins C (2016) Emotional experience with dyslexia and self-esteem: the protective role of perceived family support in late adulthood. *Aging Ment Health* 20: 284-294.
- 92 McArthur G, Castles A, Kohnen S, Banales E (2016) Low self-concept in poor readers: prevalence, heterogeneity, and risk. *PeerJ* 4: e2669.
- 93 Simões C, Santos S (2016) Comparing the quality of life of adults with and without intellectual disability. *J Intellect Disabil Res* 60: 378-388.
- 94 Ali A, King M, Strydom A, Hassiotis A (2016) Self-reported stigma and its association with socio-demographic factors and physical disability in people with intellectual disabilities: results from a cross-sectional study in England. *Soc Psychiatry Psychiatr Epidemiol* 51: 465-474.
- 95 Maïano C, Aimé A, Salvat MC, Morin AIS, Normand CL (2016) Prevalence and correlates of bullying perpetration and victimization among school-aged youth with intellectual disabilities: a systematic review. *Res Dev Disabil* 49: 181-195.
- 96 Byrne G (2017) Prevalence and psychological sequelae of sexual abuse among individuals with an intellectual disability. *J Intellect Disabil* 22: 294-310.
- 97 Maclean MJ, Sims S, Bower C, Leonard H, Stanley FJ, et al. (2017) Maltreatment Risk Among Children With Disabilities. *Pediatrics* 139: e20161817.
- 98 Baines S, Emerson E, Robertson J, Hatton C (2018) Sexual activity and sexual health among young adults with and without mild/moderate intellectual disability. *BMC Public Health* 18: 1-12.
- 99 Fogden BC, Thomas SDM, Daffern M, Ogloff JRP (2016) Crime and victimisation in people with intellectual disability: a case linkage study. *BMC Psychiatry* 16: 1-9.
- 100 Hosking FJ, Carey IM, Shah SM, Harris T, DeWilde S, et al. (2016) Mortality among adults with intellectual disability in England: comparisons with the general population. *Am J Public Health* 106: 1483-1490.
- 101 Evans J, Seri S, Cavanna AE (2016) The effects of Gilles de la Tourette syndrome and other chronic tic disorders on quality of life across the lifespan: a systematic review. *Eur Child Adolesc Psychiatry* 25: 939-948.
- 102 Malli MA, Forrester-Jones R, Murphy G (2016) Stigma in youth with Tourette's syndrome: a systematic review and synthesis. *Eur Child Adolesc Psychiatry* 25: 127-139.
- 103 Hanks CE, McGuire JF, Lewin AB, Storch EA, Murphy TK (2016) Clinical correlates and mediators of self-concept in youth with chronic tic disorders. *Child Psychiatry Hum Dev* 47: 64-74.
- 104 McGuire JF, Hanks C, Lewin AB, Storch EA, Murphy TK (2013) Social deficits in children with chronic tic disorders: phenomenology, clinical correlates and quality of life. *Compr Psychiatry* 54: 1023-1031.
- 105 Storch EA, Murphy TK, Chase RM, Keeley M, Goodman WK, et al. (2007) Peer victimization in youth with Tourette's syndrome and chronic tic disorder: relations with tic severity and internalizing symptoms. *J Psychopathol Behav Assess* 29: 211-219.
- 106 Zinner SH, Conelea CA, Glew GM, Woods DW, Budman CL (2012) Peer victimization in youth with Tourette syndrome and other chronic tic disorders. *Child Psychiatry Hum Dev* 43: 124-136.
- 107 Carter AS, O'Donnell DA, Schultz RT, Scahill L, Leckman JF, et al. (2000) Social and emotional adjustment in children affected with Gilles de la Tourette's syndrome: associations with ADHD and family functioning. *J Child Psychol Psychiatry* 41: 215-223.
- 108 Cubo E, Gonzalez C, Ausin V, Delgado V, Saez S, et al. (2017) The association of poor academic performance with tic disorders: a longitudinal, mainstream school-based population study. *Neuroepidemiology* 48: 155-163.
- 109 Pérez-Vigil A, Fernández de la Cruz L, Brander G, Isomura K, Jangmo A, et al. (2018) Association of Tourette syndrome and chronic tic disorders with objective indicators of educational attainment: a population-based sibling comparison study. *JAMA Neurol* 1194.
- 110 Fernández de la Cruz L, Rydell M, Runeson B, Brander G, Rück C, et al. (2017) Suicide in Tourette's and chronic tic disorders. *Biol Psychiatry* 82: 111-118.
- 111 Meier SM, Dalsgaard S, Mortensen PB, Leckman JF, Plessen KJ (2017) Mortality risk in a nationwide cohort of individuals with tic disorders and with Tourette syndrome. *Mov Disord* 32: 605-609.
- 112 Craig F, Operto FF, De Giacomo A, Margari L, Frolli A, et al. (2016) Parenting stress among parents of children with neurodevelopmental disorders. *Psychiatry Res* 242: 121-129.
- 113 McKechnie A, Moffat V, Johnstone E, Fletcher-Watson S (2017) Links between autism spectrum disorder diagnostic status and family quality of life. *Children* 4: 23.
- 114 Chan KKS, Lam CB, Law NCW, Cheung RYM (2018) From child autistic symptoms to parental affective symptoms: a family process model. *Res Dev Disabil* 75: 22-31.
- 115 Mak WWS, Cheung RYM (2008) Affiliate stigma among caregivers of people with intellectual disability or mental illness. *J Appl Res Intellect Disabil* 21: 532-545.
- 116 Cappe E, Bolduc M, Rougé MC, Saiag MC, Delorme R (2017) Quality of life, psychological characteristics, and adjustment in parents of children with attention-deficit/hyperactivity disorder. *Qual Life Res* 26: 1283-1294.
- 117 Cussen A, Sciberras E, Ukoumunne OC, Efron D (2012) Relationship between symptoms of attention-deficit/hyperactivity disorder and family functioning: a community-based study. *Eur J Pediatr* 171: 271-280.
- 118 Margari F, Craig F, Petruzzelli MG, Lamanna A, Matera E, et al. (2013) Parents psychopathology of children with attention deficit hyperactivity disorder. *Res Dev Disabil* 34: 1036-1043.
- 119 Karande S, Kumbhare N, Kulkarni M, Shah N (2009) Anxiety levels in mothers of children with specific learning disability. *J Postgrad Med* 55: 165-170.

- 120 Snowling MJ, Muter V, Carroll J (2007) Children at family risk of dyslexia: a follow-up in early adolescence. *J Child Psychol Psychiatry Allied Discip* 48: 609-618.
- 121 Bonifacci P, Montuschi M, Lami L, Snowling MJ (2014) Parents of children with dyslexia: cognitive, emotional and behavioural profile. *Dyslexia* 20: 175-190.
- 122 Caples M, Sweeney J (2011) Quality of life: a survey of parents of children/adults with an intellectual disability who are availing of respite care. *Br J Learn Disabil* 39: 64-72.
- 123 McFelea JT, Raver S (2012) Quality of life of families with children who have severe developmental disabilities: a comparison based on child residence. *Phys Disabil Educ Relat Serv* 31: 3-17.
- 124 Fairthorne J, de Klerk N, Leonard H (2016) Brief report: burden of care in mothers of children with autism spectrum disorder or intellectual disability. *J Autism Dev Disord* 46: 1103-1109.
- 125 Fairthorne J, Jacoby P, Bourke J, de Klerk N, Leonard H (2015) Onset of maternal psychiatric disorders after the birth of a child with intellectual disability: a retrospective cohort study. *J Psychiatr Res* 61: 223-230.
- 126 Fairthorne J, De Klerk N, Leonard H (2015) Health of mothers of children with intellectual disability or autism spectrum disorder: a review of the literature. *Med Res Arch* 3: 1-21.
- 127 Jalenques I, Auclair C, Morand D, Legrand G, Marcheix M, et al. (2017) Health-related quality of life, anxiety and depression in parents of adolescents with Gilles de la Tourette syndrome: a controlled study. *Eur Child Adolesc Psychiatry* 26: 603-617.
- 128 Takahashi M, Adachi M, Takayanagi N, Yasuda S, Tanaka M, et al. (2017) Coordination difficulties in preschool-aged children are associated with maternal parenting stress: a community-based cross-sectional study. *Res Dev Disabil* 70: 11-21.
- 129 Kim Y, Kim B, Chang JS, Kim BN, Cho SC, et al. (2014) Parental quality of life and depressive mood following methylphenidate treatment of children with attention-deficit hyperactivity disorder. *Psychiatry Clin Neurosci* 68: 506-514.
- 130 Green L (2013) The well-being of siblings of individuals with autism. *ISRN Neurol* 2013: 1-7.
- 131 Biederman J, Faraone SV, Monuteaux MC (2002) Impact of exposure to parental attention-deficit hyperactivity disorder on clinical features and dysfunction in the offspring. *Psychol Med* 32: 817-827.
- 132 Murray C, Johnston C (2006) Parenting in mothers with and without attention-deficit/hyperactivity disorder. *J Abnorm Psychol* 115: 52-61.
- 133 Arnold EH, O'Leary SG, Edwards GH (1997) Father involvement and self-reported parenting of children with attention deficit-hyperactivity disorder. *J Consult Clin Psychol* 65: 337-342.
- 134 Chronis-Tuscano A, Raggi VL, Clarke TL, Rooney ME, Diaz Y, et al. (2008) Associations between maternal attention-deficit/hyperactivity disorder symptoms and parenting. *J Abnorm Child Psychol* 36: 1237-1250.
- 135 Johnston C, Scoular DJ, Ohan JL (2004) Mothers' reports of parenting in families of children with symptoms of attention-deficit/hyperactivity disorder: relations to impression management. *Child Fam Behav Ther* 26: 45-61.
- 136 Banks T, Ninowski JE, Mash EJ, Semple DL (2008) Parenting behavior and cognitions in a community sample of mothers with and without symptoms of attention-deficit/hyperactivity disorder. *J Child Fam Stud* 17: 28-43.
- 137 Chen M, Johnston C (2007) Maternal inattention and impulsivity and parenting behaviors. *J Clin Child Adol Psychol* 36: 455.
- 138 Chronis AM, Lahey BB, Pelham WE, Williams SH, Baumann BL, et al. (2007) Maternal depression and early positive parenting predict future conduct problems in young children with attention-deficit/hyperactivity disorder. *Dev Psychol* 43: 70-82.
- 139 Harold GT, Leve LD, Barrett D, Elam K, Neiderhiser JM, et al. (2013) Biological and rearing mother influences on child ADHD symptoms: Revisiting the developmental interface between nature and nurture. *J Child Psychol Psychiatry* 54: 1038-1046.
- 140 Molina BSG, Pelham WEJ, Cheong J, Marshal MP, Gnagy EM, et al. (2012) Childhood adhd and growth in adolescent alcohol use: the roles of functional impairments, adhd symptom persistence and parental knowledge. *J Abnorm Psychol* 121: 922-935.
- 141 Tung I, Brammer WA, Li JJ, Lee SS (2015) Parenting behavior mediates the intergenerational association of parent and child offspring ADHD symptoms. *J Clin Child Adolesc Psychol* 44: 787-799.
- 142 Walther CAP, Cheong J, Molina BSG, Pelham WEJ, Wymbs BTet al. (2012) Substance use and delinquency among adolescents with childhood adhd: the protective role of parenting. *Psychol Addict Behav* 26: 585-598.
- 143 Wang CH, Mazursky-Horowitz H, Chronis-Tuscano A (2014) Delivering evidence-based treatments for child attention-deficit/hyperactivity disorder (ADHD) in the context of parental ADHD. *Curr Psychiatry Rep* 16: 474.
- 144 Vostanis P, Graves A, Meltzer H, Goodman R, Jenkins R, et al. (2006) Relationship between parental psychopathology, parenting strategies and child mental health: findings from the GB national study. *Soc Psychiatry Psychiatr Epidemiol* 41: 509-514.
- 145 Gibbs J, Appleton J, Appleton R (2007) Dyspraxia or developmental coordination disorder? unravelling the enigma. *Arch Dis Child* 92: 534-539.
- 146 Bishop DVM (2000) Why is it so hard to reach agreement on terminology? the case of developmental language disorder (DLD). *Int J Lang Commun Disord* 52: 671-680.
- 147 American Psychiatric Association (2000) Diagnostic and statistical manual of mental disorders (4th Edn), Washington, DC.
- 148 Henderson SE, Sugden DA, Barnett AL (2007) Movement Assessment Battery for Children-2 (2nd Edn). London, UK: The Psychological Corporation.
- 149 Bruininks RH, Bruininks BD (2005) BOT2: Bruininks-Oseretsky Test of Motor Proficiency (2nd Edn). Minneapolis, MN, USA: Pearson Assessments.
- 150 Wilson BN, Kaplan BJ, Crawford SG, Campbell A, Dewey D (2000) Reliability and validity of a parent questionnaire on childhood motor skills. *Am J Occup Ther* 54: 484-493.
- 151 <https://www.pearsonclinical.co.uk/Psychology/ChildCognitionNeuropsychologyandLanguage/ChildPerceptionandVisuomotorAbilities/MABC-2/Resources/LeedsConsensus06.pdf>
- 152 Maunch D, Pfefferle S, Booker C, Pustell M, Levin J (2011) Report on state services to individuals with autism spectrum disorders (ASD). Cambridge, MA.

- 153 Barry CL, Epstein AJ, Marcus SC, Kennedy-Hendricks A, Candon MK, et al. (2017) Effects of state insurance mandates on health care use and spending for autism spectrum disorder. *Health Aff* 36: 1754-1761.
- 154 Kirkovski M, Enticott PG, Fitzgerald PB (2013) A review of the role of female gender in autism spectrum disorders. *J Autism Dev Disord* 43: 2584-2603.
- 155 Van Wijngaarden-Cremers PJM, Van Eeten E, Groen WB, Van Deurzen PA, Oosterling IJ, et al. (2014) Gender and age differences in the core triad of impairments in autism spectrum disorders: a systematic review and meta-analysis. *J Autism Dev Disord* 44: 627-635.
- 156 Dean M, Harwood R, Kasari C (2017) The art of camouflage: gender differences in the social behaviors of girls and boys with autism spectrum disorder. *Autism* 21: 678-689.
- 157 Jo H, Schieve LA, Rice CE, Yeargin-Allsopp M, Tian LH, et al. (2015) Age at autism spectrum disorder (ASD) diagnosis by race, ethnicity, and primary household language among children with special health care needs, united states, 2009-2010. *Matern Child Health J* 19: 1687-1697.
- 158 Begeer S, El Bouk S, Boussaid W, Terwogt MM, Koot HM (2009) Underdiagnosis and referral bias of autism in ethnic minorities. *J Autism Dev Disord* 39: 142-148.
- 159 Emerson E (2012) Deprivation, ethnicity and the prevalence of intellectual and developmental disabilities. *J Epidemiol Community Health* 66: 218-224.
- 160 Levy SE, Giarelli E, Lee LC, Schieve LA, Kirby RS, et al. (2010) Autism spectrum disorder and co-occurring developmental, psychiatric, and medical conditions among children in multiple populations of the United States. *J Dev Behav Pediatr* 31: 267-275.
- 161 Gillberg C, Fernell E (2014) Autism plus versus autism pure. *J Autism Dev Disord* 44: 3274-3276.
- 162 Nussbaum NL (2012) ADHD and female specific concerns: a review of the literature and clinical implications. *J Atten Disord* 16: 87-100.
- 163 Woolgar M, Baldock E (2015) Attachment disorders versus more common problems in looked after and adopted children: comparing community and expert assessments. *Child Adolesc Ment Health* 20: 34-40.
- 164 Buitelaar NJL, Ferdinand RF (2016) ADHD undetected in criminal adults. *J Atten Disord* 20: 270-278.
- 165 Mason K, Rowley K, Marshall CR, Atkinson JR, Herman R, et al. (2010) Identifying specific language impairment in deaf children acquiring British sign language: implications for theory and practice. *Br J Dev Psychol* 28: 33-49.
- 166 Bishop DVM, Whitehouse AJO, Watt HJ, Line EA (2008) Autism and diagnostic substitution: evidence from a study of adults with a history of developmental language disorder. *Dev Med Child Neurol* 50: 341-345.
- 167 Takara K, Kondo T, Kuba T (2015) How and why is autism spectrum disorder misdiagnosed in adult patients? - From diagnostic problem to management for adjustment. *Ment Health Fam Med* 11: 73-88.
- 168 McAvinue L, O'Keeffe F, McMackin D, Robertson IH (2005) Impaired sustained attention and error awareness in traumatic brain injury: implications for insight. *Neuropsychol. Rehabil* 15: 569-587.
- 169 Harrison AG, Edwards MJ, Armstrong I, Parker KCH (2010) An investigation of methods to detect feigned reading disabilities. *Arch Clin Neuropsychol* 25: 89-98.
- 170 Tucha L, Fuermaier ABM, Koerts J, Groen Y, Thome J (2015) Detection of feigned attention deficit hyperactivity disorder. *J Neural Transm* 122: 123-134.
- 171 Cartwright J, Lasser J, Gottlieb MC (2017) To code or not to code: some ethical conflicts in diagnosing children. *Pract Innov* 2: 195-206.
- 172 McDowell M, O'Keeffe M (2012) Public services for children with special needs: discrimination by diagnosis? *J Paediatr Child Health* 48: 2-5.
- 173 Elsabbagh M, Divan G, Koh YJ, Kim YS, Kauchali S, et al. (2012) Global prevalence of autism and other pervasive developmental disorders. *Autism Res* 5: 160-179.
- 174 Hansen SN, Schendel DE, Parner ET (2015) Explaining the increase in the prevalence of autism spectrum disorders: the proportion attributable to changes in reporting practices. *JAMA Pediatr* 169: 56-62.
- 175 Akinbami LJ, Liu X, Pastor PN, Reuben CA (2011) Attention deficit hyperactivity disorder among children aged 5-17 years in the United States 1998-2009. *NCHS Data Brief* 70: 1-8.
- 176 Rydell M, Lundström S, Gillberg C, Lichtenstein P, Larsson H (2018) Has the attention deficit hyperactivity disorder phenotype become more common in children between 2004 and 2014? Trends over 10 years from a Swedish general population sample. *J Child Psychol Psychiatry* 59: 863-871.
- 177 Wilson BN, Neil K, Kamps PH, Babcock S (2013) Awareness and knowledge of developmental co-ordination disorder among physicians, teachers and parents. *Child Care Health Dev* 39: 296-300.
- 178 Price GR, Ansari D (2013) Dyscalculia: characteristics, causes, and treatments. *Numeracy* 6: 2.
- 179 Green JL, Rinehart N, Anderson V, Efron D, Nicholson JM, et al. (2016) Association between autism symptoms and family functioning in children with attention-deficit/hyperactivity disorder: a community-based study. *Eur Child Adolesc Psychiatry* 25: 1307-1318.
- 180 Hanson E, Cerban BM, Slater CM, Caccamo LM, Bacic J, et al. (2013) Brief report: prevalence of attention deficit/hyperactivity disorder among individuals with an autism spectrum disorder. *J Autism Dev Disord* 43: 1459-1464.
- 181 Landgren M, Pettersson R, Kjellman B, Gillberg C (1996) ADHD, DAMP and other neurodevelopmental/ psychiatric disorders in 6-year-old children: epidemiology and co-morbidity. *Dev Med Child Neurol* 38: 891-906.
- 182 Tal Saban M, Ornoy A, Parush S (2014) Executive function and attention in young adults with and without developmental coordination disorder - a comparative study. *Res Dev Disabil* 35: 2644-2650.
- 183 Ghaziuddin M, Butler E, Tsai L, Ghaziuddin N (1994) Is clumsiness a marker for asperger syndrome? *J Intellect Disabil Res* 38: 519-527.
- 184 Ghaziuddin M, Butler E (1998) Clumsiness in autism and asperger syndrome: a further report. *J Intellect Disabil Res* 42: 43-48.
- 185 Flapper BCT, Schoemaker MM (2013) Developmental coordination disorder in children with specific language impairment: co-morbidity and impact on quality of life. *Res Dev Disabil* 34: 756-763.
- 186 Crawford SG, Kaplan BJ, Dewey D (2006) Effects of coexisting disorders on cognition and behavior in children with ADHD. *J Atten Disord* 10: 192-199.
- 187 Magán-Maganto M, Bejarano-Martín Á, Fernández-Alvarez C, Narzisi

- A, García-Primo P, et al. (2017) Early detection and intervention of ASD: a European overview. *Brain Sci* 7: E159.
- 188 Chronis-Tuscano A, Wang CH, Strickland J, Almirall D, Stein MA (2016) Personalized treatment of mothers with ADHD and their young at-risk children: A SMART pilot. *J Clin Child Adolesc Psychol* 45: 510-521.
- 189 Ford T, Vostanis P, Meltzer H, Goodman R (2007) Psychiatric disorder among British children looked after by local authorities : comparison with children living in private households. *Br J Psychiatry* 190: 319-325.
- 190 Paget A, Parker C, Heron J, Logan S, Henley W, et al. (2018) Which children and young people are excluded from school? Findings from a large British birth cohort study, the Avon Longitudinal Study of Parents and Children (ALSPAC). *Child Care Health Dev* 44: 285-296.
- 191 Ashworth S (2016) Autism is underdiagnosed in prisoners. *BMJ* 353: 1-2.
- 192 Holtmann M, Becker K, Kentner-Figura B, Schmidt MH (2004) Increased frequency of rolandic spikes in ADHD children. *Epilepsia* 45: 564-566.
- 193 Kim EH, Yum MS, Kim HW, Ko TS (2014) "Attention-deficit/hyperactivity disorder and attention impairment in children with benign childhood epilepsy with centrotemporal spikes. *Epilepsy Behav* 37: 54-58.
- 194 Smith AB, Bajomo O, Pal DK (2015) A meta-analysis of literacy and language in children with rolandic epilepsy. *Dev Med Child Neurol* 57: 1019-1026.
- 195 Kirby A, Williams N, Koelewijn L, Brindley LM, Muthukumaraswamy SD, et al. (2017) Benign childhood epilepsy with centrotemporal spikes (BECTS) and developmental co-ordination disorder. *Epilepsy Behav* 72: 122-126.
- 196 Canavese C, Rigardetto R, Viano V, Vittorini R, Bassi B, et al. (2007) Are dyslexia and dyscalculia associated with Rolandic epilepsy? A short report on ten Italian patients. *Epileptic Disord* 9: 432-436.
- 197 Piccinelli P, Borgatti R, Aldini A, Bindelli D, Ferri M, et al. (2008) Academic performance in children with rolandic epilepsy. *Dev Med Child Neurol* 50: 353-356.
- 198 Clarke T, Strug LJ, Murphy PL, Bali B, Carvalho J, et al. (2007) High risk of reading disability and speech sound disorder in Rolandic epilepsy families: case-control study. *Epilepsia* 48: 2258-2265.
- 199 Kinnear SH, Link BG, Ballan MS, Fischbach RL (2016) Understanding the experience of stigma for parents of children with autism spectrum disorder and the role stigma plays in families' lives. *J Autism Dev Disord* 46: 942-953.
- 200 Russell G, Norwich B (2012) Dilemmas, diagnosis and destigmatization: parental perspectives on the diagnosis of autism spectrum disorders. *Clin Child Psychol Psychiatry* 17: 229-245.
- 201 Coon ER, Quinonez RA, Moyer VA, Schroeder AR (2014) Overdiagnosis: How our compulsion for diagnosis may be harming children. *Pediatrics* 134: 1013-1023.
- 202 Shtayermman O (2009) An exploratory study of the stigma associated with a diagnosis of asperger's syndrome: the mental health impact on the adolescents and young adults diagnosed with a disability with a social nature. *J Hum Behav Soc Environ* 19: 298-313.
- 203 Baird G, Simonoff E, Pickles A, Chandler S, Loucas T, et al. (2006) Prevalence of disorders of the autism spectrum in a population cohort of children in South Thames: the Special Needs and Autism Project (SNAP). *Lancet* 368: 210-215.
- 204 Baron-Cohen S, Scott FJ, Allison C, Williams J, Bolton P, et al. (2009) Prevalence of autism-spectrum conditions: UK school-based population study. *Br J Psychiatry* 194: 500-509.
- 205 Brugha T, McManus S, Bankart J, Scott F, Purdon S, et al. (2011) Epidemiology of autism spectrum disorders in adults in the community in England. *Arch Gen Psychiatry* 68: 459-466.
- 206 Latif AHA, Williams WR (2007) Diagnostic trends in autistic spectrum disorders in the South Wales valleys. *Autism* 11: 479-487.
- 207 Russell G, Rodgers LR, Ukoumunne OC, Ford T (2014) Prevalence of parent-reported ASD and ADHD in the UK: findings from the Millennium Cohort Study. *J Autism Dev Disord* 44: 31-40.
- 208 Dillenburger K, Jordan JA, McKerr L, Keenan M (2015) The millennium child with autism: early childhood trajectories for health, education and economic wellbeing. *Dev Neurorehabil* 18: 37-46.
- 209 CDC (2009) Prevalence of autism spectrum disorders - autism and developmental disorders monitoring network, United States, 2006. *Morb Mortal Wkly Rep* 58: 1-20.
- 210 CDC (2012) Prevalence of autism spectrum disorders - autism and developmental disabilities monitoring network, 14 sites, United States, 2008. *Morb Mortal Wkly Rep* 61: 1-19.
- 211 CDC (2014) Prevalence of autism spectrum disorder among children aged 8 years - autism and developmental disabilities monitoring network, 11 sites, United States, 2010. *Morb Mortal Wkly Rep* 63: 1-21.
- 212 Christensen DL, Baio J, Braun KVN, Bilder D, Charles JM, et al. (2016) Prevalence and characteristics of autism spectrum disorder among children aged 8 years - autism and developmental disabilities monitoring network, 11 Sites, United States, 2012. *Morb Mortal Wkly Rep* 65: 1-23.
- 213 Perou R, Bitsko RH, Blumberg SJ, Pastor P, Ghandour RM, et al. (2013) Mental health surveillance among children--United States, 2005-2011. *Morb Mortal Wkly Rep* 62: 1-35.
- 214 Rice C, Nicholas J, Baio J, Pettygrove S, Lee LC, et al. (2010) Changes in autism spectrum disorder prevalence in 4 areas of the United States. *Disabil Health J* 3: 186-201.
- 215 Windham GC, Anderson MC, Croen LA, Smith KS, Collins J, et al. (2011) Birth prevalence of autism spectrum disorders in the San Francisco Bay area by demographic and ascertainment source characteristics. *J Autism Dev Disord* 41: 1362-1372.
- 216 Zahorodny W, Shenouda J, Howell S, Rosato NS, Peng B, et al. (2014) Increasing autism prevalence in metropolitan New Jersey. *Autism* 18: 117-126.
- 217 Ford T, Goodman R, Meltzer H (2003) The British child and adolescent mental health survey 1999: The prevalence of DSM-IV disorders. *J Am Acad Child Adolesc Psychiatry* 42: 1203-1211.
- 218 Green H, McGinnity Á, Meltzer H, Ford T, Goodman R (2005) Mental health of children and young people in Great Britain 2004: Summary Report.
- 219 Holden SE, Jenkins-Jones S, Poole CD, Morgan CL, Coghill D, et al. (2013) The prevalence and incidence, resource use and financial costs of treating people with attention deficit/hyperactivity disorder (ADHD) in the United Kingdom (1998 to 2010). *Child Adolesc Psychiatry Ment Health* 7: 1-13.
- 220 McCarthy S, Wilton L, Murray ML, Hodgkins P, Asherson P, et al. (2012) The epidemiology of pharmacologically treated attention deficit hyperactivity disorder (ADHD) in children, adolescents and

adults in UK primary care. *BMC Pediatr* 12: 78.

- 221 Gimpel GA, Kuhn BR (2000) Maternal report of attention deficit hyperactivity disorder symptoms in preschool children. *Child Care Heal Dev* 26: 163-176.
- 222 Heiligenstein E, Conyers LM, Ms ARB, Smith A, Heiligenstein E, et al. (1998) Preliminary normative data on DSM-IV attention deficit hyperactivity disorder in college students. *J Am Coll Heal* 48: 185-188.
- 223 Keenan K, Shaw DS, Walsh B, Delliquadri E, Giovannelli J (1997) DSM-III-R disorders in preschool children from low-income families. *J Am Acad Child Adolesc Psychiatry* 36: 620-627.
- 224 Lavigne JV, Gibbons RD, Christoffel KK, Arend R, Rosenbaum D, et al. (1996) Prevalence rates and correlates of psychiatric disorders among preschool children. *J Am Acad Child Adolesc Psychiatry* 35: 204-214.
- 225 Lavigne JV, LeBailly SA, Hopkins J, Gouze KR, Binns HJ (2009) The prevalence of ADHD, ODD, depression, and anxiety in a community sample of 4-year-olds. *J Clin Child Adolesc Psychol* 38: 315-328.
- 226 McConaughy SH, Achenbach TM (1994) Comorbidity of empirically based syndromes in matched general population and clinical samples. *J Child Psychol Psychiatry* 35: 1141-1157.
- 227 Murphy K, Barkley RA (1996) Prevalence of DSM-IV symptoms of ADHD in adult licensed drivers: Implications for clinical diagnosis. *J Atten Disord* 1: 147-161.
- 228 Neuman RJ, Sitdhiraksa N, Reich W, Ji THC, Joyner CA, et al. (2005) Estimation of prevalence of DSM-IV and latent class-defined ADHD subtypes in a population-based sample of child and adolescent twins. *Twin Res Hum Genet* 8: 392-401.
- 229 Nolan EE, Gadow KD, Sprafkin J (2001) Teacher reports of DSM-IV ADHD, ODD, and CD symptoms in schoolchildren. *J Am Acad Child Adolesc Psychiatry* 40: 241-249.
- 230 Ramtekkar U, Reiersen AM, Todorov AA, Todd RD (2010) Sex and age differences in attention-deficit/hyperactivity disorder symptoms and diagnoses: Implications for DSM-IV and ICD-11. *J Am Acad Child Adolesc Psychiatry* 49: e1-e3.
- 231 Carlson CL, Tamm L, Gaub M (1997) Gender differences in children with ADHD, ODD, and co-occurring ADHD/ODD identified in a school population. *J Am Acad Child Adolesc Psychiatry* 36: 1706-1714.
- 232 Rasmussen ER, Neuman RJ, Heath AC, Levy F, Hay DA, et al. (2004) Familial clustering of latent class and DSM-IV defined attention-deficit / hyperactivity disorder (ADHD) subtypes. *J Child Psychol Psychiatry* 45: 589-598.
- 233 Rowland AS, Skipper BJ, Umbach DM, Rabiner DL, Campbell RA, et al. (2015) The prevalence of ADHD in a population-based sample. *J Atten Disord* 19: 741-754.
- 234 Shaffer D, Fisher P, Dulcan MK, Davies M, Piacentini J, et al. (1996) The NIMH diagnostic interview schedule for children version 2.3 (DISC- 2.3): description, acceptability, prevalence rates, and performance in the MECA study. *J Am Acad Child Adolesc Psychiatry* 35: 865-877.
- 235 Todd RD, Rasmussen ER, Neuman RJ, Reich W, Hudziak JJ, et al. (2001) Familiality and heritability of subtypes of attention deficit hyperactivity disorder in a population sample of adolescent female twins. *Am J Psychiatry* 158: 1891-1898.
- 236 Wolraich ML, Hannah JN, Pinnock TY, Baumgaertel A, Brown J (1996) Comparison of diagnostic criteria for attention-deficit hyperactivity disorder in a county-wide sample. *J Am Acad Child Adolesc Psychiatry* 35: 319-324.
- 237 Wolraich ML, Hannah JN, Baumgaertel A, Feurer ID (1998) Examination of DSM-IV criteria for attention deficit/hyperactivity disorder in a county-wide sample. *J Dev Behav Pediatr* 19: 162-168.
- 238 Wolraich ML, McKeown RE, Visser SN, Bard D, Cuffe S, et al. (2014) The prevalence of ADHD: its diagnosis and treatment in four school districts across two states. *J Atten Disord* 18: 563-575.
- 239 Costello JE, Angold A, Burns BJ (1996) The great smoky mountains study of youth goals, design, methods, and the prevalence of DSM-III-R disorders. *JAMA Psychiatry* 53: 1129-1136.
- 240 Costello J, Mustillo ES, Erkanli A, Keeler G, Angold A (2003) Prevalence and development of psychiatric disorders in childhood and adolescence. *Arch Gen Psychiatry* 60: 837-844.
- 241 DuPaul GJ, Schaughency EA, Weyandt LL, Tripp G, Kiesner J, et al. (2001) Self-report of ADHD symptoms in university students. *J Learn Disabil* 34: 370-379.
- 242 Earls F (1982) Application of DSM-III in an epidemiological study of preschool children. *Am J Psychiatry* 139: 242-243.
- 243 Egger HL, Angold A (2006) Common emotional and behavioral disorders in preschool children: Presentation, nosology, and epidemiology. *J Child Psychol Psychiatry Allied Discip* 47: 313-337.
- 244 Faraone SV, Biederman J (2005) What is the prevalence of adult adhd? Results of a population screen of 966 adults. *J Atten Disord* 9: 384-391.
- 245 Froehlich TE, Lanphear BP, Epstein JN, Barbaresi WJ, Katusic SK, et al. (2007) Prevalence, recognition, and treatment of attention-deficit/hyperactivity disorder in a national sample of US children. *Arch Pediatr Adolesc Med* 161: 857-864.
- 246 Gaub M, Carlson CL (1997) Behavioral characteristics of DSM-IV ADHD subtypes in a school-based population. *J Abnorm Child Psychol* 25: 103-111.
- 247 Lingam R, Hunt L, Golding J, Jongmans M, Emond A (2009) Prevalence of developmental coordination disorder using the DSM-IV at 7 years of age: a UK population-based study. *Pediatrics* 123: e693-e700.
- 248 Valentini NC, Olivera MA, Pangelinan MM, Whitall J, Clark JE (2017) Can the MABC discriminate and predict motor impairment? A comparison of Brazilian and North American children. *Int J Ther Rehabil* 24: 3.
- 249 Lindsay G, Strand S (2016) Children with language impairment: prevalence, associated difficulties, and ethnic disproportionality in an english population. *Front Educ* 1: 2.
- 250 Stevenson J, Richman N (1976) The prevalence of language delay in a population of three-year-old children and its association with general retardation. *Dev Med Child Neurol* 18: 431-441.
- 251 Tomblin JB, Records NL, Buckwalter P, Zhang X, Smith E, et al. (1997) Prevalence of specific language impairment in kindergarten children. *J Speech Lang Hear Res* 40: 1245-1260.
- 252 Lewis C, Hitch GJ, Walker P (1994) The prevalence of specific arithmetic difficulties and specific reading difficulties in 9- to 10-year-old boys and girls. *J Child Psychol Psychiatry* 35: 283-292.
- 253 Badian NA (1983) Dyscalculia and nonverbal disorders of learning. In: *Progress in learning disabilities* (Myklebust HR Edn). New York: Ulverscroft 5: 235-264.

- 254 Badian NA (1999) Persistent arithmetic, reading, or arithmetic and reading disability. *Ann Dyslexia* 49: 43-70.
- 255 Rutter M, Tizard J, Yule W, Graham P, Whitmore K (1976) Isle of wight studies, 1964-1974. *Psychol Med* 6: 313-332.
- 256 Yule W, Rutter M, Berger M, Thompson J (1974) Over- and under-achievement in reading: distribution in the general population. *Br J Educ Psychol* 44: 1-12.
- 257 Lindgren SD, De Renzi E, Riehman LC (1985) Cross-national comparisons of developmental dyslexia in Italy and the United States. *Child Dev* 56: 1404-1417.
- 258 Shaywitz SE, Shaywitz BA, Fletcher JM, Escobar MD (1990) Prevalence of reading disability in boys and girls: results of the connecticut longitudinal study. *J Am Med Assoc* 264: 998-1002.
- 259 Roeleveld N, Zielhuis GA (1997) The prevalence of mental retardation: a critical review of recent literature. *Dev Med Child Neurol* 39: 125-132.
- 260 Simonoff E, Pickles A, Chadwick O, Gringras P, Wood N, et al. (2006) The Croydon assessment of learning study: prevalence and educational identification of mild mental retardation. *J Child Psychol Psychiatry Allied Discip* 47: 828-839.
- 261 <https://www.gov.uk/government/statistics/special-educational-needs-in-england-january-2017>
- 262 Kraus L (2017) 2016 disability statistics annual report. Durham, USA.
- 263 Croen LA, Grether JK, Selvin S (2001) The epidemiology of mental retardation of unknown cause. *Pediatrics* 107: e86-e86.
- 264 CDC (1996) State-specific rates of mental retardation - United States, 1993. *Morb Mortal Wkly Rep* 45: 61-65.
- 265 Bhasin TK, Brocksen S, Avchen RN, Van Naarden Braun K (2006) Prevalence of four developmental disabilities among children aged 8 years - Metropolitan Atlanta Developmental Disabilities Surveillance Program, 1996 and 2000. *Morb Mortal Wkly Rep* 55: 1-9.
- 266 Boyle CA, Yeargin-Allsopp M, Doernberg NS, Holmgreen P, Murphy CC, et al. (1996) Prevalence of selected developmental disabilities in children 3-10 years of age: the Metropolitan Atlanta Developmental Disabilities Surveillance Program, 1991. *Morb Mortal Wkly Rep* 45: 1-14.
- 267 Van Naarden Braun K, Christensen D, Doernberg N, Schieve L, Rice C, et al. (2015) Trends in the prevalence of autism spectrum disorder, cerebral palsy, hearing loss, intellectual disability, and vision impairment, metropolitan Atlanta, 1991-2010. *PLoS One* 10: 1-21.
- 268 Scharf JM, Miller LL, Mathews CA, Ben-Shlomo Y (2012) Prevalence of tourette syndrome and chronic tics in the population-based avon longitudinal study of parents and children cohort. *J Am Acad Child Adolesc Psychiatry* 51: 192-201.
- 269 Hornsey H, Banerjee S, Zeitlin H, Robertson M (2001) The prevalence of tourette syndrome in 13-14-year-olds in mainstream schools. *J Child Psychol Psychiatry* 42: 1035-1039.
- 270 Mason A, Banerjee S, Eapen V, Zeitlin H, Robertson MM (1998) The prevalence of tourette syndrome in a mainstream school population. *Dev Med Child Neurol* 40: 292-296.
- 271 CDC (2009) Prevalence of diagnosed tourette syndrome in persons aged 6-17 Years --- United States, 2007. *Morb Mortal Wkly Rep* 58: 581-585.
- 272 Comings DE, Himes JA, Comings BG (1990) An epidemiologic study of tourette's syndrome in a single school district. *J Clin Psychiatry* 51: 463-469.
- 273 Gadow KD, Nolan EE, Sprafkin J, Schwartz J (2002) Tics and psychiatric comorbidity in children and adolescents. *Dev Med Child Neurol* 44: 330-338.
- 274 Kurlan R, McDermott MP, Deeley C, Como PG, Brower C, et al. (2001) Prevalence of tics in schoolchildren and association with placement in special education. *Neurology* 57: 1383-1388.
- 275 Snider LA, Seligman LD, Ketchen BR, Levitt SJ, Bates LR, et al. (2002) Tics and problem behaviors in schoolchildren: Prevalence, characterization, and associations. *Pediatrics* 110: 331-336.
- 276 Peterson BS, Pine DS, Cohen P, Brook JS (2001) Prospective, longitudinal study of tic, obsessive-compulsive, and attention-deficit/hyperactivity disorders in an epidemiological sample. *J Am Acad Child Adolesc Psychiatry* 40: 685-695.
- 277 Burd L, Kerbeshian J, Wikenheiser M, Fisher W (1986) A prevalence study of gilles de la tourette syndrome in North Dakota school-age children. *J Am Acad Child Psychiatry* 25: 552-553.
- 278 Sinzig J, Walter D, Doepfner M (2009) Attention deficit/hyperactivity disorder in children and adolescents with autism spectrum disorder: symptom or syndrome? *J Atten Disord* 13: 117-126.
- 279 Yoshida Y, Uchiyama T (2004) The clinical necessity for assessing attention deficit/hyperactivity disorder (AD/HD) symptoms in children with high-functioning pervasive developmental disorder (PDD). *Eur Child Adolesc Psychiatry* 13: 307-314.
- 280 Charman T, Ricketts J, Dockrell JE, Lindsay G, Palikara O (2015) Emotional and behavioural problems in children with language impairments and children with autism spectrum disorders. *Int J Lang Commun Disord* 50: 84-93.
- 281 De Bruin EI, Ferdinand EI, Meester S, De Nijs PFA, Verheij F (2007) High rates of psychiatric co-morbidity in PDD-NOS. *J Autism Dev Disord* 37: 877-886.
- 282 Gadow KD, Devincent CJ, Pomeroy J, Azizian A (2004) Psychiatric symptoms in preschool children with PDD and clinic and comparison samples. *J Autism Dev Disord* 34: 379-393.
- 283 Ghaziuddin M, Weidmer-Mikhail E, Ghaziuddin N (1998) Comorbidity of asperger syndrome: a preliminary report. *J Intellect Disabil Res* 42: 279-283.
- 284 Goldstein S, Schwebach AJ (2004) The comorbidity of pervasive developmental disorder and attention deficit hyperactivity disorder: results of a retrospective chart review. *J Autism Dev Disord* 34: 329-339.
- 285 Lee DO, Ousley OY (2006) Attention-deficit hyperactivity disorder symptoms in a clinic sample of children and adolescents with pervasive developmental disorders. *J Child Adolesc Psychopharmacol* 16: 737-746.
- 286 Leyfer OT, Folstein SE, Bacalman S, Davis NO, Dinh E, et al. (2006) Comorbid psychiatric disorders in children with autism: interview development and rates of disorders. *J Autism Dev Disord* 36: 849-861.
- 287 Mpaka DM, Okitundu DLEA, Ndjukendi AO, N'situ AM, Kinsala SY, et al. (2016) Prevalence and comorbidities of autism among children referred to the outpatient clinics for neurodevelopmental disorders. *Pan Afr Med J* 25: 82.

- 288 Green D, Charman T, Pickles A, Chandler S, Loucas T, et al. (2009) Impairment in movement skills of children with autistic spectrum disorders. *Dev Med Child Neurol* 51: 311-316.
- 289 Kopp S, Beckung E, Gillberg C (2010) Developmental coordination disorder and other motor control problems in girls with autism spectrum disorder and/or attention-deficit/hyperactivity disorder. *Res Dev Disabil* 31: 350-361.
- 290 Pieters S, De Block K, Scheiris J, Eyssen M, Desoete A, et al. (2012) How common are motor problems in children with a developmental disorder: rule or exception? *Child Care Health Dev* 38: 139-145.
- 291 Sumner E, Leonard HC, Hill EL (2016) Overlapping phenotypes in autism spectrum disorder and developmental coordination disorder: a cross-syndrome comparison of motor and social skills. *J Autism Dev Disord* 46: 2609-2620.
- 292 Miyahara M, Tsujii M, Hori M, Nakanishi K, Kageyama H, et al. (1997) Brief report: motor incoordination in children with asperger syndrome and learning disabilities. *J Autism Dev Disord* 27: 595-603.
- 293 Bartak L, Rutter M, Cox A (1977) A comparative study of infantile autism and specific developmental receptive language disorder. III. Discriminant function analysis. *J Autism Child Schizophr* 7: 383-396.
- 294 Delobel-Ayoub M, Ehlinger V, Klapouszczak D, Maffre T, Raynaud JP, et al. (2015) Socioeconomic disparities and prevalence of autism spectrum disorders and intellectual disability. *PLoS One* 10: 1-13.
- 295 Idring S, Lundberg M, Sturm H, Dalman C, Gumpert C, et al. (2015) Changes in prevalence of autism spectrum disorders in 2001-2011: findings from the Stockholm youth cohort. *J Autism Dev Disord* 45: 1766-1773.
- 296 Kim YS, Leventhal BL, Koh YJ, Fombonne E, Laska E, et al. (2011) Prevalence of autism spectrum disorders in a total population sample. *Am J Psychiatry* 168: 904-912.
- 297 Williams E, Thomas K, Sidebotham H, Emond A (2008) Prevalence and characteristics of autistic spectrum disorders in the ALSPAC cohort. *Dev Med Child Neurol* 50: 672-677.
- 298 Kalyva E, Kyriazi M, Vargiami E, Zafeiriou DI (2016) A review of co-occurrence of autism spectrum disorder and tourette syndrome. *Res Autism Spectr Disord* 24: 39-51.
- 299 Canitano R, Vivanti G (2007) Tics and tourette syndrome in autism spectrum disorders. *Autism* 11: 19-28.
- 300 Larson K, Russ SA, Kahn RS, Halfon N (2011) Patterns of comorbidity, functioning, and service use for US children with ADHD, 2007. *Pediatrics* 127: 462-470.
- 301 Kadesjö B, Gillberg C (1998) Attention deficits and clumsiness in Swedish 7-year-old children. *Dev Med Child Neurol* 40: 796-804.
- 302 Kadesjö B, Gillberg C (2001) The comorbidity of ADHD in the general population of Swedish school-age children. *J Child Psychol Psychiatry Allied Discip* 42: 487-492.
- 303 Piek JP, Pitcher TM, Hay DA (1999) Motor coordination and kinaesthesia in boys with attention-deficit-hyperactivity disorder. *Dev Med Child Neurol* 41: 159-165.
- 304 Pitcher TM, Piek JP, Hay DA (2003) Fine and gross motor ability in males with ADHD. *Dev Med Child Neurol* 45: 525-535.
- 305 Tsui KW, Lai KYC, Lee MMC, Shea CKS, Tong LCT (2016) Prevalence of motor problems in children with attention deficit hyperactivity disorder in Hong Kong. *Hong Kong Med J* 22: 98-105.
- 306 Steinhausen HC, Nøvik TS, Balursson G, Curatolo P, Lorenzo MJ, et al. (2006) Co-existing psychiatric problems in ADHD in the ADORE cohort. *Eur Child Adolesc Psychiatry* 15: 25-30.
- 307 Spencer T, Biederman J, Harding M, Donnell DO, Wilens T, et al. (1998) Disentangling the overlap between Tourette's disorder and ADHD. *J Child Psychol Psychiatry Allied Discip* 39: 1037-1044.
- 308 Czamara D, Tiesler CMT, Kohlböck G, Berdel D, Hoffmann B, et al. (2013) Children with ADHD symptoms have a higher risk for reading, spelling and math difficulties in the GINIplus and LISAPlus cohort studies. *PLoS One* 8: e63859.
- 309 Capano L, Minden D, Chen SX, Schachar RJ, Ickowicz A (2008) Mathematical learning disorder in school-age children with attention-deficit hyperactivity disorder. *La Rev Can Psychiatr* 53: 392-399.
- 310 Monuteaux MC, Faraone SV, Herzig K, Navsaria N, Biederman J (2005) ADHD and dyscalculia. *J Learn Disabil* 38: 86-93.
- 311 Germanò E, Gagliano A (2010) Comorbidity of ADHD and dyslexia. *Dev Neuropsychol* 35: 475-493.
- 312 Willcutt EG, Pennington BF, Olson RK, DeFries JC (2007) Understanding comorbidity: a twin study of reading disability and attention-deficit/hyperactivity disorder. *Am J Med Genet Part B Neuropsychiatr Genet* 144: 709-714.
- 313 Pauc R (2005) Comorbidity of dyslexia, dyspraxia, attention deficit disorder (ADD), attention deficit hyperactive disorder (ADHD), obsessive compulsive disorder (OCD) and Tourette's syndrome in children: a prospective epidemiological study. *Clin Chiropr* 8: 189-198.
- 314 Jacob P, Srinath S, Girimaji S, Seshadri S, Sagar J, et al. (2016) Comorbidity in attention-deficit hyperactivity disorder: a clinical study from India. *East Asian Arch Psychiatry* 26: 148-53.
- 315 Ahuja A, Martin J, Langley K, Thapar A (2013) Intellectual disability in children with attention deficit hyperactivity disorder. *J Pediatr* 163: 890-895.
- 316 Xenitidis K, Paliokosta E, Rose E, Maltezos S, Bramham J (2010) ADHD symptom presentation and trajectory in adults with borderline and mild intellectual disability. *J Intellect Disabil Res* 54: 668-677.
- 317 Lingam R, Golding J, Jongmans MJ, Hunt LP, Ellis M, et al. (2010) The association between developmental coordination disorder and other developmental traits. *Pediatrics* 126: e1109-e1118.
- 318 Kadesjö B, Gillberg C (1999) Developmental coordination disorder in Swedish 7-year-old children. *J Am Acad Child Adolesc Psychiatry* 38: 820-828.
- 319 Archibald LMD, Alloway TP (2008) Comparing language profiles: children with specific language impairment and developmental coordination disorder. *Int J Lang Commun Disord* 43: 165-180.
- 320 Alloway TP (2007) Working memory, reading, and mathematical skills in children with developmental coordination disorder. *J Exp Child Psychol* 96: 20-36.
- 321 Pieters S, Roeyers H, Rosseel Y, Van Waelvelde H, Desoete A (2015) Identifying subtypes among children with developmental coordination disorder and mathematical learning disabilities, using model-based clustering. *J Learn Disabil* 48: 83-95.
- 322 Dockrell J, Lindsay G (2000) Meeting the needs of children with specific speech and language difficulties. *Eur J Spec Needs Educ* 15: 24-41.
- 323 Paul R, Cohen DJ, Caparulo BK (1983) A longitudinal study of

- patients with severe developmental disorders of language learning. *J Am Acad Child Psychiatry* 22: 525-534.
- 324 Westerlund M, Bergkvist L, Lagerberg D, Sundelin C (2002) Comorbidity in children with severe developmental language disability. *Acta Paediatr* 91: 529-534.
- 325 Beitchman JH, Brownlie EB, Inglis A, Wild J, Ferguson B, et al. (1996) Seven-year follow-up of speech/language impaired and control children: psychiatric outcome. *J Child Psychol Psychiatry* 37: 961-970.
- 326 Finlay JCS, McPhillips M (2013) Comorbid motor deficits in a clinical sample of children with specific language impairment. *Res Dev Disabil* 34: 2533-2542.
- 327 Gaines R, Missiuna C (2007) Early identification: are speech/language-impaired toddlers at increased risk for developmental coordination disorder?. *Child Care Health Dev* 33: 325-332.
- 328 Hill EL (1998) A dyspraxic deficit in specific language impairment and developmental coordination disorder? evidence from hand and arm movements. *Dev Med Child Neurol* 40: 388-395.
- 329 Rintala P, Pienimäki K, Ahonen T, Cantell M, Kooistra L (1998) The effects of a psychomotor training programme on motor skill development in children with developmental language disorders. *Hum Mov Sci* 17: 721-737.
- 330 Visscher C, Houwen S, Scherder EJAA, Moolenaar B, Hartman E (2007) Motor profile of children with developmental speech and language disorders. *Pediatrics* 120: e158-e163.
- 331 Webster RI, Majnemer, Platt RW, Shevell MI (2005) Motor function at school age in children with a preschool diagnosis of developmental language impairment. *J Pediatr* 146: 80-85.
- 332 Webster RI, Erdos C, Evans K, Majnemer A, Kehayia E, et al. (2006) The clinical spectrum of developmental language impairment in school-aged children: language, cognitive, and motor findings. *Pediatrics* 118: e1541-e1549.
- 333 Catts HW, Fey ME, Tomblin JB, Zhang X (2002) A longitudinal investigation of reading outcomes in children with language impairments. *J Speech Lang Hear Res* 45: 1142.
- 334 Dalby M, Elbro C, Mårbjerg S (2009) Langtidsprognosen for sprogindlæringsvanskeligheder. *Ugeskr læger* 1: 37-41.
- 335 McArthur GM, Hogben JH, Edwards VT, Heath SM, Mengler ED (2000) On the 'specifics' of specific reading disability and specific language impairment. *J Child Psychol Psychiatry* 41: 869-874.
- 336 Shalev RS, Manor O, Gross-Tsur V (1997) Neuropsychological aspects of developmental dyscalculia. *Math Cogn* 3: 105-120.
- 337 Pieters S, Desoete A, Roeyers H, Vanderswalmen R, Van Waelvelde H (2012) Behind mathematical learning disabilities: what about visual perception and motor skills?. *Learn Individ Differ* 22: 498-504.
- 338 Dirks E, Spyer G, Van Lieshout ECDM, De Sonnevile L (2008) Prevalence of combined reading and arithmetic disabilities. *J Learn Disabil* 41: 460-473.
- 339 Jordan NC, Hanich LB, Kaplan D (2003) A longitudinal study of mathematical competencies in children with specific mathematics difficulties versus children with comorbid mathematics and reading difficulties. *Child Dev* 74: 834-850.
- 340 Landerl K, Moll K (2010) Comorbidity of learning disorders: prevalence and familial transmission. *J Child Psychol Psychiatry Allied Discip* 51: 287-294.
- 341 Iversen S, Berg K, Ellertsen B, Tønnessen FE (2005) Motor coordination difficulties in a municipality group and in a clinical sample of poor readers. *Dyslexia* 11: 217-231.
- 342 David M, Dieterich K, Billette de Villemeur A, Jouk PS, Counillon J, et al. (2014) Prevalence and characteristics of children with mild intellectual disability in a French county. *J Intellect Disabil Res* 58: 591-602.
- 343 Oeseburg B, Jansen DEMC, Dijkstra GJ, Groothoff JW, Reijneveld SA (2010) Prevalence of chronic diseases in adolescents with intellectual disability. *Res Dev Disabil* 31: 698-704.
- 344 Lindblad I, Gillberg C, Fernell E (2011) ADHD and other associated developmental problems in children with mild mental retardation. The use of the 'Five-To-Fifteen' questionnaire in a population-based sample. *Res Dev Disabil* 32: 2805-2809.
- 345 Huisman-van Dijk H, Van De Schoot R, Rijkeboer MM, Mathews CA, Cath DC (2016) The relationship between tics, OC, ADHD and autism symptoms: a cross-disorder symptom analysis in Gilles de la Tourette syndrome patients and their family members. *Psychiatry Res* 237: 138-146.
- 346 Baron-Cohen S, Scahill VL, Izaguirre J, Hornsey H, Robertson MM (1999) The prevalence of Gilles de la Tourette syndrome in children and adolescents with autism: a large scale study. *Psychol Med* 29: 1151-1159.
- 347 Burd L, Li Q, Kerbeshian J, Klug MG, Freeman RD (2009) Tourette Syndrome and Comorbid Pervasive Developmental Disorders. *J Child Neurol* 24 : 170-175.
- 348 Ghanizadeh A, Mosallaei S (2009) Psychiatric disorders and behavioral problems in children and adolescents with Tourette syndrome. *Brain Dev* 31: 15-19.
- 349 Freeman RD, Aabech H, Allagband-Rad J, Berlin C, Bruun R, et al. (2007) Tic disorders and ADHD: Answers from a world-wide clinical dataset on Tourette syndrome. *Eur Child Adolesc Psychiatry* 16: 15-23.
- 350 Kadesjö B, Gillberg C (2000) Tourette's disorder: epidemiology and comorbidity in primary school children. *J Am Acad Child Adolesc Psychiatry* 39: 548-555.
- 351 Tabori Kraft J, Dalsgaard S, Obel C, Thomsen PH, Henriksen TB, et al. (2012) Prevalence and clinical correlates of tic disorders in a community sample of school-age children. *Eur Child Adolesc Psychiatry* 21: 5-13.
- 352 Comings DE, Comings BG (1987) A controlled study of Tourette syndrome. I. Attention-deficit disorder, learning disorders, and school problems. *Am J Hum Genet* 41: 701-741.
- 353 Kano Y, Ohta M, Nagai Y, Scahill L (2010) Association between Tourette syndrome and comorbidities in Japan. *Brain Dev* 32: 201-207.
- 354 Comings DE, Comings BG (1984) Tourette's syndrome and attention deficit disorder with hyper activity: are they genetically related?. *J Am Acad Child Adolesc Psychiatry* 23: 138-146.
- 355 Burd L, Freeman RD, Klug MG, Kerbeshian J (2005) Tourette syndrome and learning disabilities. *BMC Pediatr* 5: 7-12.
- 356 Butwicki A, Långström N, Larsson H, Lundström S, Serlachius E, et al. (2017) Increased risk for substance use-related problems in autism spectrum disorders: a population-based cohort study. *J Autism Dev Disord* 47: 80-89.
- 357 Glidden D, Bouman WP, Jones BA (2016) Gender dysphoria and

- autism spectrum disorder : a systematic review of the literature. *Sex Med Rev* 4: 3-14.
- 358 Marín JL, Rodríguez-Franco MA, Chugani VM, Maganto MM, Villoria ED, et al. (2018) Prevalence of schizophrenia spectrum disorders in average-IQ adults with autism spectrum disorders: a meta-analysis. *J Autism Dev Disord* 48: 239-250.
- 359 Stahlberg O, Soderstrom H, Rastam M, Gillberg C (2004) Bipolar disorder, schizophrenia, and other psychotic disorders in adults with childhood onset AD/HD and/or autism spectrum disorders. *J Neural Transm* 111: 891-902.
- 360 Westwood H, Tchanturia K (2017) Autism spectrum disorder in anorexia nervosa: an updated literature review. *Curr Psychiatry Rep* 19: 41.
- 361 Bolton PF, Carcani-Rathwell I, Hutton J, Goode S, Howlin P, et al. (2011) Epilepsy in autism: Features and correlates. *Br J Psychiatry* 198: 289-294.
- 362 Wentz E, Björk A, Dahlgren J (2017) Neurodevelopmental disorders are highly over-represented in children with obesity: a cross-sectional study. *Obesity* 25: 178-184.
- 363 Zuckerman KE, Hill AP, Guion K, Voltolina L, Fombonne E (2014) Overweight and obesity: prevalence and correlates in a large clinical sample of children with autism spectrum disorder. *J Autism Dev Disord* 44: 1708-1719.
- 364 Baeza-Velasco C, Pailhez G, Bulbena A, Baghdadli A (2015) Joint hypermobility and the heritable disorders of connective tissue: clinical and empirical evidence of links with psychiatry. *Gen Hosp Psychiatry* 37: 24-30.
- 365 Broder-Fingert S, Brazauskas K, Lindgren K, Iannuzzi D, Van Cleave J (2014) Prevalence of overweight and obesity in a large clinical sample of children with autism. *Acad Pediatr* 14: 408-414.
- 366 Doshi-Velez F, Ge Y, Kohane I (2014) Comorbidity clusters in autism spectrum disorders: an electronic health record time-series analysis. *Pediatrics* 133: e54-e63.
- 367 Frye RE, Rossignol DA (2016) Identification and treatment of pathophysiological comorbidities of autism spectrum disorder to achieve optimal outcomes. *Clin Med Insights Pediatr* 10: S38337.
- 368 Giovanardi Rossi P, Posar A, Parmeggiani A (2000) Epilepsy in adolescents and young adults with autistic disorder. *Brain Dev* 22: 102-106.
- 369 Gurney JG, McPheeters ML, Davis MM (2006) Parental report of health conditions and health care use among children with and without autism. *Arch Pediatr Adolesc Med* 160: 825-830.
- 370 Konstantareas MM, Homatidis S (1987) Brief report: ear infections in autistic and normal children. *J Autism Dev Disord* 17: 585-594.
- 371 Rosenhall U, Nordin V, Sandström M, Ahlsén G, Gillberg C (1999) Autism and hearing loss. *J Autism Dev Disord* 29: 349-357.
- 372 Suren P, Bakken IJ, Aase H, Chin R, Gunnes N, et al. (2012) Autism spectrum disorder, ADHD, epilepsy, and cerebral palsy in Norwegian children. *Pediatrics* 130: e152-e158.
- 373 Farbstein I, Mansbach-Kleinfeld I, Auerbach JG, Ponizovsky AM, Apter A (2014) The Israel survey of mental health among adolescents: prevalence of attention-deficit/hyperactivity disorder, comorbidity, methylphenidate use, and help-seeking patterns. *Isr Med Assoc J* 16: 568-573.
- 374 Gillberg C, Gillberg IC, Rasmussen P, Kadesjö B, Söderström H, et al. (2004) Co-existing disorders in ADHD - Implications for diagnosis and intervention. *Eur Child Adolesc Psychiatry* 13: 80-93.
- 375 Sprafkin J, Gadow KD, Weiss MD, Schneider J, Nolan EE (2007) Psychiatric comorbidity in ADHD symptom subtypes in clinic and community adults. *J Atten Disord* 11: 114-124.
- 376 Strang JF, Kenworthy L, Dominska A, Sokoloff J, Kenealy LE, et al. (2014) Increased gender variance in autism spectrum disorders and attention deficit hyperactivity disorder. *Arch Sex Behav* 43: 1525-1533.
- 377 Jameson ND, Sheppard BK, Lateef TM, Vande Voort JL, He JP, et al. (2016) Medical comorbidity of attention-deficit/hyperactivity disorder in US adolescents. *J Child Neurol* 31: 1282-1289.
- 378 Virring A, Lambek R, Thomsen PH, Møller LR, Jennum PJ (2016) Disturbed sleep in attention-deficit hyperactivity disorder (ADHD) is not a question of psychiatric comorbidity or ADHD presentation. *J Sleep Res* 25: 333-340.
- 379 Lingam R, Jongmans MJ, Ellis M, Hunt LP, Golding J, et al. (2012) Mental health difficulties in children with developmental coordination disorder. *Pediatrics* 129: e882-e891.
- 380 Adib N, Davies K, Grahame R, Woo P, Murray KJ (2005) Joint hypermobility syndrome in childhood. A not so benign multisystem disorder? *Rheumatology* 44: 744-750.
- 381 Barnett AL, Wiggs L (2012) Sleep behaviour in children with developmental co-ordination disorder. *Child Care Health Dev* 38: 403-411.
- 382 Kirby A, Davies R (2007) Developmental coordination disorder and joint hypermobility syndrome - overlapping disorders? Implications for research and clinical practice. *Child Care Health Dev* 33: 513-519.
- 383 Wiggs L, Sparrowhawk M, Barnett AL (2016) Parent report and actigraphically defined sleep in children with and without developmental coordination disorder; links with fatigue and sleepiness. *Front Pediatr* 4: 81.
- 384 Clegg J, Hollis C, Mawhood L, Rutter M (2005) Developmental language disorders - a follow-up in later adult life. Cognitive, language and psychosocial outcomes. *J Child Psychol Psychiatry Allied Discip* 46: 128-149.
- 385 Mouridsen SE, Hauschild KM (2009) A longitudinal study of personality disorders in individuals with and without a history of developmental language disorder. *Logoped Phoniatr Vocol* 34: 135-41.
- 386 Whitehouse AJO, Watt HJ, Line EA, Bishop DVM (2009) Adult psychosocial outcomes of children with specific language impairment, pragmatic language impairment and autism. *Int J Lang Commun Disord* 44: 511-528.
- 387 Selassie GRH, Jennische M, Kyllerman M, Viggedal G, Hartelius L (2005) Comorbidity in severe developmental language disorders: neuropsychiatric and psychological considerations. *Acta Paediatr* 94: 471-478.
- 388 Cederlöf M, Östberg P, Pettersson E, Anckarsäter H, Gumpert C, et al. (2014) Language and mathematical problems as precursors of psychotic-like experiences and juvenile mania symptoms. *Psychol Med* 44: 1293-1302.
- 389 Crow TJ, Done DJ, Sacker A (1995) Childhood precursors of psychosis as clues to its evolutionary origins. *Eur Arch Psychiatry Clin Neurosci* 245: 61-69.
- 390 Willcutt EG, Petrill SA, Wu S, Boada R, DeFries JC, et al. (2013) Comorbidity between reading disability and math disability:

- concurrent psychopathology, functional impairment, and neuropsychological functioning. *J Learn Disabil* 46: 500-516.
- 391 Thomé U, da P Alves SR, Guerreiro SM, da Costa CRCM, de S Moreira F, et al. (2014) Developmental dyscalculia in children and adolescents with idiopathic epilepsies in a Brazilian sample. *Arq Neuropsiquiatr* 72: 283-288.
- 392 Rössler W, Vetter S, Müller M, Gallo WT, Haker H, et al. (2011) Risk factors at the low end of the psychosis continuum: much the same as at the upper end? *Psychiatry Res* 189: 77-81.
- 393 Törő KT, Miklósi M, Horanyi E, Kovács GP, Balázs J (2018) Reading disability spectrum: early and late recognition, subthreshold, and full comorbidity. *J Learn Disabil* 51: 158-167.
- 394 Emerson E (2003) Prevalence of psychiatric disorders in children and adolescents with and without intellectual disability. *J Intellect Disabil Res* 47: 51-58.
- 395 Deb S, Thomas M, Bright C (2001) Mental disorder in adults with intellectual disability. I: prevalence of functional psychiatric illness among a community-based population aged between 16 and 64 years. *J Intellect Disabil Res* 45: 495-505.
- 396 a Bradley E, a Summers J, Wood HL, Bryson SE (2004) Comparing rates of psychiatric and behavior disorders in adolescents and young adults with severe intellectual disability with and without autism. *J Autism Dev Disord* 34: 151-161.
- 397 Morgan VA, Leonard H, Bourke J, Jablensky A (2008) Intellectual disability co-occurring with schizophrenia and other psychiatric illness: population-based study. *Br J Psychiatry* 193: 364-372.
- 398 Myrbakk E, von Tetzchner S (2008) Psychiatric disorders and behavior problems in people with intellectual disability. *Res Dev Disabil* 29: 316-332.
- 399 Cooper S, Smiley E, Morrison J, Williamson A, Allan L (2007) Mental ill-health in adults with intellectual disabilities: prevalence and associated factors. *Br J Psychiatry* 190: 27-35.
- 400 White P, Chant D, Edwards N, Townsend C, Waghorn G (2005) Prevalence of intellectual disability and comorbid mental illness in an Australian community sample. *Aust N Z J Psychiatry* 39: 395-400.
- 401 Scuticchio D, Rossi M, Bertelli MO (2017) Personality disorders in intellectual disability: a prevalence study. *J Ment Health Res Intellect Disabil* 10: 181.
- 402 Carroll Chapman SL, Wu LT (2012) Substance abuse among individuals with intellectual disabilities. *Res Dev Disabil* 33: 1147-1156.
- 403 Coffey BJ, Biederman J, Smoller JW, Geller DA, Sarin P, et al. (2000) Anxiety disorders and tic severity in juveniles with tourette's disorder. *J Am Acad Child Adolesc Psychiatry* 39: 562-568.
- 404 Cavanna AE, Robertson MM, Critchley HD (2007) Schizotypal personality traits in Gilles de la tourette syndrome. *Acta Neurol Scand* 116: 385-391.
- 405 Robertson MM, Banerjee S, Hiley PJ, Tannock C (1997) Personality disorder and psychopathology in tourette's syndrome: a controlled study. *Br J Psychiatry* 171: 283-286.
- 406 Chen MH, Su TP, Chen YS, Hsu JW, Huang KL, et al. (2013) Attention deficit hyperactivity disorder, tic disorder, and allergy: is there a link? A nationwide population-based study. *J Child Psychol Psychiatry Allied Discip* 54: 545-551.
- 407 Hung KL (2017) Association of epilepsy and tourette syndrome in children. *Epilepsy J* 3: 1-2.
- 408 Tick B, Bolton P, Happé F, Rutter M, Rijdsdijk F (2016) Heritability of autism spectrum disorders: a meta-analysis of twin studies. *J Child Psychol Psychiatry Allied Discip* 57: 585-595.
- 409 Bertrand J, Mars A, Boyle CA, Bove F, Yeargin-Allsopp M, et al. (2001) Prevalence of autism in a United States population - The Brick Township, New Jersey, investigation. *Pediatrics* 108: 1155-1161.
- 410 Lichtenstein P, Carlström E, Råstam M, Gillberg C, Anckarsäter H (2010) The genetics of autism spectrum disorders and related neuropsychiatric disorders in childhood. *Am J Psychiatry* 167: 1357-1363.
- 411 Bishop DVM, Hayiou-Thomas ME (2008) Heritability of specific language impairment depends on diagnostic criteria. *Genes Brain Behav* 7: 365-372.
- 412 Alarcón M, DeFries JC, Light JG, Pennington BF (1997) A twin study of mathematics disability. *J Learn Disabil* 30: 617-623.
- 413 Kovas Y, Haworth CMA, Harlaar N, Petrill SA, Plomin R (2007) Overlap and specificity of genetic and environmental influences on reading and mathematics disability in 10 year-old twins. *J Child Psychol Psychiatry* 48: 914-922.
- 414 Astrom RL, Wadsworth SJ, DeFries JC (2007) Etiology of the stability of reading difficulties: the longitudinal twin study of reading disabilities. *Twin Res Hum Genet* 10: 434-439.
- 415 Hawke JL, Wadsworth SJ, DeFries JC (2006) Genetic influences on reading difficulties in boys and girls: The Colorado twin study. *Dyslexia* 12: 21-29.
- 416 LaBuda MC, DeFries JC (1988) Genetic and environmental etiologies of reading disability: a twin study. *Ann Dyslexia* 38: 131-138.
- 417 Stevenson J, Graham P, Fredman G, McLoughlin V (1987) A twin study of genetic influences on reading and spelling ability and disability. *J Child Psychol Psychiatry* 28: 229-247.
- 418 Zilhão NR, Olthof MC, Smit DJA, Cath DC, Ligthart L, et al. (2017) Heritability of tic disorders: a twin-family study. *Psychol Med* 47: 1085-1096.
- 419 Mayhew AJ, Meyre D (2017) Assessing the heritability of complex traits in humans: methodological challenges and opportunities. *Curr Genomics* 18: 332-340.